

MATH 1271 Spring 2014, Midterm #2
Handout date: Thursday 17 April 2014
Instructor: Scot Adams

PRINT YOUR NAME:

PRINT YOUR X.500 ID:

PRINT YOUR TA'S NAME:

WHAT RECITATION SECTION ARE YOU IN?

Closed book, closed notes, no calculators/PDAs; no reference materials of any kind. Turn off all handheld devices, including cell phones.

Show work; a correct answer, by itself, may be insufficient for credit. Arithmetic need not be simplified, unless the problem requests it.

I. Multiple choice

A. (5 pts) (no partial credit) Let $f(x) = \sin^2(5x^4 + 1)$. Compute $\int_2^2 f(x) dx$. Circle one of the following answers:

- (a) 0
 - (b) 2
 - (c) 6
 - (d) 20
 - (e) NONE OF THE ABOVE
-

B. (5 pts) (no partial credit) Let $f(x) = e^{3x-4}$. Recall that $M_2S_1^5 f$ denotes the midpoint Riemann sum, from 1 to 5, of f , with two subintervals. Which of these is equal to $M_2S_1^5 f$? Circle one of the following answers:

- (a) $e^5 + e^{11}$
 - (b) $e^2 + e^8$
 - (c) $2(e^5 + e^{11})$
 - (d) $2(e^2 + e^8)$
 - (e) NONE OF THE ABOVE
-

C. (5 pts) (no partial credit) Find the derivative of $(2 + x^4)^{\cos x}$ w.r.t. x . Circle one of the following answers:

- (a) $[(2 + x^4)^{\cos x}][(\cos x)(\ln(2 + x^4)) + (-\sin x)(4x^3/(2 + x^4))]$
- (b) $[(2 + x^4)^{\cos x}][(-\sin x)(4x^3/(2 + x^4))]$
- (c) $[(2 + x^4)^{\cos x}][(-\sin x)(\ln(2 + x^4)) + (\cos x)(4x^3/(2 + x^4))]$
- (d) $[(2 + x^4)^{\cos x}][(\cos x)(\ln(2 + x^4))]$
- (e) NONE OF THE ABOVE

D. (5 pts) (no partial credit) Suppose $f''(x) = -(x - 7)^3(x - 8)^4$. At most one of the following statements is true. If one is, circle it. Otherwise, circle "NONE OF THE ABOVE".

- (a) f is concave down on $(-\infty, 7]$ and up on $[7, \infty)$.
 - (b) f is concave up on $(-\infty, 7]$ and down on $[7, \infty)$.
 - (c) f is concave up on $(-\infty, 7]$, down on $[7, 8]$ and up on $[8, \infty)$.
 - (d) f is concave down on $(-\infty, 7]$, up on $[7, 8]$ and down on $[8, \infty)$.
 - (e) NONE OF THE ABOVE
-

E. (5 pts) (no partial credit) Let $f(x) = e^{2x} + 3x$. What is the iterative formula of Newton's method used to solve $f(x) = 0$? Circle one of the following answers:

- (a) $x_{n+1} = x_n + \frac{e^{2x_n} + 3}{e^{2x_n} + 3x_n}$
 - (b) $x_{n+1} = x_n + \frac{2e^{2x_n} + 3}{e^{2x_n} + 3x_n}$
 - (c) $x_{n+1} = x_n + \frac{e^{2x_n} + 3x_n}{e^{2x_n} + 3}$
 - (d) $x_{n+1} = x_n + \frac{e^{2x_n} + 3x_n}{2e^{2x_n} + 3}$
 - (e) NONE OF THE ABOVE
-

F. (5 pts) (no partial credit) Let $y = x^2 + x$. Compute dy , evaluated at $x = 10$, $dx = 0.1$. Circle one of the following answers:

- (a) 1.2
- (b) 2.1
- (c) 1.22
- (d) 2.11
- (e) NONE OF THE ABOVE

II. True or false (no partial credit):

a. (5 pts) $\frac{d}{dx} \left[\int_1^x \sin(e^t) dt \right] = \cos(e^x)$.

b. (5 pts) Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be any function with a global maximum at 7. Assume that f'' is defined on \mathbb{R} . Then $f'(7) = 0$ and $f''(7) < 0$.

c. (5 pts) If f is continuous on $[a, b]$, then $\int_a^b (f(x)) dx = \lim_{n \rightarrow \infty} [M_n S_a^b f]$.

d. (5 pts) Let $f, g : \mathbb{R} \rightarrow \mathbb{R}$ be any two differentiable functions such that, for all $x \in \mathbb{R}$, $f'(x) = g'(x)$. Then $f - g$ is a constant.

e. (5 pts) Assume that $\lim_{x \rightarrow a} [f(x)] = 0 = \lim_{x \rightarrow a} [g(x)]$. Assume also that $\lim_{x \rightarrow a} \frac{f'(x)}{g'(x)} = -\infty$.

Then $\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = -\infty$.

THE BOTTOM OF THIS PAGE IS FOR TOTALING SCORES
PLEASE DO NOT WRITE BELOW THE LINE

VERSION A

I. A,B,C

I. D,E,F

II. a,b,c,d,e

III. 1.

III. 2.

III. 3.

III. 4.

III. Computations. Show work. Unless otherwise specified, answers must be exactly correct, but can be left in any form easily calculated on a standard calculator.

1. (10 pts) Find an antiderivative w.r.t. x of $\sin^2(2x - 1)$. (Hint: $\cos(2\theta) = 1 - 2(\sin^2 \theta)$.)

2. (10 pts) Let $f(x) = \int_x^{x^4} \sqrt{t^6 + 4t^2 + 4} dt$. Compute $f'(1)$.

3. (15 pts) We are asked to design a large cup in the shape of an inverted (*i.e.*, upside down) cone. The cup is to have an open top, and must contain $\pi/3$ cubic feet of volume inside. Let r be the radius of the top of the cup. On the interval $r > 0$, find the choice of r (in feet) that minimizes the surface area, A , of the cup. (HINT: Our local precalculus expert shows us the formula that relates A to r . It is $A = (\pi r)\sqrt{r^2 + r^{-4}}$.)

4. (10 pts) A square-based pyramid is growing. Its height is always equal to the length, s , of the sides of its base. Assume that its volume is always growing at a rate of 10 cubic feet per minute. Find the rate of growth in s (in feet per minute) at the moment when the volume is 9 cubic feet. (HINT: According to our local precalculus expert, its volume, V , is given by $V = s^3/3$.)