

Berry–Esseen bounds in the local limit theorems*

Sergey G. Bobkov^a and Friedrich Götze^{b,1} 

^a School of Mathematics, University of Minnesota, Minneapolis, MN, USA

^b Faculty of Mathematics, Bielefeld University, Germany

(e-mail: bobkov@math.umn.edu; goetze@math.uni-bielefeld.de)

Received June 26, 2024; revised November 28, 2024

Abstract. Berry–Esseen-type bounds are developed in the multidimensional local limit theorem in terms of the Lyapunov coefficients and maxima of involved densities.

MSC: 60E, 60F

Keywords: central limit theorem, local limit theorem, Berry–Esseen bounds

1 Introduction

Consider the normalized sum

$$Z_n = \frac{X_1 + \cdots + X_n}{\sqrt{n}}$$

of n independent random vectors X_k in \mathbb{R}^d with mean zero and covariance matrices $\sigma_k^2 I_d$ ($\sigma_k > 0$) such that $\sigma_1^2 + \cdots + \sigma_n^2 = n$. The latter ensures that Z_n has mean zero and identity covariance matrix I_d .

Introduce the normalized Lyapunov ratios of the third and fourth orders

$$\beta_3 = \sup_{|\theta|=1} \left[\frac{1}{n} \sum_{k=1}^n \mathbf{E} |\langle \theta, X_k \rangle|^3 \right], \quad \beta_4 = \sup_{|\theta|=1} \left[\frac{1}{n} \sum_{k=1}^n \mathbf{E} \langle \theta, X_k \rangle^4 \right],$$

assuming respectively that $\mathbf{E} |X_k|^3 < \infty$ and $\mathbf{E} |X_k|^4 < \infty$ for all $k \leq n$. In dimension one, the quantity β_3 is commonly used to quantify the normal approximation for the distribution of Z_n in a weak sense via the Berry–Esseen bound

$$\sup_{x \in \mathbb{R}} |\mathbf{P}\{Z_n \leq x\} - \mathbf{P}\{Z \leq x\}| \leq \frac{C}{\sqrt{n}} \beta_3,$$

* Research supported by the NSF grant DMS-2154001 and the GRF – SFB 1283/2 2021 – 317210226.

¹ Corresponding author.

where Z is a standard normal random variable, and C is an absolute constant (see, e.g., [18]). The Lyapunov ratio β_4 also appears in a natural way, for example, for the approximation of the characteristic function of Z_n by a corrected normal characteristic function. Both quantities may also be used to control the distance between the distribution of Z_n and Z on the real line in total variation and in relative entropy; see [9].

The aim of this note is to quantify the normal approximation in a stronger sense toward a uniform convergence of densities p_n of Z_n to the standard normal density

$$\varphi(x) = \frac{1}{(2\pi)^{d/2}} e^{-|x|^2/2}, \quad x \in \mathbb{R}^d.$$

In the i.i.d. situation (where all X_k are identically distributed with $\sigma_k = 1$), the necessary and sufficient condition for the convergence of the uniform distance

$$\Delta_n = \sup_x |p_n(x) - \varphi(x)|$$

to zero as $n \rightarrow \infty$ is that p_n is bounded for some $n = n_0$ (see [5, 18]). As typically $n_0 = 1$ in applications, it is natural to consider the case where all X_k have bounded densities.

So, introduce the maximum-of-density functional $M(X) = \text{ess sup}_x p(x)$, where p denotes the density of a random vector X , and define

$$M = \max_k M(X_k), \quad \sigma^2 = \max_k \sigma_k^2 \quad (\sigma > 0).$$

The functionals β_3 , β_4 , σ , and M can be used to derive the following upper bounds (which seem to be new already in the one-dimensional situation).

Theorem 1. *With some positive absolute constant C , the density p_n of Z_n satisfies*

$$\Delta_n \leq \frac{(C\sigma)^d M^2}{\sqrt{n}} \beta_3. \quad (1.1)$$

Moreover, if $\mathbf{E}\langle \theta, X_k \rangle^3 = 0$ for all $\theta \in \mathbb{R}^d$ and $k \leq n$ (in particular, if the distributions of X_k are symmetric), then

$$\Delta_n \leq \frac{(C\sigma)^{2d} M^3}{n} \beta_4. \quad (1.2)$$

Hence, when M and σ are bounded, it is possible to strengthen the Berry–Esseen theorem with an extension to higher dimensions.

To reflect the influence of $M(X_k)$ on average (similarly to β_3), rather than via the maximal value M , some refinements of the bounds (1.1)–(1.2) are given in Section 7.

For several classes of probability distributions, the functional $M(X)$ is of the order σ^{-d} when a random vector X has a covariance matrix $\sigma^d I_d$ (modulo d -dependent constants). Here is an example involving convexity properties of distributions.

Corollary 1. *Suppose that the random vectors X_k have log-concave densities, with mean zero and identity covariance matrix. Then*

$$\Delta_n \leq \frac{C_d}{\sqrt{n}} \quad (1.3)$$

with some constant C_d depending on d only. If additionally the distributions of X_k are symmetric, then

$$\Delta_n \leq \frac{C_d}{n}. \quad (1.4)$$

Based on the application of the Fourier transform (which is typical in the study of various variants of the central limit theorems), the main arguments used in the proof of Theorem 1 employ the subadditivity property of the maximum-of-density functional $M(X)$ with respect to convolutions. This tool has been introduced in the field of limit theorems only recently and is used in [10, 11]. Another ingredient in the proof involves the extension of the Statulevičius separation theorem for characteristic functions to higher dimensions.

The paper is organized as follows. We start with remarks on isotropic constants and general bounds for $M(X)$ in terms of the covariance matrix of a random vector X (Section 2) and then discuss the question of how we can separate the absolute value of a given characteristic function $f(t)$ from 1 in terms of $M(X)$ outside a neighborhood of the origin (Sections 3–4). In Sections 5, we recall the subadditivity property of $M(X)$ and develop its applications to the integrability properties of powers of characteristic functions. Basic results on the normal approximation of products of characteristic functions are recalled in Section 6. In Section 7, we derive refined bounds on Δ_n , which are used in Section 8 to finish the proofs of Theorem 1 and Corollary 1.

2 Lower bounds on maximum of density via covariance matrix

To start with, first, let us recall the general relation

$$M^2\sigma^2 \geq \frac{1}{12}, \tag{2.1}$$

which holds for any random variable X with standard deviation σ and maximum of density $M = M(X)$. As an early reference, we can mention the paper by Statulevičius [20, p. 651], where (2.1) is stated without derivation as an obvious fact. In the multidimensional situation, (2.1) is extended in the form

$$M^{2/d}\sigma^2 \geq \frac{1}{d+2}\omega_d^{-2/d}, \tag{2.2}$$

assuming that the random vector X has a covariance matrix $\sigma^2 I_d$ (such distributions are called isotropic). Here an equality is attained for the uniform distribution on every Euclidean ball in \mathbb{R}^d and, in particular, for the unit ball $B_2 = \{x \in \mathbb{R}^d: |x| \leq 1\}$ with volume

$$\omega_d = \text{vol}_d(B_2) = \frac{\pi^{d/2}}{\Gamma(\frac{d}{2} + 1)}. \tag{2.3}$$

This extremal property of balls has been investigated in convex geometry in the context of bounding volume of slices of convex bodies; see Hensley [16] and Ball [3]. More precisely, it was shown in [3, Lemma 6] that if the density of X satisfies $p(x) \leq p(0)$ for all $x \in \mathbb{R}^d$, then

$$p(0)^{2/d} \int_{\mathbb{R}^d} |x|^2 dx \geq \frac{d}{d+2}\omega_d^{-2/d}.$$

This amounts to (2.2) in the case where X has mean zero and covariance matrix $\sigma^2 I_d$, assuming that $p(x)$ is maximized at the origin. In [12, Prop. III.1], this inequality was generalized and strengthened as the property that for any nondecreasing function $H = H(t)$ in $t \geq 0$, the moment-type functional

$$\int_{\mathbb{R}^d} H(M^{1/d}|x|)p(x) dx$$

is minimized for the uniform distribution on every Euclidean ball with center at the origin. For $H(t) = t^2$, this property implies (2.2), assuming that X has mean zero and covariance matrix $\sigma^2 I_d$. Since inequality (2.2) is translation invariant, it continues to hold without the mean zero assumption.

The quantity $M^{1/d}\sigma$ is often called an isotropic constant of the distribution of X . It is also well known that relation (2.2) is in essence dimension-free, since it implies

$$M^{2/d}\sigma^2 \geq \frac{1}{2\pi e} \quad (2.4)$$

with an optimal constant on the right-hand side (attainable asymptotically for growing dimension d). More generally, if a random vector X has a covariance matrix R , then

$$(M^2 \det R)^{1/d} \geq \frac{1}{2\pi e}.$$

This follows from the previous lower bound (2.4) by applying it to the random vector $Y = R^{-1/2}X$. It is isotropic with $\sigma(Y) = 1$ and $M(Y) = \sqrt{\det R} M(X)$.

In this connection, let us mention that in the class of isotropic distributions on \mathbb{R}^d having log-concave densities, all these lower bounds can be reversed as

$$M^{2/d}\sigma^2 \leq K_d^2 \quad (2.5)$$

in terms of the maximal isotropic constant over this class for a fixed dimension d . In the equivalent form, the hyperplane conjecture raised in the 1980s by Bourgain, which is still open, states that K_d is bounded by an absolute constant. This is true, for example, when additionally the density of X is symmetric about all coordinate axes (see, e.g., [13]). As for the general log-concave case, at this moment the best result in this direction belongs to Klartag [17] with his bound $K_d^2 \leq C \log(d+1)$ for some absolute constant C . Let us refer the interested reader to [15] and [1] for the history of the problem and related results.

3 Separation of characteristic functions (Statulevičius theorem)

To apply the Fourier methods for the derivation of density bounds in a quantitative way, we have to realize how to separate the characteristic function

$$f(t) = \mathbf{E}e^{itX}, \quad t \in \mathbb{R},$$

of a random variable X from 1 outside a neighborhood of the origin (which is potentially possible due to the Riemann–Lebesgue lemma); that is, the task is to derive estimates like $\sup_{|t| \geq t_0} |f(t)| < 1 - \delta$ with an arbitrary $t_0 > 0$ and some positive $\delta = \delta(t_0)$. An important step in this direction was made by Statulevičius [20], who derived an upper bound, which we prefer to state in an equivalent form.

Proposition 1. *Given a random variable X with standard deviation σ and finite maximum of density $M = M(X)$, its characteristic function satisfies*

$$|f(t)| \leq 1 - \frac{c}{M^2\sigma^2} \min\{\sigma^2|t|^2, 1\}, \quad t \in \mathbb{R}, \quad (3.1)$$

with some absolute constant $c > 0$.

Some generalizations of this result to the general case of unbounded densities and without moment assumptions are discussed in [8]. In fact, Statulevičius considered more complicated quantities for upper bounds reflecting the behavior of the density p of X on nonoverlapping intervals of the real line and obtained the bound

$$|f(t)| \leq \exp\left\{-\frac{t^2}{96 M^2 (2\sigma|t| + \pi)^2}\right\}$$

as a partial case of a more general relation. Therefore let us include a shorter simplified argument aimed at the bound (3.1) only (without polishing the constant c).

Proof. In a more flexible form, inequality (2.1) can be equivalently written as

$$\operatorname{ess\,sup}_x q(x)^2 \int_{-\infty}^{\infty} x^2 q(x) \, dx \geq \frac{1}{12} \left(\int_{-\infty}^{\infty} q(x) \, dx \right)^3 \tag{3.2}$$

for any nonnegative Borel measurable function q on the real line.

Consider the symmetrized random variable $\tilde{X} = X - X'$, where X' is an independent copy of X . It has a nonnegative characteristic function $|f(t)|^2$ and density

$$w(x) = \int_{-\infty}^{\infty} p(x+y)p(y) \, dy \leq M.$$

Write

$$1 - |f(2\pi t)|^2 = \int_{-\infty}^{\infty} (1 - \cos(2\pi tx))w(x) \, dx = 2 \int_{-\infty}^{\infty} \sin^2(\pi tx) w(x) \, dx.$$

To bound the last integral from below, we may use the elementary inequality

$$\frac{1}{2} |\sin(\pi x)| \geq \|x\| = \min\{|x - k| : k \in \mathbb{Z}\},$$

where both sides represent 1-periodic functions. Assuming without loss of generality that $t > 0$ and using $1 - |f(s)|^2 = (1 - |f(s)|)(1 + |f(s)|) \leq 2(1 - |f(s)|)$, this gives

$$1 - |f(2\pi t)| \geq 4 \int_W \|tx\|^2 w(x) \, dx, \tag{3.3}$$

where we restrict the integration to the set $W = \{x \in \mathbb{R} : t|x| < N + 1/2\}$ for a suitable integer $N \geq 0$. Let us split the integral in (3.3) into the sets

$$W_k = \left\{ x \in \mathbb{R} : k - \frac{1}{2} < t|x| < k + \frac{1}{2} \right\}$$

and rewrite (3.3) as

$$1 - |f(2\pi t)| \geq 4 \sum_{k=-N}^N \int_{W_k} |tx - k|^2 w(x) \, dx = 4t^2 \sum_{k=-N}^N \int_{-1/(2t)}^{1/(2t)} y^2 w\left(y + \frac{k}{t}\right) \, dy.$$

Applying (3.2) to the functions $q_k(y) = w(y + k/t) \mathbf{1}_{[-1/(2t), 1/(2t)]}(y)$ and using $w \leq M$, we have

$$\int_{-1/(2t)}^{1/(2t)} y^2 w\left(y + \frac{k}{t}\right) \, dy \geq \frac{1}{12M^2} \left[\int_{-1/(2t)}^{1/(2t)} w\left(y + \frac{k}{t}\right) \, dy \right]^3 = \frac{1}{12M^2} \left(\int_{W_k} w(x) \, dx \right)^3.$$

Hence

$$1 - |f(2\pi t)| \geq \frac{t^2}{3M^2} \sum_{k=-N}^N Q_k^3 \quad \text{where } Q_k = \int_{W_k} w(x) dx.$$

Subject to $\sum_{k=-N}^N Q_k = Q$, the sum $\sum_{k=-N}^N Q_k^3$ is minimized when $Q_k = Q/(2N+1)$. This leads to

$$1 - |f(2\pi t)| \geq \frac{1}{3M^2} \frac{t^2}{(2N+1)^2} Q^3. \quad (3.4)$$

We should now maximize the right-hand side or choose a suitable N . Since $\mathbf{E}\tilde{X} = 0$, $\mathbf{E}\tilde{X}^2 = 2\sigma^2$, we get, by Chebyshev's inequality,

$$1 - Q = \mathbf{P}\left\{|\tilde{X}| \geq \frac{N + \frac{1}{2}}{t}\right\} \leq 2\left(\frac{\sigma t}{N + \frac{1}{2}}\right)^2. \quad (3.5)$$

If $\sigma t > 1/4$, then we choose $N = [2\sigma t + 1/2]$, in which case $Q \geq 1/2$ and $2N+1 \leq 12\sigma t$. Hence

$$\frac{t^2}{(2N+1)^2} Q^3 \geq \frac{t^2}{(12\sigma t)^2 \cdot 8} = \frac{1}{1152\sigma^2},$$

and thus

$$1 - |f(2\pi t)| \geq \frac{1}{3456M^2\sigma^2}, \quad \sigma t \geq \frac{1}{4}.$$

If $\sigma t \leq 1/4$, then the choice $N = 0$ in (3.5) yields $Q \geq 1/2$. By (3.4),

$$1 - |f(2\pi t)| \geq \frac{t^2}{24M^2}, \quad \sigma t \leq \frac{1}{4}. \quad \square$$

4 Separation of characteristic functions (reduction to dimension one)

In the multidimensional case, the characteristic function

$$f(t) = \mathbf{E}e^{i\langle t, X \rangle}, \quad t \in \mathbb{R}^d,$$

admits a similar bound. The next statement is a preliminary step in the proof of Theorem 1.

Proposition 2. *Given a random vector X in \mathbb{R}^d with covariance matrix $\sigma^2 I_d$, $\sigma > 0$, and a finite maximum of density $M = M(X)$, its characteristic function satisfies*

$$|f(t)| \leq 1 - \frac{c^d}{M^2\sigma^{2d}} \min\{\sigma^2|t|^2, 1\}, \quad t \in \mathbb{R}^d, \quad (4.1)$$

with some absolute constant $c > 0$.

Note that the isotropic constant $M^{1/d}\sigma$, being bounded away from zero according to (2.4), is present on the right-hand side of (4.1). For isotropic log-concave distributions, it is also bounded from above by a d -dependent constant according to (2.5). In this case, we can certainly obtain better bounds with decay of $f(t)$ at infinity.

For the proof, it looks natural to apply the one-dimensional result to random variables $X_\theta = \langle \theta, X \rangle$ with unit vectors θ . However, it may happen that X_θ will have an unbounded density, which means that Proposition 1 is not applicable. For example, in dimension $d = 2$, suppose that $X = (X_1, X_2)$ has a uniform distribution on the unbounded region $R = \{(x_1, x_2): |x_1| \leq \exp(-c|x_2|)\}$, $c > 0$, that is, with density

$$p(x_1, x_2) = \frac{c}{4} \mathbf{1}_R(x_1, x_2), \quad x_1, x_2 \in \mathbb{R}.$$

Then X_1 takes values in $(-1, 1)$ and has the density

$$p_1(x_1) = \int_{-\infty}^{\infty} p(x_1, x_2) \, dx_2 = \frac{1}{2} \log \frac{1}{|x_1|}, \quad -1 < x_1 < 1,$$

which is unbounded near zero. It is easy to check that the random vector X is isotropic, that is, $\mathbf{E}X_1X_2 = 0$ and $\mathbf{E}X_1^2 = \mathbf{E}X_2^2$ for a suitable constant c .

Proof of Proposition 2. Since $|f(t)|$ does not change when adding any fixed vector to X , we may assume that this random vector has mean zero. Let p denote the density of X .

The above example shows that a preliminary density truncation is desirable. Introduce the random vector X_r in \mathbb{R}^d with parameter $r > 0$ with density

$$p_r(x) = \frac{1}{b_r} p(x) \mathbf{1}_{\{|x| < r\}}, \quad x \in \mathbb{R}^d,$$

where $b_r = \mathbf{P}\{|X| < r\}$ is a normalizing constant. Assuming that $d \geq 2$, we choose $r = \sigma\sqrt{2d}$, which guarantees, by Markov’s inequality, that

$$b_r = 1 - \mathbf{P}\{|X| \geq r\} \geq 1 - \frac{\mathbf{E}|X|^2}{r^2} = 1 - \frac{d\sigma^2}{r^2} = \frac{1}{2}.$$

Consider the random vectors $\tilde{X} = X - Y$ and $\tilde{X}_r = X_r - Y_r$, where Y and Y_r are independent copies of X and X_r , respectively, and denote by \tilde{p} and \tilde{p}_r their densities. The function $\tilde{p}_r(x)$ is supported on the ball $|x| < 2r$. Moreover, for all $x \in \mathbb{R}^d$,

$$\begin{aligned} \tilde{p}_r(x) &= \int_{\mathbb{R}^d} p_r(x+y)p_r(y) \, dy = \frac{1}{b_r^2} \int_{|y| < r} p(x+y)p(y) \mathbf{1}_{\{|x+y| < r\}} \, dy \\ &\leq 4 \int_{\mathbb{R}^d} p(x+y)p(y) \, dy = 4\tilde{p}(x). \end{aligned} \tag{4.2}$$

Put $t = s\theta$, $s \in \mathbb{R}$, $|\theta| = 1$, and consider the characteristic functions

$$g_r(s) = \mathbf{E}e^{i\langle t, X_r \rangle} = \mathbf{E}e^{is\langle \theta, X_r \rangle}, \quad g(s) = \mathbf{E}e^{i\langle t, X \rangle} = \mathbf{E}e^{is\langle \theta, X \rangle},$$

so that

$$|g_r(s)|^2 = \mathbf{E}e^{i\langle t, \tilde{X}_r \rangle} = \mathbf{E}e^{is\langle \theta, \tilde{X}_r \rangle}, \quad |g(s)|^2 = \mathbf{E}e^{i\langle t, \tilde{X} \rangle} = \mathbf{E}e^{is\langle \theta, \tilde{X} \rangle},$$

Then, by (4.2),

$$\begin{aligned} 1 - |g(s)|^2 &= 2 \int_{\mathbb{R}^d} \sin^2\left(\frac{\langle t, x \rangle}{2}\right) \tilde{p}(x) \, dx \geq \frac{1}{2} \int_{\mathbb{R}^d} \sin^2\left(\frac{\langle t, x \rangle}{2}\right) \tilde{p}_r(x) \, dx \\ &= \frac{1}{4} (1 - |g_r(s)|^2). \end{aligned} \quad (4.3)$$

To bound from below the right-hand side, first, note that

$$\begin{aligned} \text{Var}(\langle \theta, X_r \rangle) &= \frac{1}{2b_r^2} \int_{|x| < r} \int_{|y| < r} \langle \theta, x - y \rangle^2 p(x)p(y) \, dx \, dy \\ &\leq \frac{1}{2b_r^2} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d} \langle \theta, x - y \rangle^2 p(x)p(y) \, dx \, dy = \frac{\sigma^2}{b_r^2} \leq 4\sigma^2. \end{aligned}$$

Thus

$$\text{Var}(\langle \theta, X_r \rangle) \leq 4\sigma^2. \quad (4.4)$$

The maximum of density of the random variable $\langle \theta, X_r \rangle$ can also be related to $M = M(X)$. For simplicity, let $\theta = e_1 = (1, 0, \dots, 0)$, so that $\langle \theta, X_r \rangle$ has the one-dimensional density

$$\int_{\mathbb{R}^{d-1}} p_r(x, y) \, dy \leq \frac{1}{b_r} \int_{|y| < r} p(x, y) \, dy \leq 2\omega_{d-1} r^{d-1} M, \quad x \in \mathbb{R}.$$

Hence, for any unit vector θ in \mathbb{R}^d ,

$$M(\langle \theta, X_r \rangle) \leq 2\omega_{d-1} r^{d-1} M. \quad (4.5)$$

Applying Proposition 1 to the characteristic function $g_r(s)$ with its properties (4.4)–(4.5), it follows that

$$1 - |g_r(s)|^2 \geq \frac{c}{\omega_{d-1}^2 r^{2(d-1)} M^2 \sigma^2} \min\{\sigma^2 s^2, 1\}$$

up to some absolute constant $c > 0$. Using this in (4.3), we arrive at the similar relation

$$1 - |g(s)|^2 \geq \frac{c}{\omega_{d-1}^2 r^{2(d-1)} M^2 \sigma^2} \min\{\sigma^2 s^2, 1\},$$

that is,

$$1 - |g(s)|^2 \geq \frac{c}{\omega_{d-1}^2 (2d)^{d-1} M^2 \sigma^{2d}} \min\{\sigma^2 s^2, 1\}.$$

To simplify the constants, recall formula (2.3) and Batir's bounds for the gamma function [4]

$$\sqrt{2e} \left(\frac{x}{e}\right)^x \leq \Gamma\left(x + \frac{1}{2}\right) \leq \sqrt{2\pi} \left(\frac{x}{e}\right)^x, \quad x \geq \frac{1}{2}.$$

The lower bound gives

$$\omega_{d-1}(2d)^{(d-1)/2} = \frac{\pi^{(d-1)/2}}{\Gamma(\frac{d+1}{2})} (2d)^{(d-1)/2} \leq \frac{1}{2\sqrt{\pi ed}} (4\pi e)^{d/2}.$$

It remains to note that $1 - |g(s)|^2 \leq 2(1 - |g(s)|)$, and (4.1) follows. \square

5 Maximum of convolved densities

Convolved densities are known to have improved smoothing properties. First, let us emphasize the following general fact.

Proposition 3. *If independent random vectors X_1, \dots, X_m ($m \geq 2$) with values in \mathbb{R}^d have bounded densities, then the sum $S_m = X_1 + \dots + X_m$ has a bounded uniformly continuous density vanishing at infinity.*

Proof. Denote by q_k the densities of X_k and assume that $q_k(x) \leq M_k$ for all $x \in \mathbb{R}^d$ with some constants M_k ($k \leq m$). By the Plancherel theorem, for the characteristic functions $v_k(t) = \mathbf{E}e^{i\langle t, X_k \rangle}$, we have

$$\begin{aligned} \int_{\mathbb{R}^d} |v_k(t)|^m dt &\leq \int_{\mathbb{R}^d} |v_k(t)|^2 dt = (2\pi)^d \int_{\mathbb{R}^d} q_k(x)^2 dx \leq (2\pi)^d \int_{\mathbb{R}^d} M_k q_k(x) dx \\ &= (2\pi)^d M_k, \end{aligned}$$

where we used the property $|v_k(t)| \leq 1$, $t \in \mathbb{R}^d$. Hence by Hölder’s inequality the characteristic function $f(t) = v_1(t) \cdots v_m(t)$ of S_m is integrable and has the L^1 -norm

$$\begin{aligned} \int_{\mathbb{R}^d} |f(t)| dt &\leq \left(\int_{\mathbb{R}^d} |v_1(t)|^m dt \right)^{1/m} \cdots \left(\int_{\mathbb{R}^d} |v_m(t)|^m dt \right)^{1/m} \\ &\leq (2\pi)^d (M_1 \cdots M_m)^{1/m} < \infty. \end{aligned} \tag{5.1}$$

We may conclude that the random variable S_m has a bounded, uniformly continuous density expressed by the inversion Fourier formula

$$q(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-i\langle t, x \rangle} f(t) dt, \quad x \in \mathbb{R}^d. \tag{5.2}$$

Since f is integrable, it also follows that $q(x) \rightarrow 0$ as $|x| \rightarrow \infty$ by the Riemann–Lebesgue lemma. \square

Since, by (5.2),

$$q(x) \leq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} |f(t)| dt$$

for all $x \in \mathbb{R}^d$, inequality (5.1) also implies that

$$M(S_m) \leq (M(X_1) \cdots M(X_m))^{1/m}. \tag{5.3}$$

However, relation (5.3) does not correctly reflect the behavior of $M(S_m)$ with respect to the growing parameter m , especially in the i.i.d. situation. A more precise statement from [7] is described in the following relation, where the geometric mean of maxima is replaced with the harmonic mean.

Proposition 4. For independent random vectors X_k , $1 \leq k \leq m$, with values in \mathbb{R}^d , we have

$$\frac{1}{M(S_m)^{2/d}} \geq \frac{1}{e} \sum_{k=1}^m \frac{1}{M(X_k)^{2/d}}. \quad (5.4)$$

This bound may be viewed as a counterpart of the entropy power inequality in information theory. It is derived by applying the Hausdorff–Young inequality with best constants (due to Beckner and Lieb). The constant $1/e$ is optimal and is attained asymptotically as $d \rightarrow \infty$ for random vectors uniformly distributed on Euclidean balls. However, in dimension $d = 1$, it can be improved to $1/2$, which follows from results due to Rogozin [19] and Ball [2].

One useful consequence of (5.4) is the following bound on the L^{2m} -norms of characteristic functions.

Proposition 5. If $f(t)$ is the characteristic function of a random vector X in \mathbb{R}^d , then for any integer $m \geq 1$,

$$\frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} |f(t)|^{2m} dt \leq \left(\frac{e}{2m} \right)^{d/2} M(X). \quad (5.5)$$

Proof. We apply Proposition 4 to $2m$ summands $X_1, -X'_1, \dots, X_{m_2} - X'_m$, assuming that X_k and X'_k are independent copies of X . Introduce the symmetrized random vector $S_m = S_m - S'_m$, where S'_m is an independent copy of S_m . By (5.4) we then get

$$M(\tilde{S}_m) \leq \left(\frac{e}{2m} \right)^{d/2} M(X).$$

In addition, \tilde{S}_m has the characteristic function $|f(t)|^{2m}$. If $M(X)$ is finite, then we may apply Proposition 5 and conclude that \tilde{S}_m has a bounded continuous density $q_m(x)$ vanishing at infinity. Moreover, $q_m(x)$ is maximized at $x = 0$, and its value at this point is described by the inversion formula (5.2), which gives

$$M(\tilde{S}_m) = q_m(0) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} |f(t)|^{2m} dt. \quad \square$$

Using (5.3), we can obtain a similar relation, but without the factor $(e/(2m))^{d/2}$ in (5.5).

When $M(X)$ is finite and m is large, the bound (5.5) may be considerably sharpened asymptotically with respect to m when restricting the integration to the regions $|t| \geq \varepsilon > 0$.

Proposition 6. Let f be the characteristic function of a random variable X with covariance matrix $\sigma^2 I_d$ ($\sigma > 0$) and finite $M = M(X)$. For any $\varepsilon > 0$ and $n \geq 2$,

$$\int_{|t| \geq \varepsilon} |f(t)|^n dt \leq \left(\frac{8\pi^2 e}{n} \right)^{d/2} M \exp \left\{ -\frac{c^d n}{M^2 \sigma^{2d}} \min(\sigma^2 \varepsilon^2, 1) \right\} \quad (5.6)$$

with some absolute constant $c > 0$.

Proof. If $n = 2$, then we may extend the integration in (5.6) to the whole space and apply (5.5) with $m = 1$. Then using the lower bound (2.4), namely $M^2 \sigma^{2d} \geq (2\pi e)^{-d}$, inequality (5.6) will follow from

$$1 \leq 2e^{-4c_1^d/d}, \quad c_1 = c \cdot 2\pi e.$$

This holds with $c_1 = 1/8$, for example.

If $n \geq 3$, then write $n = 2m + k$ with $k = 1$ or $k = 2$ for $n \leq 5$ and with $m = \lceil n/3 \rceil$, $k = n - 2m$ for $n \geq 6$. By Proposition 2 the function $\delta_f(\varepsilon) = \max_{|t| \geq \varepsilon} |f(t)| < 1$ admits the upper bound

$$\delta_f(\varepsilon) \leq 1 - \frac{c^d}{M^2 \sigma^{2d}} \min(\sigma^2 \varepsilon^2, 1) \tag{5.7}$$

with some absolute constant $c > 0$. Hence, by (5.5) and (5.7),

$$\int_{|t| \geq \varepsilon} |f(t)|^n dt \leq \delta_f(\varepsilon)^k \int_{\mathbb{R}^d} |f(t)|^{2m} dt \leq \left(\frac{2\pi^2 e}{m} \right)^{d/2} M \exp \left\{ -\frac{c^d k}{M^2 \sigma^{2d}} \min(\sigma^2 \varepsilon^2, 1) \right\}.$$

It remains to note that $m \geq n/4$ and $k \geq n/5$. \square

6 Normal approximation for products of characteristic functions

Let us now recall standard results about the approximation of products of characteristic functions. Consider the sum

$$S_n = \xi_1 + \dots + \xi_n$$

of independent random variables ξ_k with mean zero and standard deviations b_k such that $b_1^2 + \dots + b_n^2 = 1$, so that S_n has mean zero and variance one.

In terms of the characteristic functions $v_k(t) = \mathbf{E}e^{it\xi_k}$, the characteristic function of S_n represents the product

$$f_n(t) = \mathbf{E}e^{itS_n} = v_1(t) \dots v_n(t), \quad t \in \mathbb{R}.$$

Under higher-order moment assumptions, it may be approximated by the standard normal characteristic function $g(t) = e^{-t^2/2}$ or by Edgeworth corrections on relatively large t -intervals by means of the Lyapunov coefficients

$$L_p = \sum_{k=1}^n \mathbf{E}|\xi_k|^p, \quad p > 2,$$

provided that they are small. We only mention such results for the particular indexes $p = 3$ and $p = 4$.

Proposition 7. *With some absolute constants $C > 0$ and $c > 0$,*

$$|f_n(t) - e^{-t^2/2}| \leq CL_3 |t|^3 e^{-ct^2}, \quad |t| \leq \frac{1}{L_3}. \tag{6.1}$$

Moreover, if $\mathbf{E}\xi_k^3 = 0$ for all $k \leq n$, then

$$|f_n(t) - e^{-t^2/2}| \leq CL_4 t^4 e^{-ct^2}, \quad |t| \leq \frac{1}{\sqrt{L_4}}. \tag{6.2}$$

Inequalities (6.1)–(6.2) are often stated in a slightly different form. For example, Petrov [18, p. 109] derived relation (6.1) on a smaller interval $|t| \leq 1/(4L_3)$ with $C = 16$ and $c = 1/3$. As we can see from the proof or properly modifying it, the interval of approximation can be increased to $|t| \leq c_0/L_3$ with some absolute constant $c_0 > 1$ at the expense of a smaller value of c and a larger value of C . Similar relations with arbitrary real $p > 2$ can be found in the review [6].

In the noninteresting case where L_3 or L_4 are greater than 1, (6.1)–(6.2) hold on the larger interval $|t| \leq 1$. This can be seen from the Taylor integral formula for the function $h(t) = f_n(t) - e^{-t^2/2}$ about the point $t = 0$, using the property that the first two and three derivatives of $h(t)$ at the origin are respectively vanishing.

In general, the function $p \rightarrow L_p^{1/(p-2)}$ is nondecreasing with $L_p \geq n^{-(p-2)/2}$. In particular,

$$\frac{1}{\sqrt{n}} \leq L_3 \leq \sqrt{L_4}. \quad (6.3)$$

Since $\mathbf{E}|\xi_k|^p \geq (\mathbf{E}\xi_k^2)^{p/2}$, we also have

$$L_p \geq b_1^p + \cdots + b_n^p \geq \left(\max_k b_k\right)^p. \quad (6.4)$$

Now, if instead of S_n , we consider the normalized sum

$$Z_n = \frac{X_1 + \cdots + X_n}{\sqrt{n}},$$

then these results can be correspondingly reformulated. Then we should assume that the independent random variables X_k have mean zero and standard deviations σ_k such that $\sigma_1^2 + \cdots + \sigma_n^2 = n$. In this setting, it is more natural to represent the Lyapunov coefficients as

$$L_p = n^{-(p-2)/2} \beta_p, \quad \beta_p = \frac{1}{n} \sum_{k=1}^n \mathbf{E}|X_k|^p.$$

Thus $\beta_p = \mathbf{E}|X|^p$ when X_k are independent copies of a random variable X .

In general, from (6.3)–(6.4) it follows that

$$1 \leq \beta_3 \leq \sqrt{\beta_4}, \quad 1 \leq \max_k \sigma_k \leq (n\beta_p)^{1/p}. \quad (6.5)$$

Finally, let us state Proposition 7 once more about the normalized sums.

Proposition 8. *With some absolute constants $C > 0$ and $c > 0$, the characteristic function $f_n(t)$ of Z_n satisfies*

$$|f_n(t) - e^{-t^2/2}| \leq \frac{C\beta_3}{\sqrt{n}} |t|^3 e^{-ct^2}, \quad |t| \leq \frac{\sqrt{n}}{\beta_3}. \quad (6.6)$$

Moreover, if $\mathbf{E}X_k^3 = 0$ for all $k \leq n$, then

$$|f_n(t) - e^{-t^2/2}| \leq \frac{C\beta_4}{n} t^4 e^{-ct^2}, \quad |t| \leq \frac{\sqrt{n}}{\sqrt{\beta_4}}. \quad (6.7)$$

7 Refinement of Theorem 1

Let us return to the setting of Theorem 1 and assume that the independent random vectors X_k have mean zero, covariance matrices $\sigma_k^2 I_d$ such that $\sigma_1^2 + \cdots + \sigma_n^2 = n$, and finite maxima of densities $M_k = M(X_k)$. Recall the notations

$$\beta_3 = \sup_{|\theta|=1} \left[\frac{1}{n} \sum_{k=1}^n \mathbf{E}|\langle \theta, X_k \rangle|^3 \right], \quad \beta_4 = \sup_{|\theta|=1} \left[\frac{1}{n} \sum_{k=1}^n \mathbf{E}\langle \theta, X_k \rangle^4 \right],$$

and

$$\Delta_n = \sup_x |p_n(x) - \varphi(x)|.$$

Here we prove this theorem in a somewhat more general form (although more complicated).

Theorem 2. *With some positive absolute constants C and c , the density p_n of Z_n satisfies*

$$\Delta_n \leq \frac{C^d \beta_3}{\sqrt{n}} + C^d (M_1 \cdots M_n)^{1/n} \exp \left\{ -c^d \sum_{k=1}^n \frac{1}{M_k^2 \sigma_k^{2d}} \min \left(\frac{\sigma_k^2}{\beta_3^2}, 1 \right) \right\}. \tag{7.1}$$

Moreover, if $\mathbf{E}\langle \theta, X_k \rangle^3 = 0$ for all $\theta \in \mathbb{R}^d$ and $k \leq n$ (in particular, if the distributions of X_k are symmetric), then

$$\Delta_n \leq \frac{C^d \beta_4}{n} + C^d (M_1 \cdots M_n)^{1/n} \exp \left\{ -c^d \sum_{k=1}^n \frac{1}{M_k^2 \sigma_k^{2d}} \min \left(\frac{\sigma_k^2}{\beta_4}, 1 \right) \right\}. \tag{7.2}$$

Proof. In the case $n = 1$, it is assumed that $\sigma_1 = 1$, implying that $M_1 \geq (2\pi e)^{d/2}$ according to (2.4). Since also $\sqrt{\beta_4} \geq \beta_3 \geq 1$ (see (6.5)), inequalities (7.1)–(7.2) are fulfilled automatically. So, assume that $n \geq 2$.

In terms of the characteristic functions $v_k(t) = \mathbf{E}e^{i\langle t, X_k \rangle}$, the characteristic function of Z_n is given by the product

$$f_n(t) = \mathbf{E}e^{i\langle t, Z_n \rangle} = v_1 \left(\frac{t}{\sqrt{n}} \right) \cdots v_n \left(\frac{t}{\sqrt{n}} \right).$$

Applying Hölder’s inequality and then Proposition 5, we get

$$\int_{\mathbb{R}^d} |f_n(t)| dt \leq \prod_{k=1}^n \left(\int_{\mathbb{R}^d} \left| v_k \left(\frac{t}{\sqrt{n}} \right) \right|^n dt \right)^{1/n} < \infty.$$

Hence we may apply the Fourier inversion formula to represent the densities of Z_n as

$$p_n(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{-i\langle t, x \rangle} f_n(t) dt, \quad x \in \mathbb{R}^d.$$

Using a similar representation for the standard normal density, we get

$$\Delta_n \leq \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} |f_n(t) - e^{-|t|^2/2}| dt. \tag{7.3}$$

Next, we split the integration in (7.3) into two regions. From Proposition 6 applied to the characteristic function $v_k(t)$ of X_k , we obtain that for any $\varepsilon > 0$,

$$\int_{|t| \geq \varepsilon} |v_k(t)|^n dt \leq \left(\frac{C}{\sqrt{n}} \right)^d M_k \exp \left\{ -c^d \frac{n}{M_k^2 \sigma_k^{2d}} \min(\sigma_k^2 \varepsilon^2, 1) \right\}$$

with some absolute constants $C > 0$ and $c > 0$. Equivalently,

$$\int_{|t| \geq \varepsilon \sqrt{n}} \left| v_k \left(\frac{t}{\sqrt{n}} \right) \right|^n dt \leq C^d M_k \exp \left\{ -c^d \frac{n}{M_k^2 \sigma_k^{2d}} \min(\sigma_k^2 \varepsilon^2, 1) \right\}.$$

Applying once more Hölder's inequality

$$\int_{|t| \geq \varepsilon \sqrt{n}} |f_n(t)| dt \leq \prod_{k=1}^n \left(\int_{|t| \geq \varepsilon \sqrt{n}} \left| v_k \left(\frac{t}{\sqrt{n}} \right) \right|^n dt \right)^{1/n},$$

we then get

$$\int_{|t| \geq \varepsilon \sqrt{n}} |f_n(t)| dt \leq C^d (M_1 \cdots M_n)^{1/n} \exp \left\{ -c^d \sum_{k=1}^n \frac{\min(\sigma_k^2 \varepsilon^2, 1)}{M_k^2 \sigma_k^{2d}} \right\}.$$

Since

$$\int_{|t| \geq \varepsilon \sqrt{n}} e^{-|t|^2/2} dt \leq C^d e^{-n\varepsilon^2/4}$$

for some absolute constant $C > 0$, it follows that

$$\int_{|t| \geq \varepsilon \sqrt{n}} |f_n(t) - e^{-|t|^2/2}| dt \leq C^d e^{-n\varepsilon^2/4} + C^d (M_1 \cdots M_n)^{1/n} \exp \left\{ -c^d \sum_{k=1}^n \frac{\min(\sigma_k^2 \varepsilon^2, 1)}{M_k^2 \sigma_k^{2d}} \right\}. \quad (7.4)$$

We now turn to the other region $|t| < \varepsilon \sqrt{n}$. By the mean zero and isotropy assumption on the distribution of X_k we have $\mathbf{E}\langle \theta, X_k \rangle = 0$ and $\mathbf{E}\langle \theta, X_k \rangle^2 = \sigma_k^2$ for any unit vector θ in \mathbb{R}^d . Therefore we are in position to apply Proposition 8 to the random variables $\langle \theta, X_k \rangle$. Write $t = s\theta$ for $s > 0$ and $|\theta| = 1$. Since $s \mapsto f_n(s\theta)$ represents the characteristic function of the normalized sum of $\langle \theta, X_k \rangle$, it satisfies, by (6.6),

$$|f_n(s\theta) - e^{-s^2/2}| \leq \frac{C\beta_3(\theta)}{\sqrt{n}} s^3 e^{-cs^2}, \quad s \leq \frac{\sqrt{n}}{\beta_3(\theta)}, \quad (7.5)$$

where

$$\beta_3(\theta) = \frac{1}{n} \sum_{k=1}^n \mathbf{E}|\langle \theta, X_k \rangle|^3.$$

In (7.5), $\beta_3(\theta)$ can be replaced with the larger value β_3 , which leads to

$$|f_n(t) - e^{-|t|^2/2}| \leq \frac{C\beta_3}{\sqrt{n}} |t|^3 e^{-c|t|^2}, \quad |t| \leq T_n = \frac{\sqrt{n}}{\beta_3}.$$

This readily yields

$$\int_{|t| \leq T_n} |f_n(t) - e^{-|t|^2/2}| dt \leq \frac{C^d \beta_3}{\sqrt{n}}$$

with some absolute constant C . We can now combine this inequality with (7.4) by choosing $\varepsilon = 1/\beta_3$ and apply these inequalities in (7.3), which yields

$$\Delta_n \leq C^d \left[\frac{\beta_3}{\sqrt{n}} + e^{-n/(4\beta_3^2)} + (M_1 \cdots M_n)^{1/n} \exp \left\{ -c^d \sum_{k=1}^n \frac{\min(\sigma_k^2 \varepsilon^2, 1)}{M_k^2 \sigma_k^{2d}} \right\} \right].$$

Since $\beta_3 \geq 1$, we arrive at (7.1).

In the second scenario, we similarly apply inequality (6.7) and combine it once more with (7.4) by choosing $\varepsilon = 1/\sqrt{\beta_4}$. \square

8 Proofs of Theorem 1 and Corollary 1

The right-hand sides in (7.1)–(7.2) may be bounded and simplified in terms of the functionals

$$M = \max_k M(X_k), \quad \sigma^2 = \max_k \sigma_k^2.$$

Proof of Theorem 1. The k th term in the sum in (7.1),

$$\frac{1}{M_k^2 \sigma_k^{2d}} \min \left(\frac{\sigma_k^2}{\beta_3^2}, 1 \right),$$

is a decreasing function with respect to σ_k^2 and M_k^2 . Hence the second summand on the right-hand side in (7.1) does not exceed

$$C^d M \exp \left\{ -\frac{c^d n}{M^2 \sigma^{2d}} \min \left(\frac{\sigma^2}{\beta_3^2}, 1 \right) \right\} \leq C^d M \exp \left\{ -\frac{c^d n}{M^2 \sigma^{2d} \beta_3^2} \right\},$$

where we used $\sigma^2 \geq 1$ and $\beta_3 \geq 1$, implying that $\min(\sigma^2/\beta_3^2, 1) \geq \beta_3^{-2}$. Using $e^{-x} < x^{1/2}$ ($x > 0$), from (7.1) we obtain that

$$\Delta_n \leq \frac{C^d \beta_3}{\sqrt{n}} + C^d \frac{M^2 \sigma^d \beta_3}{\sqrt{n}}$$

with some absolute constant C . In view of the relation $M \sigma^d \geq (2\pi e)^{-d/2}$, the above estimate is simplified to (1.1).

With similar arguments, the second summand on the right-hand side in (7.2) does not exceed

$$C^d M \exp \left\{ -\frac{c^d n}{M^2 \sigma^{2d} \beta_4} \right\}.$$

Using $e^{-x} < x^{-1}$ ($x > 0$), from (7.2) we therefore obtain that

$$\Delta_n \leq \frac{C^d \beta_4}{n} + C^d \frac{M^3 \sigma^{2d} \beta_4}{n}.$$

Again, the first term on the right-hand side is dominated by the second one up to the multiple, so that the above estimate is simplified to (1.2). This proves Theorem 1. \square

Proof of Corollary 1. First, we need to mention that if a random vector X has a log-concave density, then so does any linear functional $\xi = \langle \theta, X \rangle$. This is a consequence of a well-known characterization of log-concave

measures due to Borell [14]. He also derived a large deviation bound for norms under log-concave measures, which implies in dimension one that the L^p -norms of ξ are equivalent to each other. More precisely,

$$(\mathbf{E}|\xi|^p)^{1/p} \leq Cp(\mathbf{E}\xi^2)^{1/2}$$

for all $p > 2$ with some absolute constant C . Applying this with $X = X_k$, we conclude that

$$\begin{aligned} \frac{1}{n} \sum_{k=1}^n \mathbf{E}|\langle \theta, X_k \rangle|^p &\leq \frac{(Cp)^p}{n} \sum_{k=1}^n (\mathbf{E}|\langle \theta, X_k \rangle|^2)^{p/2} = \frac{(Cp)^p}{n} \sum_{k=1}^n \sigma_k^p \\ &\leq (Cp)^p, \quad |\theta| = 1, \end{aligned}$$

where the last inequality is due to the assumption that $\sigma_1^2 + \dots + \sigma_n^2 = n$. Hence β_3 and β_4 are bounded by absolute constants, so that inequalities (1.1)–(1.2) are respectively simplified to

$$\Delta_n \leq \frac{C^d M^2}{\sqrt{n}}, \quad \Delta_n \leq \frac{C^d M^3}{n}.$$

To prove (1.3)–(1.4), it remains to recall the bound (2.5), which implies $M \leq K_d^d$, where K_d is the maximal isotropic constant for the class of log-concave probability measures on \mathbb{R}^d . \square

Acknowledgment. We would like to thank the referee for a careful reading of the manuscript and useful comments.

Funding. Open Access funding enabled and organized by Projekt DEAL.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>

References

1. S. Artstein-Avidan, A. Giannopoulos, and V. D. Milman, *Asymptotic Geometric Analysis. Part I*, Math. Surv. Monogr., Vol. 202, AMS, Providence, RI, 2015.
2. K.M. Ball, Cube slicing in \mathbb{R}^n , *Proc. Am. Math. Soc.*, **97**(3):465–473, 1986.
3. K.M. Ball, Logarithmically concave functions and sections of convex bodies, *Stud. Math.*, **88**:69–84, 1988.
4. N. Batir, Inequalities for the gamma function, *Arch. Math.*, **91**(6):554–563, 2008.
5. R.N. Bhattacharya and R. Ranga Rao, *Normal Approximation and Asymptotic Expansions*, John Wiley & Sons, New York, 1976. Also published by SIAM, Philadelphia, 2010.
6. S.G. Bobkov, Asymptotic expansions for products of characteristic functions under moment assumptions of non-integer orders, in E. Carlen, M. Madiman, and E.M. Werner (Eds.), *Convexity and Concentration*, IMA Vol. Math. Appl., Vol. 161, Springer, New York, 2017, pp. 297–357.

7. S.G. Bobkov and G.P. Chistyakov, Bounds for the maximum of the density of the sum of independent random variables, *J. Math. Sci., New York*, **199**(2):100–106, 2014. Translated from *Zap. Nauchn. Semin. POMI*, **408**:62–73, 2012.
8. S.G. Bobkov, G.P. Chistyakov, and F. Götze, Bounds for characteristic functions in terms of quantiles and entropy, *Electron. Commun. Probab.*, **17**(21):9, 2012.
9. S.G. Bobkov, G.P. Chistyakov, and F. Götze, Berry–Esseen bounds in the entropic central limit theorem, *Probab. Theory Relat. Fields*, **159**(3–4):435–478, 2014.
10. S.G. Bobkov, G.P. Chistyakov, and F. Götze, Richter’s local limit theorem, its refinement, and related results, *Lith. Math. J.*, **63**(2):138–160, 2023.
11. S.G. Bobkov and F. Götze, Central limit theorem for Rényi divergence of infinite order, *Ann. Probab.*, 2025 (to appear), arXiv:2402.02259v2.
12. S.G. Bobkov and M. Madiman, The entropy per coordinate of a random vector is highly constrained under convexity conditions, *IEEE Trans. Inf. Theory*, **57**(8):4940–4954, 2011.
13. S.G. Bobkov and F.L. Nazarov, On convex bodies and log-concave probability measures with unconditional basis, in V.D. Milman and G. Schechtman (Eds.), *Geometric Aspects of Functional Analysis. Israel Seminar 2001–2002*, Lect. Notes Math., Vol. 1807, Springer, Berlin, 2003, pp. 53–69.
14. C. Borell, Convex measures on locally convex spaces, *Ark. Mat.*, **12**:239–252, 1974.
15. S. Brazitikos, A. Giannopoulos, P. Valettas, and B.-H. Vritsiou, *Geometry of Isotropic Convex Bodies*, Math. Surv. Monogr., Vol. 196, AMS, Providence, RI, 2014.
16. D. Hensley, Slicing convex bodies—bounds for slice area in terms of the body’s covariance, *Proc. Am. Math. Soc.*, **79**(4):619–625, 1980.
17. B. Klartag, Logarithmic bounds for isoperimetry and slices of convex sets, *Ars Inventionis Analyticae*, Paper No. 4, 2023.
18. V.V. Petrov, *Sums of Independent Random Variables*, Ergeb. Math. Grenzgeb., Band 82, Springer, New York, Heidelberg, 1975.
19. B.A. Rogozin, An estimate for the maximum of the convolution of bounded densities, *Theory Probab. Appl.*, **32**(1):48–56, 1987. Translated from *Teor. Veroyatn. Primen.*, **32**(1):53–61, 1987.
20. V. A. Statulevičius, Limit theorems for densities and the asymptotic expansions for distributions of sums of independent random variables, *Theory Probab. Appl.*, **10**(4):582–595, 1965. Translated from *Teor. Veroyatn. Primen.*, **10**:645–659, 1965.