

# Risk & Asset Allocation

## Case Solution for Spring 1

John A. Dodson

January 23, 2013

Say  $X$  is the invariant random variable that will drive market prices  $P_{T+\tau} \triangleq g(X; p_T)$  between today, time  $T$ , and the next decision date, time  $T + \tau$  with an investment horizon of  $\tau$  (all time measured in years) and say that we have used historical data to estimate the parameters of the characterization of the random variable  $\tilde{X}$  under a different (usually shorter) horizon,  $\tilde{\tau}$ ; e.g., from a timeseries sample  $(p_T, p_{T-\tilde{\tau}}, p_{T-2\tilde{\tau}}, \dots)$ .

Furthermore, say our invariant mapping is additive in the sense that

$$g(\tilde{x}_1; g(\tilde{x}_2, p_{T-2\tilde{\tau}})) = g(\tilde{x}_1 + \tilde{x}_2; p_{T-2\tilde{\tau}})$$

which is true, for example, with the continuous version of total return (although the simple version of total return is often an adequate approximation).

Then projecting  $X$  from  $\tilde{X}$  essentially involves the transformation

$$X \triangleq \overbrace{\tilde{X} + \tilde{X} + \dots}^{\frac{\tau}{\tilde{\tau}} \text{ i.i.d. copies}}$$

We know from the properties of characteristic functions that as long as increments are independent and identically distributed,

$$\phi_X(\omega) = \phi_{\tilde{X}}(\omega)^{\frac{\tau}{\tilde{\tau}}}$$

If the first two moments exist, we also know that

$$\begin{aligned} \mathbb{E} X &= -i \left. \frac{d\phi_X}{d\omega'} \right|_0 \\ &= -i \frac{\tau}{\tilde{\tau}} \phi_{\tilde{X}}^{\frac{\tau}{\tilde{\tau}}-1} \left. \frac{d\phi_{\tilde{X}}}{d\omega'} \right|_0 \\ &= \frac{\tau}{\tilde{\tau}} \mathbb{E} \tilde{X} \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}(XX') &= - \left. \frac{d^2\phi_X}{d\omega' d\omega} \right|_0 \\ &= - \frac{\tau}{\tilde{\tau}} \left( \frac{\tau}{\tilde{\tau}} - 1 \right) (\phi_{\tilde{X}})^{\frac{\tau}{\tilde{\tau}}-2} \frac{d\phi_{\tilde{X}}}{d\omega'} \frac{d\phi_{\tilde{X}}}{d\omega} - \frac{\tau}{\tilde{\tau}} (\phi_{\tilde{X}})^{\frac{\tau}{\tilde{\tau}}-1} \left. \frac{d^2\phi_{\tilde{X}}}{d\omega' d\omega} \right|_0 \\ &= \frac{\tau}{\tilde{\tau}} \left( \frac{\tau}{\tilde{\tau}} - 1 \right) \mathbb{E} \tilde{X} \mathbb{E} \tilde{X}' + \frac{\tau}{\tilde{\tau}} \mathbb{E} (\tilde{X} \tilde{X}') \\ &= \mathbb{E} X \mathbb{E} X' + \frac{\tau}{\tilde{\tau}} \left( \mathbb{E} (\tilde{X} \tilde{X}') - \mathbb{E} \tilde{X} \mathbb{E} \tilde{X}' \right) \end{aligned}$$

Since the covariance is defined as

$$\text{cov } \tilde{X} = E(XX') - E X E X'$$

we have that

$$\text{cov } X = \frac{\tau}{\tilde{\tau}} \text{cov } \tilde{X}$$

Furthermore, since

$$\text{std } \tilde{X} = \text{diag} \sqrt{\text{diag} \text{diag} \text{cov } \tilde{X}}$$

we have the “square-root rule” for time-scaling market invariants.

$$\text{std } X = \sqrt{\frac{\tau}{\tilde{\tau}}} \text{std } \tilde{X}$$

This is valid regardless of the distribution of  $\tilde{X}$  (as long as it has two moments).

Note that in general  $X$  will not belong to the same family of random variables as  $\tilde{X}$ .