

Copulas and Dependence

MFM Practitioner Module: Quantitative Risk Management

John Dodson

January 25, 2017

We have discussed continuous multivariate random variables and some broad parametric classes. It is clear that each instance of these involves at least one class of univariate random variable in the form of the marginals for the components. But it is also clear that the characterization of the original multivariate r.v. is not simply a collection of these marginal characterizations. There is a structure, with its own parameters, that connects them together.

- ▶ This is the **copula**.

For me, this is the prototypical example; and multivariate random variables are a rich source for parametric copulas. But it is not the only source. In fact, any random variable whose sample space is a unit hypercube with standard uniform margins is a copula.

To the extent that the *joint* density is not just a product of the *marginal* densities, there is dependence.

Factorization

This ratio can be expressed as

$$c(F_{X_1}(x_1), F_{X_2}(x_2), \dots) \triangleq \frac{f_{(X_1, X_2, \dots)}(x_1, x_2, \dots)}{f_{X_1}(x_1) f_{X_2}(x_2) \cdots}$$

Copula


Skalar's theorem says this is always possible. More generally, $c = f_U : [0, 1]^d \mapsto \mathbb{R}^+$ is a density function that characterizes a new random variable, U , that encapsulates the dependence structure of X .

Note that independence means $c \equiv 1$

Normal (Gaussian) Copula

When dependence can be entirely described by correlation, the Gaussian copula can be appropriate. For $d = 2$,

$$c(u) = \frac{1}{\sqrt{1-\rho^2}} \exp \left[\frac{-\rho}{1-\rho^2} \left(\rho \operatorname{erfc}^{-1}(2u_1)^2 \dots \right. \right. \\ \left. \left. + \rho \operatorname{erfc}^{-1}(2u_2)^2 - 2 \operatorname{erfc}^{-1}(2u_1) \operatorname{erfc}^{-1}(2u_2) \right) \right]$$



Gaussian copula density for $\rho = \frac{1}{2}$

Introduction

Concordance

Normal Mixture
CopulasArchimedean
Copulas

Upper & Lower Tail Dependence

Tail dependence is a pair-wise measure of the concordance of extreme outcomes.

$$\lambda_U = \lim_{p \uparrow 1} P \{X > F_X^{\leftarrow}(p) | Y > F_Y^{\leftarrow}(p)\}$$

$$\lambda_L = \lim_{p \downarrow 0} P \{X \leq F_X^{\leftarrow}(p) | Y \leq F_Y^{\leftarrow}(p)\}$$

The normal copula fails to exhibit tail dependence: extreme outcomes are essentially independent.

This is a problem, because in practice an extreme outcome in one component often acts to cause extreme outcomes in other components. Developing practical alternatives that include this **contagion effect** is an active area of research.

Several measures of **concordance** have been developed. Their definitions are motivated by the properties of their estimators, which we will not discuss just yet. Each ranges from -1 to 1 , with 0 for independence. In order of generality, we have

1. **Pearson's rho**. This is the classical linear correlation measure $\text{cov}(X, Y) / \sqrt{\text{var } X \text{ var } Y}$.
2. **Spearman's rho**. This is linear correlation applied to the **grades**, $F_X(X)$. It is a simple measure of dependence that is not sensitive to margins.
3. **Kendall's tau**. This is based strictly on the rank order of pairs of observations of pairs of components. It has useful theoretical and statistical properties.

N.B.: While independence implies zero concordance (under any of these definitions), zero concordance does not imply independence.

Kendall's tau can be defined as

$$\tau = 4 E C(U_1, U_2) - 1$$

where C is the distribution function characterizing the copula of X . It is the probability of concordance minus the probability of discordance for two independent draws of X .

Relationship with other measures

In general Spearman's rho is bounded by


$$\frac{3|\tau| - 1}{2} \operatorname{sgn} \tau \quad \& \quad \frac{1 + 2|\tau| - \tau^2}{2} \operatorname{sgn} \tau$$

For a Gaussian copula, Pearson's rho is

$$\rho = \sin\left(\frac{\pi}{2}\tau\right)$$

We use this to define **pseudo-correlation** for any elliptical r.v..

The relationship between Kendall's tau and Spearman's and Pearson's rho is illustrated by this graph.



For a given level of Kendall's tau, Spearman's rho is bounded by the two outer curves. Pearson's rho for a Gaussian copula is the curve through the origin.

Introduction

Concordance

Normal Mixture
CopulasArchimedean
Copulas

A normal mixture copula is simply the copula from a normal mixture multivariate random variable. The elliptical copula is an important subclass.

Elliptical Copula

An elliptical random variable is described by a mean vector, a dispersion matrix, and a characteristic generator function. It should be clear that the mean vector has no role in the copula. It should also be clear that the diagonal entries of the dispersion matrix also do not play a role.

Generally, an elliptical copula is parameterized by a semi-definite matrix with unit diagonals, which describe pair-wise dependence, and one or several **shape parameters** related to the characteristic generator.

The Gaussian copula is an example. Another important example is the t_ν copula, which we will work with in this week's exercise.

Introduction

Concordance

Normal Mixture
Copulas

Archimedean
Copulas

There are on the order of $d^2/2$ parameters to estimate for an elliptical copula. If you are dealing with a very large dimension, such as in a retail or securitization context, you either need a factor model to reduce the dimension or you should consider an Archimedean copula.

Archimedean Copulas

An Archimedean copula is defined in terms of a generator, a decreasing continuous function $\psi : [0, \infty) \mapsto [0, 1]$ with $\psi(0) = 1$ and $\lim_{t \rightarrow \infty} \psi(t) = 0$. The copula distribution is

$$C(u_1, u_2, \dots, u_d) = \psi(\psi^{-1}(u_1) + \psi^{-1}(u_2) + \dots + \psi^{-1}(u_d))$$

Three common single-parameter examples are the Gumbel, Clayton, and Frank.

Introduction

Concordance

Normal Mixture
CopulasArchimedean
Copulas