

(November 26, 2021)

## Complex analysis examples 04

Paul Garrett [garrett@math.umn.edu](mailto:garrett@math.umn.edu) <http://www.math.umn.edu/~garrett/>

[This document is [http://www.math.umn.edu/~garrett/m/complex/examples\\_2021-22/cx\\_ex\\_04.pdf](http://www.math.umn.edu/~garrett/m/complex/examples_2021-22/cx_ex_04.pdf)]

If you want feedback from me on your treatment of these examples, please email your work to me by Friday, Dec 03.

[04.1] Prove that, given holomorphic  $f, g$  on a non-empty open set  $U$ , and given a *simple* zero  $z_o$  of  $f$ , for all small-enough complex  $\varepsilon$  there is a unique zero of  $f + \varepsilon g$  nearest  $z_o$  and it is *simple*.

[04.2] For small  $w \in \mathbb{C}$ , let  $f(w)$  be the *simple* zero of  $z^5 - z + w = 0$  near 0. Determine a few terms of the power series expansion of  $f(w)$  at  $w = 0$ .

[04.3] Exhibit a linear fractional transformation mapping 1, 2, 3 to  $z_1, z_2, z_3$ .

[04.4] Exhibit a linear fractional transformation mapping the circle  $|z| = 1$ , minus a point, to the line  $\operatorname{Re}(z) = \operatorname{Im}(z)$ .

[04.5] Let  $z, z'$  be points in the open upper half-plane  $\mathfrak{H}$ . Exhibit a linear fractional transformation stabilizing  $\mathfrak{H}$  and mapping  $z$  to  $z'$ .

[04.6] Let  $z, z'$  be points in the open unit disk  $\mathfrak{D}$ . Exhibit a linear fractional transformation stabilizing  $\mathfrak{D}$  and mapping  $z$  to  $z'$ .

[04.7] Exhibit a conformal map of the open half disk  $\{z : |z| < 1, \operatorname{Re}(z) > 0\}$  to the open unit disk.

[04.8] Exhibit a conformal map of the open unit disk with  $[0, 1)$  removed to the open unit disk.

[04.9] Exhibit a conformal map of the sector  $\{re^{i\theta} : r > 0, 0 < \theta < \frac{\pi}{4}\}$  to the open unit disk.

[04.10] Exhibit a conformal map from the strip  $\{z = x + iy : c < ax + by < c'\}$  to the crescent

$$\Omega = \{z : |z| < 1, |z - \frac{1}{2}| > \frac{1}{2}\}$$

[04.11] Let  $f$  be holomorphic on  $\mathbb{C}$ , and meromorphic at infinity, with a pole of order  $N$ . Show that  $f$  is a polynomial of degree  $N$  (and conversely).

[04.12] Let holomorphic  $f : \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{C}\mathbb{P}^1$  be 2-to-1. Show that there are two linear fractional transformations  $\alpha, \beta$  such that  $\alpha \circ f \circ \beta$  is the map  $z \rightarrow z^2$ .