

(April 3, 2013)

Functions on circles: Fourier series, I

Paul Garrett garrett@math.umn.edu <http://www.math.umn.edu/~garrett/>

[This document is http://www.math.umn.edu/~garrett/m/fun/notes_2012-13/04_fourier_series_I.pdf]

1. Provocative example
2. Natural function spaces on the circle $S^1 = \mathbb{R}/2\pi\mathbb{Z}$
3. Topology on $C^\infty(S^1)$
4. Pointwise convergence of Fourier series
5. Distributions: generalized functions
6. Invariant integration, periodicization
7. Hilbert space theory of Fourier series
8. Levi-Sobolev inequality, Levi-Sobolev imbedding
9. $C^\infty(S^1) = \lim C^k(S^1) = \lim H^s(S^1)$
10. Distributions, generalized functions, again
11. The provocative example explained
12. Appendix: products and limits of topological vector spaces
13. Appendix: Fréchet spaces and limits of Banach spaces
14. Appendix: Urysohn and density of C^o

The simplest physical object with an interesting function theory is the circle, $S^1 = \mathbb{R}/2\pi\mathbb{Z}$, which inherits *group structure* and *translation-invariant* differential operator d/dx from the real line \mathbb{R} . The exponential functions $x \rightarrow e^{inx}$ for $n \in \mathbb{Z}$ are group homomorphisms $\mathbb{R}/2\pi\mathbb{Z} \rightarrow \mathbb{C}^\times$ and are eigenfunctions for d/dx on $\mathbb{R}/2\pi\mathbb{Z}$. Finite or infinite linear combinations

$$\sum_{n \in \mathbb{Z}} c_n e^{inx}$$

are *Fourier series*. [1] Conveniently, a function so expressed is a linear combination of eigenvectors for d/dx . That is, on functions *with* Fourier expansions [2] the linear operator of differentiation is *diagonalized*. However, infinite-dimensional linear algebra is subtler than finite-dimensional. Some fundamental questions are [3]

In what sense(s) can a function be expressed as a Fourier series?

Can a Fourier series be differentiated term-by-term?

How cautious must we be in differentiating functions that are only piecewise differentiable?

What will derivatives of discontinuous functions be?

[1] In the early 19th century, J. Fourier was an impassioned advocate of the use of such sums, of course writing sines and cosines rather than complex exponentials. Euler, the Bernouillis, and others had used such sums in similar fashions and for similar ends, but Fourier made a claim extravagant for the time, namely that *all functions* could be expressed in such terms. Unfortunately, in those days there was no clear idea of what a *function* was, no vocabulary to specify *classes* of functions, and no specification of what it would mean to *represent* a function by such a series.

[2] The notion of *has a Fourier expansion* would need to clarify what *has such an expansion* means. Must it mean that pointwise values can be retrieved from the Fourier series? Less? More?

[3] At about the time Fourier was promoting Fourier series, Abel proved that convergent power series *can* be differentiated term-by-term in the interior of their interval (on \mathbb{R}) or disk (in \mathbb{C}) of convergence, and *are* infinitely-differentiable functions. Abel's result fit the optimistic expectations of the time, but created unreasonable expectations for the behavior of Fourier series.

Several further issues are *implicit*, and the *best* answers need viewpoints created first in 1906 by Beppo Levi, 1907 by G. Frobenius, in the 1930's by Sobolev, and Schwartz post-1949, enabling legitimate discussion of *generalized functions* (a.k.a., *distributions*).^[4] There are natural technical questions, like

Why define generalized functions as dual spaces?

In brief, Schwartz' 1940's insight to define generalized functions as *dual spaces* is a natural consequence of one natural *relaxation* of the notion of *function*. Rather than demand that functions produce *pointwise values*, which precipitated endless classical discussion of what to do with jump discontinuities, instead declare that *functions* in the broadest sense are merely things that can be *integrated against*. For given φ , the map that integrates against it

$$f \longrightarrow \int f(x) \varphi(x) dx$$

is a *functional* (\mathbb{C} -valued linear map), and is, or ought to be, probably *continuous* in any reasonable topology. To consider the collection of *all* continuous linear functionals is a reasonable way to enlarge the collection of functions, as things to be integrated against.

From the other side, it might have been that this generalization of *function* is needlessly extravagant, but it turns out that every distribution is a high-order derivative of a continuous function. Thus, since we *do* want to be able to take derivatives indefinitely, there is no waste.

Further, in any of the several natural topologies on distributions, very nice ordinary functions are *dense*, and the space of distributions is *complete* in a sense subsuming that for metric spaces. Thus, taking limits *yields* all distributions, *and* produces no excess.

This discussion is easiest on the circle, or products of circles, making use of Fourier series, and clarifying many technical questions about Fourier series.^[5] This story is a prototype for more complicated examples.

There is an important auxiliary technical point. Natural spaces of functions *do not* have structures of Hilbert spaces, but typically, of Banach spaces. Nevertheless, the simplicity of Hilbert spaces motivates comparisons of natural function spaces with related Hilbert spaces. Such comparisons are *Levi-Sobolev imbeddings* or *Levi-Sobolev inequalities*.

1. Provocative example

Let $s(x)$ be the *sawtooth function*^[6]

$$s(x) = x - \pi \quad (\text{for } 0 \leq x < 2\pi)$$

and made *periodic* by demanding $s(x + 2\pi n) = s(x)$ for all $n \in \mathbb{Z}$. In other words, letting $\llbracket x/2\pi \rrbracket$ be the greatest integer less than or equal $x/2\pi$,

$$s(x) = x - 2\pi \cdot \left\llbracket \frac{x}{2\pi} \right\rrbracket - \pi \quad (\text{for } x \in \mathbb{R})$$

^[4] K. Friedrichs' important 1934-5 discussions of semi-bounded unbounded operators on Hilbert spaces used norms defined in terms of derivatives, but only internally in proofs, while for Levi, Frobenius, and Sobolev these norms were significant objects themselves.

^[5] The classic reference is A. Zygmund, *Trigonometric Series, I, II*, first published in Warsaw in 1935, reprinted several times, including a 1959 Cambridge University Press edition. The present discussion neglects many interesting details, *but* is readily adaptable to more complicated situations, so necessarily our treatment is different from Zygmund's.

^[6] One may also take $s(x) = x$ for $-\pi < x < \pi$ and extend by periodicity. This definition avoids the subtraction of π , and has the same operational features. In the end, it doesn't matter.

Away from $2\pi\mathbb{Z}$, the sawtooth function is infinitely differentiable, with derivative 1. At $x \in 2\pi\mathbb{Z}$ the sawtooth jumps down from value π to value $-\pi$. There is no reason to worry about defining a value at $x \in 2\pi\mathbb{Z}$.

The exponential functions $\psi_n(x) = e^{inx}$ are not quite an *orthonormal* basis for the Hilbert space $L^2[0, 2\pi]$, but are *orthogonal*:

$$\int_0^{2\pi} \psi_m(x) \cdot \overline{\psi_n(x)} dx = \begin{cases} 0 & (\text{for } m \neq n) \\ 2\pi & (\text{for } m = n) \end{cases}$$

Anticipating that Fourier coefficients $\widehat{f}(n)$ of $2\pi\mathbb{Z}$ -periodic functions f are computed^[7] by integrating against $\psi_n(x) = e^{inx}$ (conjugated):

$$\widehat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(x) e^{-inx} dx$$

integration by parts gives

$$\widehat{s}(n) = \frac{1}{2\pi} \int_0^{2\pi} s(x) \cdot e^{-inx} dx = \begin{cases} \frac{1}{-in} & (\text{for } n \neq 0) \\ 0 & (\text{for } n = 0) \end{cases}$$

Thus, in whatever sense a function *is* its Fourier expansion, we anticipate that

$$s(x) \sim \sum_{n \in \mathbb{Z}} \widehat{s}(n) \cdot e^{inx} = \sum_{n \neq 0} \frac{1}{-in} \cdot e^{inx}$$

Even though this series does not converge absolutely for *any* value of x , we will see below^[8] that it *does* converge to the value of $s(x)$ for $x \notin 2\pi\mathbb{Z}$. Since $s(x)$ has discontinuities at $2\pi\mathbb{Z}$ *anyway*, this is hardly surprising. Nothing disturbing has happened.

Now differentiate. The sawtooth function *is* differentiable away from $2\pi\mathbb{Z}$, with value 1, and with uncertain value at $2\pi\mathbb{Z}$. With exogenous reasons to differentiate the Fourier series term-by-term, with or without confidence in doing so, and the blatant differentiability of $s(x)$ away from $2\pi\mathbb{Z}$ suggests it's not entirely ridiculous to differentiate term-by-term. Then

$$s'(x) = \begin{cases} 1 & (\text{for } x \notin 2\pi\mathbb{Z}) \\ ? & (\text{for } x \in 2\pi\mathbb{Z}) \end{cases} \sim - \sum_{n \neq 0} e^{inx}$$

The right-hand side is hard to interpret, certainly as having pointwise values. On the other hand, reasonably interpreted, it is still ok to integrate against this sum: letting $\widehat{f}(n)$ be the n^{th} Fourier coefficient of a *smooth* function f , and not worrying about justifications,

$$\begin{aligned} \int_0^{2\pi} f(x) \left(- \sum_{n \neq 0} e^{inx} \right) dx &= - \sum_{n \neq 0} \int_0^{2\pi} f(x) e^{inx} dx = -2\pi \sum_{n \neq 0} \widehat{f}(-n) \\ &= 2\pi \widehat{f}(0) - 2\pi \sum_{n \in \mathbb{Z}} \widehat{f}(n) e^{in \cdot 0} = 2\pi \widehat{f}(0) - 2\pi f(0) = \int_0^{2\pi} f(x) dx - 2\pi \cdot f(0) \end{aligned}$$

[7] Apparently at first Fourier did not have this expression for the Fourier coefficients!

[8] We will prove that at points where *left and right derivatives* exist, *piecewise continuous* functions' Fourier series do converge to their pointwise values.

The map

$$f \longrightarrow \int_0^{2\pi} f(x) dx - 2\pi \cdot f(0)$$

has a sense for continuous f , and gives a *functional*. That the derivative of the sawtooth is *mostly* 1 gives the integral of f (against 1) over $[0, 2\pi]$. Further, the $-2\pi f(0)$ term forcefully suggests that *the derivative of the discontinuity of the sawtooth function is the (periodic) evaluation-at-0 functional $f \rightarrow f(0)$ multiplied by -2π* . [9]

[1.1.1] **Remark:** A truly disastrous choice at this point would be to think that since $s'(x)$ is *almost everywhere* 1 (in a measure-theoretic sense) that its singularities are somehow *removable*, and thus pretend that $s'(x) = 1$. This would give $s''(x) = 0$, and make the following worse than it is, and impossible to explain.

Still, $s'(x)$ is differentiable away from $2\pi\mathbb{Z}$, and by repeated differentiation

$$s^{(k+1)}(x) = \begin{cases} 0 & (\text{for } x \notin 2\pi\mathbb{Z}) \\ ? & (\text{for } x \in 2\pi\mathbb{Z}) \end{cases} \sim -(i)^k \sum_{n \neq 0} n^k \cdot e^{inx}$$

By now the right-hand sides are vividly not convergent. The summands do not go to zero, in fact, are *unbounded*.

One can continue differentiating in this symbolic sense, but the meaning is unclear.

One reaction is to simply object to differentiating a non-differentiable function, even if its discontinuities are mild. This is not productive.

Another unproductive viewpoint is to deny that Fourier series reliably represent the functions that produced their coefficients.

A happier and more useful response is to suspect that the above computation is *correct*, though the question mark needs explanation, *and* that the right-hand side is correct and meaningful, *despite* its divergence in classical senses. The question is *what* meaning to attach. This requires preparation.

We will establish a context in which the derivatives of the sawtooth, and derivatives of other discontinuous functions, are *things to integrate against*, rather than *things to evaluate pointwise*, and see that termwise differentiation of Fourier series *does* capture an extended notion of function and derivative.

2. Natural function spaces on the circle $S^1 = \mathbb{R}/2\pi\mathbb{Z}$

We review natural families of functions. In all cases, the object is to give the vector space of functions a metric (if possible) which makes it *complete*, to allow *taking limits* inside the same class of functions. For example, *pointwise* limits of continuous functions easily fail to be continuous. [10]

[2.1] Continuous functions and sup-norm

First, we care about *continuous* complex-valued functions. Although we have in mind continuous functions on the circle $S^1 = \mathbb{R}/2\pi\mathbb{Z}$, the basic result depends only upon the *compactness* of $\mathbb{R}/2\pi\mathbb{Z}$.

[9] The *jump* is downward rather than upward.

[10] Awareness of such possibilities and figuring out how to avoid them was the fruit of embarrassing errors and experimentation throughout the 19th century. Unifying abstract notions such as *metric space* and general *topological space* only became available in the early 20th century, with the work of Hausdorff, Fréchet, and others.

We give the set $C^o(K)$ of (complex-valued) continuous functions on a compact topological space K the metric

$$d(f, g) = \sup_{x \in K} |f(x) - g(x)|$$

The sup is *finite* because K is compact and $f - g$ is continuous. ^[11] The right-hand side of this last equation arises from the (*sup*) *norm* ^[12]

$$|f|_\infty = |f|_{C^o} = \sup_{x \in K} |f(x)|$$

and $d(f, g) = |f - g|_{C^o}$. A main feature of continuous functions is that they have *pointwise values*. We have the unsurprising but important

[2.1.1] Claim: With the $C^o(K)$ topology, for $x \in K$ the *evaluation functional* ^[13] $C^o(K) \rightarrow \mathbb{C}$ by $f \rightarrow f(x)$ is *continuous*.

Proof: The inequality

$$|f(x) - g(x)| \leq \sup_{y \in K} |f(y) - g(y)| \quad (\text{for } f, g \in C^o(K))$$

proves the continuity of evaluation. ///

[2.1.2] Theorem: The space $C^o(K)$ of (complex-valued) continuous functions on a compact topological space K is *complete*. ^[14]

[2.1.3] Remark: Thus, being complete with respect to the metric arising in this fashion from a *norm*, by definition $C^o(K)$ is a *Banach space*.

Proof: This is a typical three-epsilon argument. The point is the *completeness*, namely that a Cauchy sequence of continuous functions has a *pointwise* limit which is a continuous function. First we observe that a Cauchy sequence f_i does have a pointwise limit. Given $\varepsilon > 0$, choose N large enough such that for $i, j \geq N$ we have $|f_i - f_j| < \varepsilon$. Then, for any x in K , $|f_i(x) - f_j(x)| < \varepsilon$. Thus, the sequence of values $f_i(x)$ is a Cauchy sequence of complex numbers, so has a limit $f(x)$. Further, given $\varepsilon' > 0$, choose $j \geq N$ sufficiently large such that $|f_j(x) - f(x)| < \varepsilon'$. Then for all $i \geq N$

$$|f_i(x) - f(x)| \leq |f_i(x) - f_j(x)| + |f_j(x) - f(x)| < \varepsilon + \varepsilon'$$

Since this is true for every positive ε'

$$|f_i(x) - f(x)| \leq \varepsilon \quad (\text{for all } i \geq N)$$

^[11] Recall how to prove that a continuous function f on a compact set K is *bounded*: let U_n be $\{x \in K : |f(x)| < n\}$, for $n = 1, 2, \dots$. The union of these open U_n 's is all of K . By compactness, there is a finite subcover, proving the boundedness. But it's not trivial to prove that closed intervals $[a, b]$ are compact in this sense.

^[12] Recall that a *norm* $v \rightarrow |v|$ on a complex vector space V is a non-negative real-valued function $v \rightarrow |v|$ which is *positive* (meaning that $|v| = 0$ only for $v = 0$), *homogeneous* (meaning that $|\alpha \cdot v| = |\alpha|_{\mathbb{C}} \cdot |v|$ for complex α , where $|\alpha|_{\mathbb{C}}$ is the usual complex absolute value), and satisfies the *triangle inequality* (that $|v + w| \leq |v| + |w|$). The first two properties are readily verified for the sup norm, and the triangle inequality follows from the readily verifiable fact that the *sup of the sum* is less than or equal the *sum of the sups*.

^[13] As usual, a (*continuous*) *functional* is a (continuous) linear map to \mathbb{C} .

^[14] There are many not-continuous *pointwise* limits of continuous functions, but these are not *uniform* pointwise limits. In the early 19th century the distinction between the two notions was not clearly established, precipitating technical confusion.

This holds for every x in K , so the pointwise limit is uniform in x .

Now prove that $f(x)$ is continuous. Given $\varepsilon > 0$, let N be large enough so that for $i, j \geq N$ we have $|f_i - f_j| < \varepsilon$. From the previous paragraph

$$|f_i(x) - f(x)| \leq \varepsilon \quad (\text{for every } x \text{ and for } i \geq N)$$

Fix $i \geq N$ and $x \in K$, and choose a small enough neighborhood U of x such that $|f_i(x) - f_i(y)| < \varepsilon$ for any y in U . Then

$$|f(x) - f(y)| \leq |f(x) - f_i(x)| + |f_i(x) - f_i(y)| + |f(y) - f_i(y)| < \varepsilon + \varepsilon + \varepsilon$$

Thus, the pointwise limit f is continuous at every x in U . ///

[2.2] Differentiation on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$

To talk about *differentiability* return to the concrete situation of \mathbb{R} and its quotient $S^1 = \mathbb{R}/2\pi\mathbb{Z}$.

The continuous quotient map $q : \mathbb{R} \rightarrow \mathbb{R}/2\pi\mathbb{Z}$ yields continuous functions under composition $f \circ q$ for $f \in C^0(S^1) = C^0(\mathbb{R}/2\pi\mathbb{Z})$. More is true, namely, that a continuous function F on \mathbb{R} is of the form $f \circ q$ if and only if F is *periodic* in the sense that $F(x + 2\pi n) = F(x)$ for all $x \in \mathbb{R}$ and $n \in \mathbb{Z}$. Indeed, the periodicity gives a *well-defined* function f on $\mathbb{R}/2\pi\mathbb{Z}$. Then the continuity of f follows immediately from the definition of the quotient topology on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$.

As usual, a real-valued or complex-valued function f on \mathbb{R} is *continuously differentiable* if it has a derivative itself a continuous function. That is, the limit

$$\frac{df}{dx}(x) = f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

is required to exist for all x , and the function f' is in $C^0(\mathbb{R})$. Let $f^{(1)} = f'$, and inductively define

$$f^{(i)} = \left(f^{(i-1)}\right)' \quad (\text{for } i > 1)$$

when the corresponding limits exist.

We must make explicit our expectation that *differentiation* on the circle $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ is descended from differentiation on the real line. That is, *characterize* differentiation on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ in terms of such a compatibility relation. Thus, for $f \in C^k(S^1)$, require that the differentiation D on S^1 be related to the differentiation on \mathbb{R} by

$$(Df) \circ q = \frac{d}{dx}(f \circ q)$$

Via the quotient map $q : \mathbb{R} \rightarrow \mathbb{R}/2\pi\mathbb{Z}$, make a *preliminary* definition of the collection of k -times continuously differentiable functions on S^1 , with a topology, by

$$C^k(S^1) = \{f \text{ on } S^1 : f \circ q \in C^k(\mathbb{R})\}$$

with the C^k -norm^[15]

$$|f|_{C^k} = \sum_{0 \leq i \leq k} |(f \circ q)^{(i)}|_{\infty} = \sum_{0 \leq i \leq k} \sup_x |(f \circ q)^{(i)}(x)|$$

[15] Granting that the sup norm on *continuous* functions is a norm, verification that the C^k -norm is a norm is straightforward.

where $F^{(i)}$ is the (continuous!) i^{th} derivative of F on \mathbb{R} . The associated metric on $C^k(S^1)$ is

$$d(f, g) = \|f - g\|_{C^k}$$

[2.2.1] **Remark:** Among other features, the norm on the spaces C^k makes continuity of the *differentiation* map $C^k \rightarrow C^{k-1}$ completely obvious.

[2.2.2] **Remark:** Implicit in this definition is that, viewed as functions on $[0, 2\pi]$, the values and derivatives *must agree at the endpoints*: $f(0) = f(2\pi)$ for f continuous on S^1 , $f'(0) = f'(2\pi)$ for $f \in C^1(S^1)$, and so on. This is not whimsical, but is intrinsic to the structure of S^1 .

An often-seen equivalent version of the norm is

$$\|f\|_{C^k}^{\text{var}} = \sup_{0 \leq i \leq k} \|(f \circ q)^{(i)}\|_{\infty} = \sup_{0 \leq i \leq k} \sup_x |(f \circ q)^{(i)}(x)|$$

These two norms give the same topology, since for complex numbers a_0, \dots, a_k

$$\sup_{0 \leq i \leq k} |a_i| \leq \sum_{0 \leq i \leq k} |a_i| \leq (k+1) \cdot \sup_{0 \leq i \leq k} |a_i|$$

[2.2.3] **Claim:** There is a unique, well-defined, continuous (differentiation) map $D : C^k(S^1) \rightarrow C^{k-1}(S^1)$ giving a commutative diagram

$$\begin{array}{ccc} C^k(\mathbb{R}) & \xrightarrow{d/dx} & C^{k-1}(\mathbb{R}) \\ \uparrow -\circ q & & \uparrow -\circ q \\ C^k(S^1) & \xrightarrow{D} & C^{k-1}(S^1) \end{array}$$

[2.2.4] **Remark:** One might feel that this proof is needlessly complicated. However, it is worthwhile to do it this way. This approach applies broadly, *and* is as terse as possible without ignoring important details.

Proof: The point is that differentiation of periodic functions yields periodic functions. That is, we claim that, for $f \in C^k(S^1)$, the pullback $f \circ q$ has derivative $\frac{d}{dx}(f \circ q)$ which is the pullback $g \circ q$ of a unique function $g \in C^{k-1}(S^1)$. To see this, first recall that, by definition of the quotient topology, a continuous function F on \mathbb{R} descends to a continuous function on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ if and only if it is $2\pi\mathbb{Z}$ -invariant, that is $F(x + 2\pi n) = F(x)$ for all $x \in \mathbb{R}$ and $n \in \mathbb{Z}$. Then, from our definition of $C^k(S^1)$, a function $F \in C^k(\mathbb{R})$ is a pullback via q from $C^k(\mathbb{R}/2\pi\mathbb{Z})$ exactly when $F^{(i)}(x + 2\pi n) = F^{(i)}(x)$ for all $x \in \mathbb{R}$, $n \in \mathbb{Z}$, and $0 \leq i \leq k$, since then these continuous functions descend to the circle. Let

$$(T_y F)(x) = F(x + y) \quad (\text{for } x, y \in \mathbb{R})$$

Since $\frac{d}{dx}$ is a linear, constant-coefficient differential operator, the operations T_y and $\frac{d}{dx}$ commute, that is, $\frac{\partial F}{\partial x}(x + y) = \frac{\partial}{\partial x}(F(x + y))$, which is to say

$$T_y \circ \frac{d}{dx} = \frac{d}{dx} \circ T_y$$

In particular, for $n \in \mathbb{Z}$,

$$T_{2\pi n} \left(\frac{d}{dx}(f \circ q) \right) = \frac{d}{dx}(T_{2\pi n}(f \circ q)) = \frac{d}{dx}(f \circ q)$$

This shows that a (continuous) derivative is periodic when the (continuously differentiable) function is periodic.

From the definition of the C^k -norm,

$$|Df|_{C^{k-1}} \leq |f|_{C^k}$$

so differentiation is continuous. ///

[2.2.5] **Remark:** In light of the uniqueness of differentiation on S^1 , from now on write d/dx for the differentiation D on S^1 , and $f^{(k)}$ for $D^k f$. Now rewrite the description of $C^k(S^1)$ more simply, as

$$C^k(S^1) = \{f \text{ on } S^1 : f \circ q \in C^k(\mathbb{R})\}$$

with the C^k -norm

$$|f|_{C^k} = \sum_{0 \leq i \leq k} |f^{(i)}|_{\infty} = \sum_{0 \leq i \leq k} \sup_x |f^{(i)}(x)|$$

where $f^{(i)}$ is the (continuous!) i^{th} derivative of f . The *associated metric* on $C^k(S^1)$ still is

$$d(f, g) = |f - g|_{C^k}$$

There is the alternative norm

$$|f|_{C^k}^{\text{var}} = \sup_{0 \leq i \leq k} \sup_x |f^{(i)}(x)| = \sup_{0 \leq i \leq k} |f^{(i)}|_{\infty}$$

Again, these two norms give the same topology, for the same reason as before.

[2.2.6] **Claim:** With the topology above, the space $C^k(S^1)$ is *complete*, so is a *Banach* space.

Proof: The case $k = 1$ illustrates all the points. For a Cauchy sequence $\{f_n\}$ in $C^1(S^1)$, both $\{f_n\}$ and $\{f'_n\}$ are Cauchy in $C^0(S^1)$, so converge uniformly pointwise: let

$$f(x) = \lim_n f_n(x) \quad g(x) = \lim_n f'_n(x)$$

The convergence is uniformly pointwise, so f and g are C^0 . If we knew that f were pointwise differentiable, then the demonstrated continuity of $\frac{d}{dx} : C^1(S^1) \rightarrow C^0(S^1)$ gives the expected conclusion, that $f' = g$.

What could go wrong? The issue is whether f is *differentiable!*

By the fundamental theorem of calculus, for any index i , since f_i is continuous, [16]

$$f_i(x) - f_i(a) = \int_a^x f'_i(t) dt$$

Interchanging limit and integral [17] shows that the limit of the right-hand side is

$$\lim_i \int_a^x f'_i(t) dt = \int_a^x \lim_i f'_i(t) dt = \int_a^x g(t) dt$$

[16] The fundamental theorem of calculus for integrals of *continuous* functions needs only the simplest notion of an integral.

[17] For example, interchange of limit and integral is justified by the simplest form of Lebesgue's Dominated Convergence Theorem. It may seem to be overkill to invoke the Dominated Convergence Theorem in this context, but systematic attention to such details helps us avoid many of the gaffes of early 19th-century analysis. Lebesgue's deserved fame is due to his overcoming the murkiness of earlier theories of integration.

Thus, the limit of the left-hand side is

$$f(x) - f(a) = \int_a^x g(t) dt$$

from which $f' = g$. That the derivative f' of the limit f is the limit of the derivatives is not a surprise, since if f is differentiable, what else could its derivative be? The point is that f is differentiable, ascertained by computing its derivative, which happens to be g . ///

[2.2.7] **Remark:** Again, the differentiation map $C^1(S^1) \rightarrow C^0(S^1)$ is continuous *by design*. Thus, if a limit of C^1 functions f_n is differentiable, its derivative must be the obvious thing, namely, the limit of the derivatives f'_n . The issue was whether the limit of the f_n is differentiable. The proof shows that it is differentiable by computing its derivative via the Mean Value Theorem.

By construction, and from the corresponding result for C^0 ,

[2.2.8] **Claim:** With the C^k -topology, for $x \in S^1$ and integer $0 \leq i \leq k$, the *evaluation functional* $C^k(S^1) \rightarrow \mathbb{C}$ by

$$f \longrightarrow f^{(i)}(x)$$

is *continuous*. ///

This applies to Fourier series, without any claim about what functions are representable as Fourier series. With $\psi_n(x) = e^{inx}$,

[2.2.9] **Claim:** For complex numbers c_n , when

$$\sum_n |c_n| \cdot |n|^k < +\infty$$

the Fourier series $\sum c_n \psi_n$ converges to a function in $C^k(S^1)$, and its derivative is computed by termwise differentiation

$$\frac{d}{dx} \sum c_n \psi_n = \sum (in) c_n \psi_n \in C^{k-1}(S^1)$$

Proof: The $C^0(S^1)$ norm of a Fourier series is easily estimated, by

$$\left| \sum_{|n| \leq N} c_n \psi_n(x) \right| \leq \sum_{|n| \leq N} |c_n| \quad (\text{for all } x \in S^1)$$

The right-hand side is independent of $x \in S^1$, so bounds the sup over $x \in S^1$. Similarly, estimate derivatives (of partial sums) by

$$\left| \left(\sum_{|n| \leq N} c_n \psi_n \right)^{(k)} \right| \leq \sum_{|n| \leq N} |c_n| n^k$$

Thus, the hypothesis of the claim implies that the partial sums form a Cauchy sequence in C^k . The partial sums of a Fourier series are *finite* sums, so can be differentiated term-by-term. Thus, we have a Cauchy sequence of C^k functions, which converges to a C^k function, by the completeness of C^k . That is, the given estimate assures that the Fourier series converges to a C^k function.

Further, since differentiation is a continuous map $C^k \rightarrow C^{k-1}$, it maps Cauchy sequences to Cauchy sequences. In particular, the Cauchy sequence of derivatives of partial sums converges to the derivative of the limit of the original Cauchy sequence. ///

We want the following to hold. Unsurprisingly, it does:

[2.2.10] Claim: The inclusion $C^k(S^1) \subset C^{k-1}(S^1)$ is continuous. [18]

Proof: The point is that, for $f \in C^k(S^1)$ the obvious inequality

$$\|f\|_{C^{k-1}} \leq \|f\|_{C^k}$$

gives an explicit estimate for the continuity. ///

3. Topology on $C^\infty(S^1)$

Next, we care about infinitely differentiable [19] functions, that is, *smooth* functions, denoted $C^\infty(S^1)$. At least as *sets* (or vector spaces),

$$C^\infty(S^1) = \bigcap_k C^k(S^1)$$

However, this space $C^\infty(S^1)$ of smooth functions provably *does not* have a structure of Banach space. Observing that a descending intersection is a *(projective) limit* [20] we should declare that

$$C^\infty(S^1) = \lim_k C^k(S^1)$$

[3.1.1] Remark: Another natural space of functions is $C_c^\circ(\mathbb{R})$, the space of compactly-supported continuous functions on \mathbb{R} . Since (as a set)

$$C_c^\circ(\mathbb{R}) = \bigcup_N \{f \in C_c^\circ(\mathbb{R}) : \text{support } f \subset [-N, +N]\} = \text{colim}_N \{f \in C_c^\circ(\mathbb{R}) : \text{support } f \subset [-N, +N]\}$$

$C_c^\circ(\mathbb{R})$ is an *ascending union* (colimit) of Banach spaces. [21] Colimits of Banach spaces are even less likely to be Banach spaces than are limits. In this example, the colimit is not even *metrizable*, since, as we'll see later, a topology in which it is suitably *complete* makes it violate the conclusion of Baire's theorem: $C_c^\circ(\mathbb{R})$ is a countable union of nowhere dense subsets. [22] We will address this when looking at functions on \mathbb{R} .

Returning to $C^\infty(S^1) = \lim_k C^k(S^1)$, unfortunately we are temporarily insufficiently sophisticated about what kind of object this limit might be. In particular, we do not know what kind of auxiliary objects to use

[18] In fact, the image of C^k in C^{k-1} is *dense*, but, we will prove this later as a side-effect of sharper results.

[19] Use of *infinitely* here is potentially misleading, but is standard. Sometimes the phrase *indefinitely differentiable* is used, but this also offers its own potential for confusion. A better (and standard) contemporary usage is *smooth*.

[20] Although descending intersections are (projective) limits, with *projections* being the natural *inclusions*, not every limit should be construed as a descending intersection. Indeed, solenoids cannot usefully be construed as descending intersections. The notion of limit is broader than either of these. Similarly, ascending unions are *colimits*, but there are much more general types of colimits.

[21] That the limitands are Banach spaces is easy: first, the set of continuous functions with support in $[-N, N]$ is the subspace of the Banach space $C^\circ[-N, N]$ defined by $f(-N) = f(N) = 0$. These evaluation functionals are continuous, so this is a *closed* subspace of a Banach space, and is Banach. Since $[-N, N]$ is *compact*, the space $C^\circ[-N, N]$ with the sup norm is Banach, as we saw earlier.

[22] The Baire Category Theorem asserts that a complete metric space cannot be a countable union of nowhere dense subsets. A nowhere dense subset is one whose closure contains no non-empty open sets.

in the very definition of *limit*. By now we know that limits of *topological spaces* exist (and are necessarily unique), but we need more structure. To explain this requires a general notion of *topological vector space*.

[3.1.2] **Remark:** Too optimistic speculation about what the limit might be leads to trouble: as it happens, this limit is provably *not* a Banach space (nor Hilbert space).^[23] A limit of topological spaces has a unique topology, whatever it may be.

We anticipate that a limit has a vector space structure compatible with the limit's topology.

[3.1.3] **Remark:** There is the disquieting question of what *test objects* Z we should consider in the diagrammatic characterization, with compatible mappings $Z \rightarrow C^k(S^1)$ to *define* the limit. Similarly, we should wonder how large a class of vector spaces with topologies is needed. That is, *existence* is perhaps non-trivial.

The broadest necessary class of vector spaces with topologies is the following. A *topological vector space* is what one would reasonably imagine, namely, a (complex) vector space V with a topology such that

$$V \times V \rightarrow V \quad \text{by} \quad v \times w \rightarrow v + w \quad \text{is continuous}$$

and such that

$$\mathbb{C} \times V \rightarrow V \quad \text{by} \quad \alpha \times v \rightarrow \alpha \cdot v \quad \text{is continuous}$$

and such that the topology is *Hausdorff*.^[24] We require that the topological vector spaces be *locally convex* in the sense that there is a local basis at 0 consisting of *convex* sets.^[25] It is easy to prove that Hilbert and Banach spaces are locally convex, which is why the issue is invisible in that context. Dismayingly, there are easily constructed complete (invariantly) metrized topological vector spaces which are *not* locally convex.^[26]

Returning to the discussion of limits of topological vector spaces: since the continuity requirements for a topological vector space are of the form $A \times B \rightarrow C$ (rather than having the arrow going the other direction), there is a *diagrammatic* argument that the continuous algebraic operations on the limitands induce continuous algebraic operations on the limit, in the limit topology (as limit of topological spaces).

[23] The non-Banach-ness of $C^\infty(S^1)$ is not the main point, but it is reasonable to wonder how this is proven. Briefly, with a definition of *topological vector space*, we will prove that a topological vector space is *normable* if and only if there is a local basis at 0 consisting of *bounded* opens. This is independent of *completeness*. The relevant sense of bounded *cannot* be the usual metric sense. Instead, a set E in a topological vector space is *bounded* when, for every open neighborhood U of 0, there is $t > 0$ such that $E \subset z \cdot U$ for all complex z with $|z| \geq t$. That is, sufficiently large *dilates* of opens eventually contain E . But we will eventually that open balls in $C^k(S^1)$ are *not* contained in *any* dilate of any open ball in $C^{k+1}(S^1)$. The definition of the limit topology then shows that $C^\infty(S^1)$ is not normable. A more detailed discussion will be given later.

[24] In fact, soon after giving the definition, one can show that the weaker condition that *points are closed*, implies the Hausdorff condition in topological spaces which are vector spaces with continuous vector addition and scalar multiplication. Indeed, the inverse image of $\{0\}$ under $x \times y \rightarrow x - y$ is the diagonal.

[25] This sense of convexity is the usual: a set X in a vector space is convex when, for all tuples x_1, \dots, x_n of points in X and all tuples t_1, \dots, t_n of non-negative reals with $\sum_i t_i = 1$, the sum $\sum_i t_i x_i$ is again in X .

[26] The simplest examples of complete metric topological vector spaces which are *not* locally convex are spaces ℓ^p with $0 < p < 1$. The metric comes from a norm-like function which is *not* a norm: $\|\{c_n\}\|_p = \sum_n |c_n|^p$. No, there is no p^{th} root taken, unlike the spaces ℓ^p with $p \geq 1$, and this causes the function $\|\cdot\|_p$ to *lose* the homogeneity it would need to be a norm. Nevertheless, such a space is *complete*. It is an amusing exercise to prove that it is not locally convex.

[3.1.4] **Claim:** Products and limits of topological vector spaces exist. Products and limits of locally convex spaces are locally convex. (Proof in appendix.)

[3.1.5] **Remark:** As usual, if they exist at all, then products and limits are unique up to unique isomorphism.

[3.1.6] **Remark:** Coproducts and colimits are more fragile, and will be treated, as needed, in discussion of function theory on non-compact spaces such as \mathbb{R} .

Thus, $C^\infty(S^1)$ has a (limit) topology for general reasons.

[3.1.7] **Claim:** Differentiation $f \rightarrow f'$ is a *continuous* map $C^\infty(S^1) \rightarrow C^\infty(S^1)$.

[3.1.8] **Remark:** *Of course* differentiation maps the smooth functions to themselves. Continuity of differentiation in the *limit* topology is less clear.

Proof: We already know that differentiation d/dx gives a continuous map $C^k(S^1) \rightarrow C^{k-1}(S^1)$. Differentiation is compatible with the inclusions among the $C^k(S^1)$. Thus, we have a commutative diagram

$$\begin{array}{ccccccc}
 C^\infty(S^1) & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & C^k(S^1) & \xrightarrow{\quad} & C^{k-1}(S^1) \xrightarrow{\quad} \dots \\
 & & & & \nearrow \frac{d}{dx} & & \nearrow \frac{d}{dx} \\
 C^\infty(S^1) & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & C^k(S^1) & \xrightarrow{\quad} & C^{k-1}(S^1) \xrightarrow{\quad} \dots
 \end{array}$$

Composing the projections with d/dx gives (dashed) induced maps from $C^\infty(S^1)$ to the limitands, inducing a unique (dotted) map to the limit, as in

$$\begin{array}{ccccccc}
 C^\infty(S^1) & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & C^k(S^1) & \xrightarrow{\quad} & C^{k-1}(S^1) \xrightarrow{\quad} \dots \\
 \uparrow \frac{d}{dx} & & & & \nearrow \text{dashed} & & \nearrow \text{dashed} \\
 C^\infty(S^1) & \xrightarrow{\quad} & \dots & \xrightarrow{\quad} & C^k(S^1) & \xrightarrow{\quad} & C^{k-1}(S^1) \xrightarrow{\quad} \dots \\
 & & & & \nearrow \text{dotted} & & \nearrow \text{dotted}
 \end{array}$$

This proves the continuity of differentiation, in the limit topology. ///

[3.1.9] **Corollary:** When a Fourier series $\sum_n c_n \psi_n$ satisfies

$$\sum_m |c_n| |n|^N < +\infty \quad (\text{for every } N)$$

the series is a smooth function, which can be differentiated term-by-term, and its derivative is

$$\sum_m c_n (in) \psi_n$$

Proof: The hypothesis assures that the Fourier series lies in C^k for every k . Differentiation is continuous in the limit topology on C^∞ . ///

[3.1.10] **Remark:** This continuity is necessary to define differentiation of *distributions* below.

4. Pointwise convergence of Fourier series

Before proceeding further, a classical criterion for pointwise convergence of Fourier series is comforting.

We can prove pointwise convergence even before proving that the exponentials give an orthonormal *basis* for $L^2[0, 2\pi]$. The hypotheses of the convergence claim below are far from optimal, but are sufficient for some purposes, and are tangibly verifiable.

The only slightly subtle thing needed is

[4.1.1] Claim: (*Riemann-Lebesgue*) For $f \in L^2(S^1)$, the Fourier coefficients $\widehat{f}(n)$ of f go to 0.

Proof: The L^2 norm of ψ_n is $\sqrt{2\pi}$. *Bessel's inequality*^[27]

$$\|f\|_{L^2}^2 \geq \sum_n \left| \left\langle f, \frac{\psi_n}{\sqrt{2\pi}} \right\rangle \right|^2$$

from abstract Hilbert-space theory applies to an orthonormal *set*, whether or not it is an orthonormal *basis*. Thus, the sum on the right converges, so by Cauchy's criterion the summands go to 0. ///

A function f on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ is (*finitely*) *piecewise C^o* when there are finitely many real numbers $a_0 \leq a_1 \leq \dots \leq a_{n-1} \leq a_n = a_0 + 1$ and C^0 functions f_i on $[a_i, a_{i+1}]$ such that

$$f_i(x) = f(x) \quad \text{on } [a_i, a_{i+1}] \quad (\text{except possibly at the endpoints})$$

Thus, while $f_i(a_{i+1})$ may differ from $f_{i+1}(a_{i+1})$, and $f(a_{i+1})$ may be different from both of these, the function f is continuous in the interiors of the intervals, and behaves well *near* the endpoints, if not *at* the endpoints.

Write

$$\langle f, F \rangle = \int_0^{2\pi} f(x) \overline{F(x)} dx$$

and

$$\widehat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(x) \overline{\psi_n(x)} dx = \frac{1}{2\pi} \langle f, \psi_n \rangle$$

[4.1.2] Claim: Let f be piecewise C^o on S^1 . Let x_o be a point at which f has both *left and right derivatives* (even if they do not agree), and is *continuous*. Then the Fourier series of f evaluated at x_o converges to $f(x_o)$. That is,

$$f(x_o) = \sum_{n \in \mathbb{Z}} \widehat{f}(n) \psi_n(x_o) \quad (\text{a convergent sum})$$

[4.1.3] Remark: The hypotheses of the claim are designed to be indifferent to the requirement that values and derivatives at endpoints of $[0, 2\pi]$ agree. We certainly will require this matching subsequently, but this preliminary result is useful.

^[27] Proof of Bessel's inequality is straightforward: for *finite* orthonormal set $\{e_i\}$ and a vector v in a Hilbert space,

$$0 \leq \|v - \sum_i \langle v, e_i \rangle e_i\|^2 = \|v\|^2 - 2 \sum_i \langle v, e_i \rangle \overline{\langle v, e_i \rangle} + \sum_i |\langle v, e_i \rangle|^2 = \|v\|^2 - \sum_i |\langle v, e_i \rangle|^2$$

For an *arbitrary* orthonormal set, the sum is the sup of the finite sub-sums, so the general Bessel inequality follows.

Proof: First, make reductions to unclutter the notation. By considering $f(x) - f(x_o)$, and observing that *constants* are represented pointwise by their Fourier expansions, we can assume that $f(x_o) = 0$. The Fourier coefficients of *translates* of a function are expressible in terms of the Fourier coefficients of the function itself:

$$\int_0^{2\pi} f(x + x_o) \bar{\psi}_n(x) dx = \int_0^{2\pi} f(x) \bar{\psi}_n(x - x_o) dx = \psi_n(x_o) \int_0^{2\pi} f(x) \bar{\psi}_n(x) dx$$

The left-hand side is 2π times the n^{th} Fourier coefficient of $f(x + x_o)$, that is, the n^{th} Fourier *term* of $f(x + x_o)$ evaluated at 0, while the right-hand side is 2π times the n^{th} Fourier *term* of $f(x)$ evaluated at x_o . Thus, simplify further by taking $x_o = 0$, without loss of generality.

A partial sum of the Fourier expansion evaluated at 0 is

$$\begin{aligned} \sum_{-M \leq n < N} \frac{1}{2\pi} \int_0^{2\pi} f(x) \bar{\psi}_n(x) dx &= \frac{1}{2\pi} \int_0^{2\pi} f(x) \sum_{-M \leq n < N} \bar{\psi}_n(x) dx \\ &= \frac{1}{2\pi} \int_0^{2\pi} \frac{f(x)}{\psi_{-1}(x) - 1} (\bar{\psi}_N(x) - \bar{\psi}_{-M}(x)) dx = \frac{1}{2\pi} \left\langle \frac{f}{\psi_{-1} - 1}, \psi_N \right\rangle - \frac{1}{2\pi} \left\langle \frac{f}{\psi_{-1} - 1}, \psi_{-M} \right\rangle \end{aligned}$$

The latter two terms are Fourier coefficients of $f/(\psi_{-1} - 1)$, so go to 0 by the Riemann-Lebesgue lemma for $f(x)/(\psi_{-1}(x) - 1)$ in $L^2(S^1)$. Since $x_o = 0$ and $f(x_o) = 0$

$$\frac{f(x)}{\psi_{-1}(x) - 1} = \frac{f(x)}{x} \cdot \frac{x}{\psi_{-1}(x) - 1} = \frac{f(x) - f(x_o)}{x - x_o} \cdot \frac{x - x_o}{e^{-ix} - e^{-ix_o}}$$

The existence of left and right derivatives of f at $x_o = 0$ is exactly the hypothesis that this expression has left and right limits at x_o , even if they do not agree.

At all other points the division by $\psi_{-1}(x) - 1$ does not disturb the continuity. Thus, $f/(\psi_{-1} - 1)$ is still at least *continuous* on each interval $[a_i, a_{i+1}]$ on which f was essentially a C^o function. Therefore, ignoring the endpoints, which do not contribute to the integrals, $f/(\psi_{-1} - 1)$ is continuous on a finite set of closed (finite) intervals, so bounded on each one. Thus, $f/(\psi_{-1} - 1)$ is indeed L^2 , and we can invoke Riemann-Lebesgue to see that the integral goes to $0 = f(x_o)$. ///

[4.1.4] **Corollary:** The Fourier series of a function in $C^1(S^1)$ converges pointwise to the function's values. ///

[4.1.5] **Remark:** Pointwise convergence is not L^2 convergence, and we have *not* yet proven that the exponentials are an orthonormal *basis* for the Hilbert space $L^2(S^1)$. The pointwise result just proven is suggestive, but not decisive.

5. Distributions: generalized functions

Although much amplification is needed, a topology on $C^\infty(S^1)$ allows the bare definition: a *distribution* or *generalized function*^[28] on S^1 is a *continuous linear functional*^[29]

$$u : C^\infty(S^1) \longrightarrow \mathbb{C}$$

[28] What's in a name? In this case, *generalized function* expresses the *intention* to think of distributions as extensions of ordinary functions, not as abstract things in a dual space.

[29] The standard usage is that a *functional* on a complex vector space V is a \mathbb{C} -linear map from V to \mathbb{C} . Continuity may or may not be required, and the topology in which continuity is required may vary. It is in this sense that there is a subject *functional analysis*.

Why a dual space? Unsurprisingly, as soon as we have a good *intrinsic* notion of *integral* on S^1 in the next section, a function $\varphi \in C^o(S^1)$ gives rise to a distribution u_φ by *integration against* φ ,

$$u_\varphi(f) = \int_{S^1} f(x) \varphi(x) dx \quad (\text{for } f \in C^\infty(S^1))$$

Thus, we relax our notion of *function*, no longer requiring *pointwise values*, but only that a function can be *integrated against*. Then it may make sense to declare functionals in a dual space to be generalized functions. The vector space of distributions is denoted

$$\text{distributions} = \text{continuous dual of } C^\infty(S^1) = \text{Hom}_{\mathbb{C}}^o(C^\infty(S^1), \mathbb{C}) = C^\infty(S^1)^*$$

That is, given a reasonable notion of integral, we have a continuous imbedding

$$C^o(S^1) \subset C^\infty(S^1)^* \quad \text{by } \varphi \longrightarrow u_\varphi \quad \text{where (again) } u_\varphi(f) = \int_{S^1} f(x) \varphi(x) dx \quad (f \in C^\infty(S^1))$$

Typically, the dual of a limit of topological vector spaces is not the colimit of the duals of the limitands. Duals of *colimits* do behave well, in the sense that in reasonable situations

$$\text{Hom}(\text{colim}_i X_i, Z) \approx \text{lim}_i \text{Hom}(X_i, Z)$$

But $C^\infty(S^1)$ is a *limit*, not a colimit. Luckily, the dual of a limit of *Banach spaces* is the colimit of the duals:

[5.1.1] Theorem: Let $X = \text{lim}_i B_i$ be a limit of Banach spaces B_i with projections $p_i : X \rightarrow B_i$. Any $\lambda \in X^* = \text{Hom}_{\mathbb{C}}^o(X, \mathbb{C})$ *factors through* some B_i . That is, there is $\lambda_j : B_j \rightarrow \mathbb{C}$ such that

$$\lambda = \lambda_j \circ p_j : X \rightarrow \mathbb{C}$$

Therefore,

$$(\text{lim}_i B_i)^* \approx \text{colim}_i B_i^*$$

Proof: Without loss of generality, each B_i is the closure of the image of X , since otherwise replacement of each B_i by that closure.

Let U be an open neighborhood of 0 in $X = \text{lim}_i B_i$ such that $\lambda(U)$ is inside the open unit ball at 0 in \mathbb{C} , by the continuity at 0. By properties of the limit topology^[30] there are finitely-many indices i_1, \dots, i_n and open neighborhoods V_{i_t} of 0 in B_{i_t} such that

$$\bigcap_{t=1}^n p_{i_t}^{-1} V_{i_t} \subset U \quad (\text{projections } p_i \text{ from the limit } X)$$

To have λ factor (continuously) through a limitand B_j , we need a *single* condition to replace the conditions from i_1, \dots, i_n . Let j be *any* index^[31] with $j \geq i_t$ for all t , and

$$V'_j = \bigcap_{t=1}^n p_{i_t, j}^{-1} V_{i_t} \subset B_j$$

[30] Recall that $X = \text{lim}_i B_i$ is the closed subspace (with the subspace topology) of the product $Y = \prod_i B_i$ of all tuples $\{b_i\}$ in which $p_{ij} : b_i \rightarrow b_j$ for $i > j$ under the transition maps $p_{ij} : B_i \rightarrow B_j$. A local basis at 0 in the product consists of products $V = \prod_i V_i$ of opens V_i in B_i with $V_i = B_i$ for all but finitely-many i , say i_1, \dots, i_n .

[31] The index set need not be the positive integers, but must be a *poset* (partially ordered set), *directed*, in the sense that for any two indices i, j there is an index k such that $k > i$ and $k > j$.

By the compatibility

$$p_{i_t}^{-1} = p_j^{-1} \circ p_{i_t, j}^{-1}$$

we have a single sufficient condition, namely $p_j^{-1}V'_j \subset U$. By the linearity of λ , for $\varepsilon > 0$

$$\lambda(\varepsilon \cdot p_j^{-1}V_j) = \varepsilon \cdot \lambda(p_j^{-1}V_j) \subset \varepsilon\text{-ball in } \mathbb{C}$$

By continuity^[32] of scalar multiplication on B_j , $\varepsilon \cdot V'_j$ is an open containing 0 in B_j .

We claim that λ factors through $p_j X$ with the subspace topology from B_j . This makes $p_j X$ a *normed* space, if not Banach.^[33] Simplifying notation, let $\lambda : X \rightarrow \mathbb{C}$ and $p : X \rightarrow N$ be continuous linear to a normed space N , with

$$\lambda(p^{-1}V) \subset \text{unit ball in } \mathbb{C} \quad (\text{for some neighborhood } V \text{ of } 0 \text{ in } N)$$

We claim that λ factors through $p : X \rightarrow N$ as a (continuous) linear map. Indeed, by the linearity of λ ,

$$\lambda\left(\frac{1}{n} \cdot p^{-1}V\right) \subset \frac{1}{n}\text{-ball in } \mathbb{C}$$

so

$$\lambda\left(\bigcap_n \frac{1}{n} \cdot p^{-1}V\right) \subset \frac{1}{m}\text{-ball} \quad (\text{for all } m)$$

Then

$$\lambda\left(\bigcap_n \frac{1}{n} \cdot p^{-1}V\right) \subset \bigcap_m \frac{1}{m}\text{-ball} = \{0\}$$

Thus,

$$\bigcap_n p^{-1}\left(\frac{1}{n} \cdot V\right) = \bigcap_n \frac{1}{n} \cdot p^{-1}V \subset \ker \lambda$$

For x, x' in X with $px = px'$, certainly $px - px' \in \frac{1}{n}V$ for all $n = 1, 2, \dots$. Therefore,

$$x - x' \in \bigcap_n p^{-1}\left(\frac{1}{n}V\right) \subset \ker \lambda$$

and $\lambda x = \lambda x'$. This proves the subordinate claim that λ factors through $p : X \rightarrow N$ via a (not necessarily continuous) linear map $\mu : N \rightarrow \mathbb{C}$. For the continuity of μ , by its linearity

$$\mu(\varepsilon \cdot V) = \varepsilon \cdot \mu V \subset \varepsilon\text{-ball in } \mathbb{C}$$

proving the continuity of $\mu : N \rightarrow \mathbb{C}$.^[34] This proves the claim.

[32] Multiplication by a non-zero scalar is a *homeomorphism*: scalar multiplication by $\varepsilon \neq 0$ is continuous, scalar multiplication by ε^{-1} is continuous, and these are mutual inverses, so these scalar multiplications are *homeomorphisms*.

[33] Recall that a *normed* space is a topological vector with topology given by a *norm* $\|\cdot\|$ as in a Banach space, but *without* the requirement that the space is *complete* with respect to the metric $d(x, y) = \|x - y\|$. This slightly complicated assertion is correct: in most useful situations $p_j X$ is rarely all of B_j , even when B_j is a *completion* of $p_j X$.

[34] Here we need V to be *open*, not merely a *set* containing 0. Continuity at 0 is all that is needed for continuity of *linear* maps, since $|\lambda(x)| < \varepsilon$ for $|x| < \delta$ gives $|\lambda(x - x')| < \varepsilon$ for $|x - x'| < \delta$.

The claim gives continuous linear $\lambda_j : p_j X \rightarrow \mathbb{C}$ through which λ factors.

Then $\lambda_j : p_j X \rightarrow \mathbb{C}$ extends by continuity^[35] to the closure of $p_j X$ in B_j , which is B_j , giving the desired map. ///

[5.1.2] **Remark:** The same proof shows that a continuous linear map from a limit of Banach spaces to a normed space factors through a limitand.

[5.1.3] **Corollary:** The space of distributions on S^1 is the ascending union (colimit)

$$C^\infty(S^1)^* = (\lim_k C^k(S^1))^* = \operatorname{colim}_k C^k(S^1)^* = \bigcup_k C^k(S^1)^*$$

of duals of the Banach spaces $C^k(S^1)$. ///

The *order* of a distribution u is the smallest k such that $u \in C^k(S^1)^*$. Since for the circle the space of all distributions is exactly this colimit, the order of a distribution is well-defined.^[36]

Distributions as generalized *functions* should be *differentiable*, compatibly with the differentiation of *functions*. The idea is that differentiation of distributions should be compatible with *integration by parts* for distributions given by integration against C^1 functions. Assuming an integral on S^1 as in the next section, for *functions* f, g , by integration by parts,

$$\int_{S^1} f(x) g'(x) dx = - \int_{S^1} f'(x) g(x) dx$$

with no boundary terms because S^1 has empty boundary. Note the negative sign. Motivated by this, define the *distributional derivative* u' of $u \in C^\infty(S^1)^*$ to be another distribution defined by

$$u'(f) = -u(f') \quad (\text{for any } f \in C^\infty(S^1))$$

The continuity of differentiation $\frac{d}{dx} : C^\infty(S^1) \rightarrow C^\infty(S^1)$ assures that u' is a distribution, since

$$u' = -(u \circ \frac{d}{dx}) : C^\infty(S^1) \rightarrow \mathbb{C}$$

[35] The extension by continuity is unambiguous, since λ_j is *linear*. In more detail: for λ a continuous linear function on a dense subspace Y of a topological vector space X , given $\varepsilon > 0$, take *convex* neighborhood U of 0 in X such that $|\lambda y| < \varepsilon$ for $y \in U$. We may suppose $U = -U$ by replacing U by $-U \cap U$. Let y_i be a Cauchy net approaching $x \in X$. For y_i and y_j inside $x + \frac{1}{2}U$, $|\lambda y_i - \lambda y_j| = |\lambda(y_i - y_j)|$, using the linearity. By the symmetry $U = -U$, since $y_i - y_j \in \frac{1}{2} \cdot 2U = U$, this gives $|\lambda y_i - \lambda y_j| < \varepsilon$. Then unambiguously define λx to be the limit of the λy_i .

[36] The *Riesz representation theorem* asserts that the dual of $C^0(S^1)$ is *Borel measures* on S^1 , so order-zero distributions are *Borel measures*. For example, elements η of $L^2(S^1)$ are Borel measures, by giving integrals $f \rightarrow \int_{S^1} f(x) \eta(x) dx$ for $f \in C^0(S^1)$. Thus, integrating continuous functions against Borel measures is a semi-classical instance of *generalizing functions* in our present style, integrating against *measures*. However, the duals of the higher $C^k(S^1)$'s don't have such a classical interpretation. The fact that $C^1(S^1)$ can be construed as *distributional derivatives of Borel measures* is not strongly related to Radon-Nikodym derivatives of measures, because, for example, the distributional derivative of a point-mass measure is not a measure.

6. Invariant integration, periodicization

We an (*invariant*) *integral* on the circle $S^1 = \mathbb{R}/2\pi\mathbb{Z}$. The main property required is *translation invariance*, meaning that, for a (for example) continuous function f on S^1 ,

$$\int_{S^1} f(x+y) dx = \int_{S^1} f(x) dx \quad (\text{for all } y \in S^1)$$

This invariance is sufficient to prove that various important integrals *vanish*.

For example, let $\psi_m(x) = e^{imx}$. As an instance of an important idea, without explicit calculus-like computations,

[6.1.1] **Claim:** (*Cancellation Lemma*) For $m \neq n$, for *any* reasonable translation-invariant integral on S^1

$$\int_{S^1} \psi_m(x) \bar{\psi}_n(x) dx = 0$$

Proof: For $m \neq n$, the function $f(x) = \psi_m(x) \bar{\psi}_n(x)$ is a *non-trivial* (not identically 1) continuous group homomorphism $S^1 \rightarrow \mathbb{C}^\times$, meaning that there is $y \in S^1$ such that $f(y) \neq 1$. The *change of variables* $x \rightarrow x + y$ in the integral does not change the overall value of the integral, so

$$\int_{S^1} f(x) dx = \int_{S^1} f(x+y) dx = \int_{S^1} f(x) \cdot f(y) dx = f(y) \int_{S^1} f(x) dx$$

Thus, the integral I has the property that $I = t \cdot I$ where $t \neq 1$. This gives $(1-t) \cdot I = 0$, so $I = 0$ since $t \neq 1$. ///

[6.1.2] **Remark:** This *vanishing trick* is impressive, since nothing specific about the continuous group homomorphism f or topological group (S^1 here) is used, apart from the finiteness of the total measure of the group, which comes from its *compactness*. That is, the same proof would show that *integrals over compact groups of non-trivial group homomorphisms are 0*. However, a notion of *invariant measure*^[37] for general groups requires effort. Nevertheless, with an invariant measure, the same argument succeeds.

Less critically than the invariance, we want a *normalization*^[38]

$$\int_{S^1} 1 dx = \text{vol}(S^1) = \text{vol}(\mathbb{R}/2\pi\mathbb{Z}) = 2\pi$$

Then

$$\int_{S^1} |\psi_n(x)|^2 dx = \int_{S^1} 1 dx = 2\pi$$

Thus, without any explicit presentation of the integral or measure, we have proven that the distinct exponentials are an *orthogonal set* with norms $\sqrt{2\pi}$ with respect to the inner product

$$\langle f, g \rangle = \int_{S^1} f(x) \bar{g}(x) dx$$

[37] Translation-invariant measures on topological groups are called *Haar measures*. General proof of their *existence* takes a little work, and invokes the Riesz representation theorem. *Uniqueness* can be made to be an example of a more general argument about uniqueness of invariant functionals.

[38] The measure of the circle need not be normalized to be 2π , but this is natural when presenting it as $\mathbb{R}/2\pi\mathbb{Z}$.

An *integration by parts* formula should be expected, with no *boundary terms* since $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ has empty boundary. Indeed, without *constructing* the invariant integral, we prove what we want from its *properties*:

[6.1.3] **Claim:** Let $f \rightarrow \int_{S^1} f(x) dx$ be an invariant integral on S^1 , for $f \in C^0(S^1)$. Then for $f \in C^1(S^1)$

$$\int_{S^1} f'(x) dx = 0$$

and we have the *integration by parts* formula for $f, g \in C^1(S^1)$

$$\int_{S^1} f(x) g'(x) dx = - \int_{S^1} f(x)' g(x) dx$$

[6.1.4] **Remark:** Vanishing of integrals of derivatives does *not* depend on the particulars of the situation. The same argument succeeds on an arbitrary group possessing (translation) invariant differentiation(s) and an invariant integral. Thus, the specific geometry of the circle is *not* needed to argue that $\int_{S^1} f'(x) dx = \int_0^{2\pi} f(x) dx = f(2\pi) - f(0) = 0$ because f is periodic. The latter classical argument is valid, but fails to show a generally applicable mechanism. The same independence of particulars applies to the integration by parts rule.

Proof: The translation invariance of the integral makes the integral of a derivative 0, by direct computation, as follows. We interchange a *differentiation* and an *integral*.^[39]

$$\int_{S^1} f'(x) dx = \int_{S^1} \frac{\partial}{\partial t} \Big|_{t=0} f(x+t) dx = \frac{d}{dt} \Big|_{t=0} \int_{S^1} f(x+t) dx = \frac{d}{dt} \Big|_{t=0} \int_{S^1} f(x) dx = 0$$

by changing variables in the integral. Then apply this to the function $(f \cdot g)' = f'g + fg'$ to obtain

$$\int_{S^1} f'(x) g(x) dx + \int_{S^1} f(x) g'(x) dx = 0$$

which gives the integration by parts formula. ///

The usual (Lebesgue) integral on the uniformizing \mathbb{R} has the corresponding property of translation invariance. Since we present the circle as a quotient $\mathbb{R} \rightarrow \mathbb{R}/2\pi\mathbb{Z} = S^1$ of \mathbb{R} we expect a *compatibility*^[40]

$$\int_{\mathbb{R}} F(x) dx = \int_{\mathbb{R}/2\pi\mathbb{Z}} \left(\sum_{n \in \mathbb{Z}} F(x + 2\pi n) \right) dx$$

for at least *compactly-supported* continuous functions F on \mathbb{R} .

[39] The argument bluntly demands this interchange of limit and differentiation, so *justification* of it is secondary to the act itself. In the near future this and many other necessary interchanges are definitively justified via *Gelfand-Pettis* (also called *weak*) integrals. In the present concrete situation elementary (but opaque) arguments could be invoked, but we do not do this.

[40] In contrast to many sources, this compatibility is *not* about choosing representatives in $[0, 2\pi)$ or anywhere else for \mathcal{S}^1 . Rather, this compatibility would be required for a group G (here \mathbb{R}), a discrete subgroup Γ (here $2\pi\mathbb{Z}$), and the quotient G/Γ (here S^1), whether or not that quotient is otherwise identifiable. This compatibility is a sort of *Fubini theorem*. The usual Fubini theorem applies to products $X \times Y$, whose quotients $(X \times Y)/X \approx Y$ are simply the factors, but another version applies to quotients that are not necessarily factors.

Indeed, we can *define* integrals of functions on S^1 by this compatibility relation, by expressing a continuous function f on S^1 as a *periodicization* (or *automorphization*)

$$f(x) = \sum_{n \in \mathbb{Z}} F(x + 2\pi n)$$

of a compactly supported continuous function F on \mathbb{R} , and *define*

$$\int_{S^1} f(x) dx = \int_{\mathbb{R}} F(x) dx$$

We still need to prove that this value is independent of the choice of F for given f .

The properties required of an integral on S^1 are clear. Sadly, we are not in a good position (yet) either to prove *uniqueness* or to give a *construction* as gracefully as these ideas deserve.

Postponing a systematic approach, we neglect any proof of uniqueness, and for a construction revert to an ugly-but-tangible reduction of the problem to integration on an interval. That is, note that in the quotient $q : \mathbb{R} \rightarrow \mathbb{R}/2\pi\mathbb{Z} = S^1$ the interval $[0, 2\pi]$ maps surjectively, with the endpoints being identified (and no other points identified). In traditional terminology, $[0, 2\pi]$ is a *fundamental domain*^[41] for the action of \mathbb{Z} on \mathbb{R} . Then define the integral of f on S^1 by

$$\int_{S^1} f(x) dx = \int_0^{2\pi} (f \circ q)(x) dx$$

with usual (Lebesgue) measure on the unit interval. Verification of the *compatibility* with integration on \mathbb{R} is silly, from this viewpoint.

This (bad) definition does allow explicit computations, but makes *translation invariance* harder to prove, since the unit interval gets pushed off itself by translation. But we can still manage the verification.^[42] Take $y \in \mathbb{R}$, and compute

$$\begin{aligned} \int_{S^1} f(x+y) dx &= \int_0^{2\pi} (f \circ q)(x+y) dx = \int_{-y}^{2\pi-y} (f \circ q)(x) dx \\ &= \int_{-y}^0 (f \circ q)(x) dx + \int_0^{2\pi-y} (f \circ q)(x) dx = \int_{-y}^0 (f \circ q)(x-2\pi) dx + \int_0^{2\pi-y} (f \circ q)(x) dx \end{aligned}$$

since $(f \circ q)(x) = (f \circ q)(x-2\pi)$ by periodicity. Then, replacing x by $x+2\pi$ in the first integral, this is

$$\int_{2\pi-y}^{2\pi} (f \circ q)(x) dx + \int_0^{2\pi-y} (f \circ q)(x) dx = \int_0^{2\pi} (f \circ q)(x) dx$$

[41] The notion of *fundamental domain* for the action of a group Γ on a set X has an obvious appeal, at least that it is more concrete than the notion of *quotient* $\Gamma \backslash X$. However, it is rarely possible to determine an exact fundamental domain, *and* one eventually discovers that the details are seldom useful even if this is possible. Instead, the *quotient* should be treated directly.

[42] While suppressing our disgust.

7. Hilbert space theory of Fourier series

The spaces of functions $C^k(S^1)$ are Banach spaces, not Hilbert spaces. ^[43] The geometry of Hilbert spaces motivates finding a way to compare these Banach spaces with Hilbert spaces. Fourier series provide a good device for this comparison.

As discussed above, we do not need an *explicit* or *formulaic* integral on S^1 , only the *property of translation invariance*, choice of *normalization*, and the property of *compatibility* with integration on \mathbb{R} . Nevertheless, for expediency, we may integrate functions on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ by pulling back to \mathbb{R} and integrating on $[0, 1]$.

As usual, $L^2(S^1) \approx L^2[0, 2\pi]$ can be described as the collection of measurable functions f on $[0, 2\pi]$ such that

$$\int_0^{2\pi} |f(x)|^2 dx < +\infty$$

modulo the equivalence relation of almost-everywhere equality, with the inner product

$$\langle f, g \rangle = \int_0^{2\pi} f(x) \bar{g}(x) dx$$

This definition makes $L^2[0, 2\pi]$ a *Hilbert space*. ^[44]

As observed in the previous section, the exponential functions

$$\psi_n(x) = e^{inx} \quad (\text{for } n \in \mathbb{Z})$$

form an orthogonal *set* in $L^2(S^1)$. It is not clear that they form an orthogonal *basis*. That is, we should show that the finite linear combinations of the exponential functions ψ_n are *dense* in $L^2(S^1)$. The *Fourier coefficients* of an L^2 function f are essentially inner products:

$$n^{\text{th}} \text{ Fourier coefficient of } f = \hat{f}(n) = \frac{1}{2\pi} \langle f, \psi_n \rangle = \frac{1}{2\pi} \int_0^{2\pi} f(x) \bar{\psi}(x) dx$$

The *Fourier expansion* of f in $L^2(S^1)$ is

$$f \sim \sum_{n \in \mathbb{Z}} \hat{f}(n) \psi_n = \sum_{n \in \mathbb{Z}} \frac{\langle f, \psi_n \rangle}{2\pi} \psi_n$$

[7.0.1] Remark: We do not write *equality* here, for several reasons, clarified in the sequel. We *do* expect an L^2 equality, although this is not a priori related to *pointwise* equality, which is one of the major issue addressed below.

^[43] It is not trivial to prove that there are no Hilbert space structures on these spaces of functions, without knowing the ramifications of the geometry of Hilbert spaces. Even when the given Banach space structure is not Hilbert, this does not preclude existence of a *different* topology coming from a Hilbert structure. More to the point is that the *natural* structures are Banach.

^[44] The completeness requires proof. The kernel of the idea is that a Cauchy sequence of L^2 functions has a subsequence that converges pointwise off a set of measure 0. This gives a pointwise definition of the candidate limit. A different definition of $L^2(S^1)$, namely as the completion of $C^o(S^1)$ with respect to the L^2 metric has the virtue that the completeness is defined into it, but the workload has been shifted to proving that these two definitions are the same.

[7.1] Completeness in $L^2(S^1)$

There are many different proofs that the normalized exponentials $\psi_n/\sqrt{2\pi}$ form an orthonormal basis in $L^2(S^1)$. At one extreme, a completely analogous result holds for any *compact abelian* topological group, without further structure. Even the abelian-ness can be dropped without much harm, though with complications. Further, formulation and proof of that general result use methods of broad utility, so we will give it later. For the moment, we take advantage of the particulars of the circle to give a 19th-century-style argument. We *do* take this opportunity to introduce the notion of *approximate identity* (below).

[7.1.1] **Theorem:** The normalized exponentials $\psi_n/\sqrt{2\pi}$ are an orthonormal basis for $L^2(S^1)$.

[7.1.2] **Remark:** The proof does *not* prove that Fourier series converge pointwise (which they often do not).

[7.1.3] **Remark:** The not-necessarily-uniform pointwise convergence of Fourier series of C^1 functions does *not* instantly yield L^2 convergence, nor does it instantly yield *uniform* pointwise convergence, which *would* imply L^2 convergence. Modifying the earlier pointwise convergence idea to achieve these goals motivates introduction of an *approximate identity* in the proof.

Proof: By *Urysohn's Lemma* (see appendix), the continuous functions $C^o(S^1)$ are dense in $L^2(S^1)$. Thus, it suffices to prove that C^o functions are approximable in L^2 by finite sums of the exponentials ψ_n , and it suffices to prove that finite sums of exponentials approximate C^o functions in the C^o topology: the total measure of the space S^1 is finite, so the L^2 norm of a continuous function is dominated by its sup norm, and density in sup norm implies density in L^2 norm. [45]

We review the situation. As in the proof of not-necessarily-uniform pointwise convergence of Fourier series for piecewise C^1 functions, the N^{th} partial sum of the Fourier series of a function f on $S^1 = \mathbb{R}/2\pi\mathbb{Z}$ can be described usefully as an *integral operator*[46] by

$$\begin{aligned} \frac{1}{2\pi} \sum_{|k|\leq n} \langle f, \psi_k \rangle \psi_k(x) &= \frac{1}{2\pi} \int_{S^1} f(y) \sum_{|k|\leq n} \overline{\psi_k}(y) \psi_k(x) dy = \frac{1}{2\pi} \int_{S^1} f(y) \sum_{|k|\leq n} \psi_k(x-y) dy \\ &= \frac{1}{2\pi} \int_{S^1} f(y) \frac{\psi_{n+1}(x-y) - \psi_{-n}(x-y)}{\psi_1(x-y) - 1} dy \end{aligned}$$

by summing finite geometric series. Let $K_n(x)$ be the summed geometric series

$$K_n(x) = \frac{\psi_{n+1}(x) - \psi_{-n}(x)}{\psi_1(x) - 1} = \frac{e^{(n+1)ix} - e^{-inx}}{e^{-ix} - 1} = \frac{e^{(n+\frac{1}{2})x} - e^{-(n+\frac{1}{2})x}}{e^{ix/2} - e^{-ix/2}} = \frac{\sin(n + \frac{1}{2})x}{\sin \frac{x}{2}}$$

so, dropping the $1/2\pi$,

$$\sum_{|k|\leq n} \langle f, \psi_k \rangle \psi_k(x) = \int_{S^1} f(y) K_n(x-y) dy$$

[45] Fortunately, proving that C^o functions are approximable in the sup norm by finite sums of exponentials does *not* require proving that Fourier series of continuous functions converge pointwise, which is not generally true. That is, we are *not* compelled to prove that the partial sums of the Fourier series are the approximating sums, which admittedly *are* the obvious candidate sequence. It turns out that this cannot possibly succeed. Examples of continuous functions whose Fourier series diverge were suggested by Riemann, made rigorous by Weierstraß, and treated carefully in Fejér, L, (1910) *Beispiele stetiger Funktionen mit divergenter Fourierreihe* Journal Reine Angew. Math. 137, pp. 1-5. Existential arguments use the *Baire category theorem*.

[46] The notion of *integral operator* T is general, and without genuine precise definition. The *form* is that $Tf(x) = \int K(x, y) f(y) dy$, and the function $K(x, y)$ is the *kernel* of the operator. The class of functions in which the kernel lies, and in which the input and output lie, varies enormously with context. Schwartz' kernel theorem hugely generalizes this idea.

Granting that it is futile to prove that the partial sums converge pointwise for continuous functions, we might try to see what related but different integral operators would work better.

This is our excuse to introduce *approximate identities* in this situation. [47] The *rough* idea of *approximate identity* is of a sequence $\{\varphi_n\}$ of functions φ_n approximating a *point-mass measure* [48] at $0 \in S^1$. Precisely, a sequence of continuous functions φ_n on S^1 is an approximate identity when

$$\varphi_n(x) \geq 0 \quad (\text{for all } n, x) \quad \int_{S^1} \varphi_n(x) dx = 1 \quad (\text{for all } n)$$

and if for every $\varepsilon > 0$ and for every $\delta > 0$ there is n_o such that for all $n \geq n_o$

$$\int_{|x| < \delta} \varphi_n(x) dx > 1 - \varepsilon \quad (\text{equivalently, } \int_{\delta \leq |x| < \frac{1}{2}} \varphi_n(x) dx < \varepsilon)$$

where we use coordinates in \mathbb{R} for $S^1 = \mathbb{R}/2\pi\mathbb{Z}$. That is, the functions φ_n are non-negative, their integrals are all 1, and their *mass bunches up* at $0 \in S^1$. It is not surprising that the integral operators made from an approximate identity have a useful property:

[7.1.4] **Claim:** For $f \in C^o(S^1)$ on S^1 and for an approximate identity φ_n , in the topology of $C^o(S^1)$,

$$\lim_n \int_{S^1} f(y) \varphi_n(x - y) dy = f(x)$$

Granting this claim, making an approximate identity out of finite sums of exponentials will prove that such finite sums are dense in $C^o(S^1)$. [49]

Proof: (of claim) Given $f \in C^o(S^1)$, and given $\varepsilon > 0$, by *uniform* continuity of f on the compact S^1 , there is $\delta > 0$ such that

$$|f(x) - f(y)| < \varepsilon \quad (\text{for } |x - y| < \delta)$$

Take n large enough so that

$$\int_{|x| < \delta} \varphi_n(x) dx > 1 - \varepsilon$$

Since the total mass of φ_n is 1,

$$\begin{aligned} \int_{S^1} f(y) \varphi_n(x - y) dy - f(x) &= \int_{S^1} (f(y) - f(x)) \varphi_n(x - y) dy \\ &= \int_{|y-x| < \delta} (f(y) - f(x)) \varphi_n(x - y) dy + \int_{\delta \leq |y-x| \leq \frac{1}{2}} (f(y) - f(x)) \varphi_n(x - y) dy \end{aligned}$$

The second integral is easily estimated by

$$\left| \int_{\delta \leq |y-x| \leq \frac{1}{2}} (f(y) - f(x)) \varphi_n(x - y) dy \right| \leq 2|f|_{C^o} \int_{\delta \leq |y-x| \leq \frac{1}{2}} \varphi_n(x - y) dy \leq 2|f|_{C^o} \cdot \varepsilon$$

[47] Versions of *approximate identity* are useful and important on *topological groups* generally.

[48] A *point-mass* measure at a point x_o is a measure which gives the point x_o measure 1 and gives a set not containing x_o measure 0. These are also called *Dirac measures*. It is easy to give a too-naive formulation of the notion of approximate identity, which *fails*. In particular, the requirement that φ_n form an approximate identity is *strictly stronger* than the condition that φ_n approach a Dirac measure in a *distributional* sense. Specifically, the *non-negativity* condition on an approximate identity is indispensable.

[49] Similarly, to give S. Bernstein's tangible proof of the Stone-Weierstraß theorem that polynomials are dense in $C^o(K)$ for compact K in \mathbb{R}^n , an *approximate identity* is exhibited consisting of *polynomials*.

Estimation of the integral near 0 uses the *positivity* of the φ_n :

$$\left| \int_{|y-x|<\delta} (f(y) - f(x)) \varphi_n(x-y) dy \right| \leq \varepsilon \int_{|y-x|<\delta} \varphi_n(x-y) dy \leq \varepsilon \int_{S^1} \varphi_n(x-y) dy = \varepsilon \cdot 1$$

This holds for all $\varepsilon > 0$ and uniformly in x , so the integrals approach $f(x)$ in the C^0 topology, proving the claim. ///

As noted above, to prove the completeness, we need only exhibit an approximate identity made from finite sums of exponentials. A failing of the *Dirichlet kernels*

$$K_n(x) = \frac{e^{(n+\frac{1}{2})x} - e^{-(n+\frac{1}{2})x}}{e^{ix/2} - e^{-ix/2}} = \frac{\sin(n+\frac{1}{2})x}{\sin \frac{x}{2}}$$

that yield the partial sums is that these kernels are not non-negative. Still, the fact (proven above) that a stronger hypothesis of C^1 -ness gives a limited result in the direction we want *suggests* that the masses of the K_n *do* bunch up near 0. Indeed, the expression for K_n in terms of sines *does* show that these functions are *real-valued*. Thus, a plausible choice for an approximate identity φ_n is the (essentially) *Fejér kernels*

$$\varphi_n = K_n^2 \times (\text{constant depending on } n)$$

with constant chosen to give total mass 1. Computing directly in terms of exponentials, expanding the square of the sum,

$$\begin{aligned} \int_{S^1} K_n(x)^2 dx &= \int_{S^1} (e^{-inx} + \dots + e^{inx})^2 dx \\ &= \int_{S^1} e^{-2nix} + 2e^{-(2n-1)ix} + 3e^{-(2n-2)ix} + 4e^{-(2n-3)ix} + \dots + (2n+1) \cdot 1 + \dots + 2e^{(2n-1)ix} + e^{2nix} dx \\ &= \int_{S^1} 2n+1 dx = 2\pi \cdot (2n+1) \end{aligned}$$

since all the non-trivial exponentials integrate to 0. Thus, take

$$\varphi_n(x) = \frac{K_n(x)^2}{2\pi \cdot (2n+1)}$$

The discussion so far gives non-negativity and total mass 1. We must show that the mass of the φ_n 's bunches up at 0. For this, revert to the expression for K_n in terms of *sines*. For $\delta \leq |x| \leq \pi$,

$$\varphi_n(x) = \frac{\sin^2(n+\frac{1}{2})x}{2\pi(2n+1)\sin^2 \frac{x}{2}} \leq \frac{1}{2\pi(2n+1)x^2} \quad (\text{since } |\sin \frac{x}{2}| \geq |\frac{x}{2}| \text{ for } |x| \leq \pi)$$

For x bounded away from 0 we get inequalities such as

$$0 \leq \varphi_n(x) \leq \frac{1}{2\pi(2n+1)x^2} \leq \frac{1}{2\pi(2n+1)n^{-2/3}} \leq \frac{1}{2\pi(2n^{1/3}+n^{-2/3})} \leq \frac{1}{2\pi n^{1/3}} \quad (\text{for } |x| \geq n^{-1/3})$$

This sup-norm estimate for φ_n on the part of S^1 covered by $n^{-1/3} \leq |x| \leq \pi$ gives

$$\int_{n^{-1/3} \leq |x| \leq \pi} \varphi_n(x) dx \leq 2\pi \cdot \frac{1}{2\pi n^{1/3}} \leq \frac{1}{n^{1/3}}$$

Thus, given $\varepsilon > 0$ and $\delta > 0$, take n large enough so that $n > (2\varepsilon)^3$, and $n > \delta^3$, to meet the mass-concentration criterion for approximate identities. ///

Therefore, having shown that the exponential functions form an orthonormal basis, for $f \in L^2[0, 1]$ we have an L^2 -equality

$$f = \sum_n \frac{\langle f, \psi_n \rangle}{2\pi} \psi_n = \sum_n \widehat{f}(n) \psi_n \quad (\text{in } L^2(S^1))$$

and the *Plancherel-Parseval theorem* from general Hilbert space theory gives a Plancherel-Parseval theorem here:

$$|f|^2 = \sum_n \left| \frac{\langle f, \psi_n \rangle}{2\pi} \right|^2 = \sum_n |\widehat{f}(n)|^2$$

[7.1.5] **Remark:** Again, L^2 convergence says *nothing* directly about *pointwise* convergence. Nor is there anything to deny the possibility that a Fourier series *does converge* at a point, but converges to a value *different* from the value of f there. For example, we proved pointwise convergence of C^1 functions, but what about C^0 ?

[7.1.6] **Remark:** Fourier already worried about pointwise convergence of Fourier series, as did Cantor. From a later viewpoint than theirs, since L^2 functions are only defined *almost everywhere*, pointwise convergence of a Fourier series would distinguish a special function in the equivalence class in $L^2[0, 1]$, which might be suspicious. Nevertheless, L. Carleson showed [50] that, given $\sum_n |c_n|^2 < \infty$, the Fourier series $\sum_n c_n e^{inx}$ converges almost everywhere. Thus, given $f \in L^2[0, 2\pi]$, the Fourier series of f *does* converge almost everywhere to f , and *does* distinguish an element in that almost-everywhere-equal equivalence class. Fortunately, this difficult result does not play a role here.

8. *Levi-Sobolev inequalities, Levi-Sobolev imbeddings*

The simplest L^2 theory of Fourier series addresses neither continuity nor differentiability. Yet it *would be* advantageous on general principles to be able to talk about differentiability in the context of Hilbert spaces, since Hilbert spaces have easily understood *dual spaces*. Beppo Levi, Frobenius, and Sobolev made useful comparisons. The idea is to compare C^k norms to norms coming from Hilbert spaces whose inner products refer to derivatives, the Levi-Sobolev spaces.

[8.1] Levi-Sobolev inequalities

First, we have an easy estimate for a variant C^k norm:

$$\left| \sum_{|n| \leq N} c_n e^{inx} \right|_{C^k} = \sup_{0 \leq j \leq k} \sup_x \left| \sum_{|n| \leq N} c_n (in)^j e^{inx} \right|_{\mathbb{C}} \leq \sum_{|n| \leq N} |c_n| \cdot (1 + n^2)^{k/2}$$

all for elementary reasons. [51] Perhaps surprisingly, rather try to directly obtain a sup norm estimate on this sum, Cauchy-Schwarz-Bunyakovsky is invoked: for any $s \in \mathbb{R}$

$$\begin{aligned} \left| \sum_{|n| \leq N} c_n e^{inx} \right|_{C^k} &\leq \sum_{|n| \leq N} |c_n| \cdot (1 + n^2)^{s/2} \cdot \frac{1}{(1 + n^2)^{(s-k)/2}} \\ &\leq \left(\sum_{|n| \leq N} |c_n|^2 \cdot (1 + n^2)^s \right)^{1/2} \cdot \left(\sum_{|n| \leq N} \frac{1}{(1 + n^2)^{s-k}} \right)^{1/2} \end{aligned}$$

[50] L. Carleson, *On convergence and growth of partial sums of Fourier series*, Acta Math. **116** (1966), 135-157.

[51] The awkward expression $(1 + n^2)^{1/2}$ is approximately n . However, for $n = 0$ we cannot divide by n , and replacing n by $(1 + n^2)^{1/2}$ is the traditional device stunt to avoid this annoyance.

Convergence of the elementary sum is easy to understand:

$$\sum_{n \in \mathbb{Z}} \frac{1}{(1+n^2)^{s-k}} < +\infty \quad (\text{for } s > k + \frac{1}{2})$$

Thus, for any $s > k + \frac{1}{2}$ we have a *Levi-Sobolev inequality*

$$\begin{aligned} \left| \sum_{|n| \leq N} c_n \psi_n \right|_{C^k} &\leq \left(\sum_{n \in \mathbb{Z}} \frac{1}{(1+n^2)^{s-k}} \right)^{1/2} \cdot \left(\sum_{|n| \leq N} |c_n|^2 \cdot (1+n^2)^s \right)^{1/2} \\ &\leq \left(\sum_{n \in \mathbb{Z}} \frac{1}{(1+n^2)^{s-k}} \right)^{1/2} \cdot \left(\sum_{n \in \mathbb{Z}} |c_n|^2 \cdot (1+n^2)^s \right)^{1/2} \end{aligned}$$

which is summarized as

$$\left| \sum_{n \in \mathbb{Z}} c_n \psi_n \right|_{C^k} \leq \left(\sum_{n \in \mathbb{Z}} \frac{1}{(1+n^2)^{s-k}} \right)^{1/2} \cdot \left(\sum_{n \in \mathbb{Z}} |c_n|^2 \cdot (1+n^2)^s \right)^{1/2} \quad (\text{for } s > k + \frac{1}{2})$$

Existence of this comparison makes the right side interesting. Taking away from the right-hand side the uniform constant

$$\omega_{s-k} = \left(\sum_{n \in \mathbb{Z}} \frac{1}{(1+n^2)^{s-k}} \right)^{1/2}$$

gives the s^{th} *Levi-Sobolev norm*

$$s^{\text{th}} \text{ Levi-Sobolev norm} = \left| \sum_{n \in \mathbb{Z}} c_n \psi_n \right|_{H^s} = \left(\sum_{n \in \mathbb{Z}} |c_n|^2 \cdot (1+n^2)^s \right)^{1/2}$$

Paraphrasing, we have proven the dominance relation

$$\left| \cdot \right|_{C^k} \leq \omega_{s-k} \cdot \left| \cdot \right|_{H^s} \quad (\text{for any } s > k + \frac{1}{2})$$

[8.2] Levi-Sobolev imbeddings

For $s \geq 0$, the s^{th} *Levi-Sobolev space* is [52]

$$H^s(S^1) = \{f \in L^2(S^1) : \sum_n |\widehat{f}(n)|^2 \cdot (1+n^2)^s < +\infty\}$$

The inner product on $H^s(S^1)$ is

$$\left\langle \sum_n a_n \psi_n, \sum_n b_n \psi_n \right\rangle = 2\pi \sum_n a_n \bar{b}_n (1+n^2)^s$$

[52] This definition is fine for $s \geq 0$, but not sufficient for $s < 0$. We will give the broader definition below. Keep in mind that $L^2(S^1)$ contains $C^0(S^1)$ and all the $C^k(S^1)$'s.

[8.2.1] **Remark:** This definition of $H^s(S^1)$ defines a useful space of functions or generalized functions only for $s \geq 0$, since for $s < 0$ the constraint $f \in L^2(S^1)$ is *stronger* (from the Plancherel theorem) than the condition defining $H^s(S^1)$ in the previous display.

[8.2.2] **Remark:** The 0^{th} Levi-Sobolev space is just $L^2(S^1)$.

[8.2.3] **Corollary:** For $s > k + \frac{1}{2}$ there is a continuous inclusion

$$H^s(S^1) \subset C^k(S^1)$$

Proof: For $s > k + \frac{1}{2}$, whenever a Fourier series has a finite Levi-Sobolev norm

$$\left| \sum_n c_n \psi_n \right|_{H^s} = \left(\sum_{n \in \mathbb{Z}} |c_n|^2 \cdot (1 + n^2)^s \right)^{1/2} < +\infty$$

the partial sums of the Fourier series are Cauchy in H^s , hence Cauchy in C^k , so converge in the Banach space C^k :

$$\sum_n c_n \psi_n = C^k \text{ function on } S^1$$

Proof: Apply the Levi-Sobolev inequality $|f|_{C^k} \leq \omega \cdot |f|_{H^s}$ to finite linear combinations f of exponentials. Such finite linear combinations are C^k , and the inequality implies that an infinite sum of such, convergent in $H^s(S^1)$, has sequence of partial sums convergent in $C^k(S^1)$. That is, by the completeness of $C^k(S^1)$, the limit is still k times continuously differentiable. Thus, we have the containment. Given the containment, the inequality of norms implies the continuity of the inclusion. ///

[8.3] Levi-Sobolev Hilbert spaces

[8.3.1] **Claim:** The s^{th} Levi-Sobolev space $H^s(S^1)$ (with $0 \leq s \in \mathbb{R}$) is a Hilbert space. In particular, the sequences of Fourier coefficients of functions in $H^s(S^1)$ are *all* two-sided sequences $\{c_n : n \in \mathbb{Z}\}$ of complex numbers meeting the condition

$$\sum_n |c_n|^2 \cdot (1 + n^2)^s < +\infty$$

[8.3.2] **Remark:** It is clear that the exponentials ψ_n are an *orthogonal* basis for $H^s(S^1)$, although their norms depend on the index s . In particular, the collection of finite linear combinations of exponentials is *dense* in $H^s(S^1)$.

[8.3.3] **Remark:** Again, we do want to *define* these positively-indexed Levi-Sobolev spaces as subspaces of genuine spaces of functions, *not* as sequences of Fourier coefficients meeting the condition, and then *prove* the second assertion of the claim. This does leave open, for the moment, the question of how to define negatively-indexed Levi-Sobolev spaces.

Proof: In effect, this is the space of L^2 functions on which the H^s -norm is finite. If we prove the second assertion of the claim, then invoke the usual proof that L^2 spaces are complete to know that $H^s(S^1)$ is complete, since it is simply a weighted L^2 -space. Given a two-sided sequence $\{c_n\}$ of complex numbers such that

$$\sum_n |c_n|^2 \cdot (1 + n^2)^s < +\infty$$

since $s \geq 0$,

$$\sum_n |c_n|^2 < +\infty$$

and, by Plancherel,

$$\sum_n c_n \psi_n \in L^2(S^1)$$

This shows that $H^s(S^1)$ is a Hilbert space for $s \geq 0$. ///

[8.3.4] Remark: Insisting on viewing $L^2(S^1)$ as *equivalence classes* of functions may mislead us into making a needlessly complicated assertion about Levi-Sobolev imbeddings $H^s(S^1) \subset C^k(S^1)$ for $s > k + \frac{1}{2}$, by insisting that $H^s(S^1)$ consists of almost-everywhere equivalence classes of $L^2(S^1)$ functions, only *one* of which is in $C^k(S^1)$. This is not a genuine issue.

[8.4] Levi-Sobolev norms in terms of derivatives

[8.4.1] Remark: Apart from having the virtue of giving inner-product structures, the expressions appearing in these Levi-Sobolev norms are *natural* because they have meaning in terms of L^2 -norms of derivatives. For $f = \sum c_n \psi_n \in C^k(S^1)$, by Plancherel

$$\begin{aligned} (\text{norm via derivatives}) &= |f|^2 + |f'|^2 + |f''|^2 + \dots + |f^{(k)}|^2 \\ &= \sum_n |c_n|^2 \cdot (1 + n^2 + n^4 + \dots + n^{2k}) \leq \sum_n |c_n|^2 \cdot (1 + n^2)^k \end{aligned}$$

Conversely,

$$(1 + n^2)^k \leq C_k \cdot (1 + n^2 + n^4 + n^6 + \dots + n^{2k}) \quad (\text{for some constant } C_k)$$

so

$$(\text{norm via Fourier coefficients}) = \sum_n |c_n|^2 \cdot (1 + n^2)^k \leq C_k \cdot (|f|^2 + |f'|^2 + |f''|^2 + \dots + |f^{(k)}|^2)$$

Thus, the two definitions of Levi-Sobolev norms, in terms of weighted L^2 norms of Fourier series, or in terms of L^2 norms of derivatives, give comparable Hilbert space structures. In particular, the *topologies* are identical.

[8.4.2] Corollary: For $k \geq 0$,

$$C^k(S^1) \subset H^k(S^1)$$

Proof: For $k = 0$, the assertion is that $C^0(S^1) \subset L^2(S^1)$, which holds because S^1 is compact. Similarly, the relevant derivatives of $f \in C^k(S^1)$ are in $L^2(S^1)$, so $f \in H^k(S^1)$. ///

[8.4.3] Remark: One can work out the corresponding inequalities for Fourier series in several variables, proving that $(k + \frac{n}{2} + \varepsilon)$ -fold L^2 differentiability (for any $\varepsilon > 0$) in dimension n is needed to assure k -fold continuous differentiability. This is L^2 Levi-Sobolev theory.

[8.5] Uniform pointwise convergence, convergence in $C^k(S^1)$

At this moment it is very easy to give a *straightforward*, if not *sharp*, result about convergence of C^k functions on S^1 , via the Levi-Sobolev spaces:

[8.5.1] Corollary: The Fourier series of $f \in C^k(S^1)$ converges to f in $C^{k-1}(S^1)$.

Proof: A function in $C^k(S^1)$ is in the Hilbert space $H^k(S^1)$, meaning that the finite partial sums of the Fourier expansion converge to f in $H^k(S^1)$. The $H^k(S^1)$ norm dominates that of $C^{k-1}(S^1)$, so the Fourier series converges to f in $C^{k-1}(S^1)$. ///

[8.5.2] Remark: It may seem mildly peculiar that the Fourier series of a C^k function can converge to it only in C^{k-1} .

[8.6] L^2 -differentiation

[8.6.1] Claim: For every $s \geq 0$, the differentiation map

$$\frac{d}{dx} : \text{finite Fourier series} \longrightarrow \text{finite Fourier series}$$

is continuous when the source is given the $H^s(S^1)$ topology and the target is given the $H^{s-1}(S^1)$ topology.

Proof: This continuity is by design:

$$\begin{aligned} \left| \frac{d}{dx} \sum_{|n| \leq N} c_n e^{inx} \right|_{H^{s-1}}^2 &= \left| \sum_{|n| \leq N} c_n in e^{inx} \right|_{H^{s-1}}^2 \leq \sum_{|n| \leq N} |nc_n|^2 \cdot (1+n^2)^{s-1} \\ &\leq \sum_{|n| \leq N} |c_n|^2 \cdot (1+n^2)^s = \left| \sum_{|n| \leq N} c_n e^{inx} \right|_{H^s}^2 \end{aligned}$$

proving the continuity on finite Fourier series. ///

Therefore, we can *extend* $\frac{d}{dx}$ *by continuity* to obtain continuous linear maps

$$(L^2\text{-differentiation}) = (\text{extension by continuity of}) \frac{d}{dx} : H^s(S^1) \longrightarrow H^{s-1}(S^1)$$

[8.6.2] Remark: In these terms, *extra* L^2 -differentiability is needed to assure comparable classical *continuous* differentiability. Specifically, $(k + \frac{1}{2} + \varepsilon)$ -fold L^2 -differentiability (for any $\varepsilon > 0$) suffices for k -fold *continuous* differentiability, in this one-dimensional example. *The comparable computations on $(S^1)^{\times n}$ show that the gap widens as the dimension grows.*

9. $C^\infty = \lim C^k = \lim H^s = H^\infty$

For larger purposes, the specific comparisons of indices in the containments

$$H^s(S^1) \subset C^k(S^1) \quad (\text{for } s > k + \frac{1}{2})$$

$$C^k(S^1) \subset H^s(S^1) \quad (\text{for } k \geq s)$$

are secondary, since we are more interested in *smooth functions* $C^\infty(S^1)$ than functions with *limited* continuous differentiability.

Thus, the point is that the Levi-Sobolev spaces and $C^k(S^1)$ spaces are *cofinal* under taking *descending intersections*. That is, letting $H^\infty(S^1)$ be the *intersection* of all the $H^s(S^1)$, as *sets* we have

$$C^\infty(S^1) = \bigcap_k C^k(S^1) = \bigcap_{s \geq 0} H^s(S^1) = H^\infty(S^1)$$

Since descending nested intersections are *limits*, the topologies behave well for trivial reasons:

[9.0.1] Theorem: As topological vector spaces

$$C^\infty(S^1) = \lim_k C^k(S^1) = \lim_{s \geq 0} H^s(S^1) = H^\infty(S^1)$$

Proof: The cofinality of the C^k 's and the H^s 's gives a natural isomorphism of the two limits, since they can be combined in a larger limit in which each is cofinal. ///

Again, in general duals of limits are not colimits, but we did show earlier that the dual of a limit of *Banach* spaces is the colimit of the duals of the Banach spaces. Thus,

[9.0.2] Corollary: The space of distributions on S^1 is

$$C^\infty(S^1)^* = \operatorname{colim}_k C^k(S^1)^* = \operatorname{colim}_{s \geq 0} H^s(S^1)^* = H^\infty(S^1)^*$$

(and the duals $H^s(S^1)^*$ admit further explication, below). ///

Expressing $C^\infty(S^1)$ as a limit of the Hilbert spaces $H^s(S^1)$, as opposed to its more natural expression as a limit of the Banach spaces $C^k(S^1)$, is convenient when taking *duals*, since by the *Riesz-Fischer theorem*^[53] we have explicit expressions for Hilbert space duals. We exploit this possibility below.

10. Distributions, generalized functions, again

We will see that distributions on S^1 have Fourier expansions, greatly facilitating their study.^[54]

The exponential functions ψ_n are in $C^\infty(S^1)$, so for any distribution u we can compute Fourier coefficients of u by

$$(n^{\text{th}} \text{ Fourier coefficient of } u) = \widehat{u}(n) = \frac{1}{2\pi} \cdot u(\psi_{-n})$$

Write

$$u \sim \sum_n \widehat{u}(n) \cdot \psi_n$$

even though *pointwise* convergence of the indicated sum is certainly not expected. Define Levi-Sobolev spaces for all $s \in \mathbb{R}$ by

$$H^s(S^1) = \{u \in C^\infty(S^1)^* : \sum_n |u(\psi_{-n})|^2 \cdot (1 + n^2)^s < \infty\}$$

and the s^{th} Levi-Sobolev norm $|u|_{H^s}$ is

$$|u|_{H^s}^2 = \sum_n |u(\psi_{-n})|^2 \cdot (1 + n^2)^s$$

For $0 \leq s \in \mathbb{Z}$, this definition is visibly compatible with the previous definition via derivatives.

[10.0.1] Remark: The formation of the Levi-Sobolev spaces of both positive and negative indices portrays the classical *functions* of various degrees of (continuous) differentiability together with *distributions* of various orders as fitting together as comparable objects. By contrast, thinking only in terms of the spaces $C^k(S^1)$ does not immediately suggest a comparison with distributions.

[53] The Riesz-Fischer theorem asserts that the (continuous) dual V^* of a Hilbert space V is \mathbb{C} -conjugate linearly isomorphic to V . The isomorphism from V to V^* attaches the linear functional $v \rightarrow \langle v, w \rangle$ to an element $w \in V$. Since our hermitian inner products \langle, \rangle are *conjugate*-linear in the second argument, the map $w \rightarrow \langle, w \rangle$ is conjugate linear.

[54] In contrast, discussion of distributions on the real line \mathbb{R} is more complicated, due to the non-compactness of \mathbb{R} . *Not* every distribution on \mathbb{R} is the Fourier transform of a *function*. Distributions which *admit* Fourier transforms, *tempered* distributions, constitute a proper subset of all distributions on \mathbb{R} .

For convenience, define a *weighted* version $\ell^{2,s}$ of (a two-sided version of) the classical Hilbert space ℓ^2 by [55]

$$\ell^{2,s} = \left\{ \{c_n : n \in \mathbb{Z}\} : \sum_{n \in \mathbb{Z}} |c_n|^2 \cdot (1+n^2)^s < \infty \right\}$$

with the weighted version of the usual hermitian inner product, namely,

$$\langle \{c_n\}, \{d_n\} \rangle = \sum_{n \in \mathbb{Z}} c_n \overline{d_n} \cdot (1+n^2)^s$$

[10.0.2] **Claim:** The complex bilinear *pairing*

$$\langle \cdot, \cdot \rangle : \ell^{2,s} \times \ell^{2,-s} \longrightarrow \mathbb{C}$$

by

$$\langle \{c_n\}, \{d_n\} \rangle = \sum_n c_n d_{-n}$$

identifies these two Hilbert spaces as mutual duals, where

$$\ell^{2,-s} \longrightarrow (\ell^{2,s})^* \quad \text{by} \quad \{d_n\} \rightarrow \lambda_{\{d_n\}} \quad \text{where} \quad \lambda_{\{d_n\}}(\{c_n\}) = \sum_n c_n d_{-n}$$

[10.0.3] **Remark:** The minus sign in the subscript in the last formula is not the main point, but is a necessary artifact of our change from a *hermitian* form to a *complex bilinear* form. It is (thus) necessary to maintain compatibility with the Plancherel theorem for ordinary functions.

Proof: The Cauchy-Schwarz-Bunyakovsky inequality gives the continuity of the functional attached to $\{d_n\}$ in $\ell^{2,-s}$ by

$$\begin{aligned} \left| \sum_n c_n \cdot d_{-n} \right| &\leq \sum_n |c_n| (1+n^2)^{s/2} \cdot |d_{-n}| (1+n^2)^{-s/2} \\ &\leq \left(\sum_n |c_n|^2 (1+n^2)^s \right)^{1/2} \cdot \left(\sum_n |d_n|^2 (1+n^2)^{-s} \right)^{1/2} = |\{c_n\}|_{\ell^{2,s}} \cdot |\{d_n\}|_{\ell^{2,-s}} \end{aligned}$$

proving the continuity. [56] To prove the surjectivity we adapt the Riesz-Fischer theorem by a renormalization. That is, given a continuous linear functional λ on $\ell^{2,s}$, by Riesz-Fischer there is $\{a_n\} \in \ell^{2,s}$ such that

$$\lambda(\{c_n\}) = \langle \{c_n\}, \{a_n\} \rangle_{\ell^{2,s}} = \sum_n c_n \cdot \overline{a_n} \cdot (1+n^2)^s$$

Take

$$d_n = \overline{a_{-n}} \cdot (1+n^2)^s$$

Check that this sequence of complex numbers is in $\ell^{2,-s}$, by direct computation, using the fact that $\{a_n\} \in \ell^{2,s}$,

$$\sum_n |d_n|^2 \cdot (1+n^2)^{-s} = \sum_n |\overline{a_{-n}} \cdot (1+n^2)^s|^2 \cdot (1+n^2)^{-s} = \sum_n |a_n|^2 \cdot (1+n^2)^s < +\infty$$

[55] The Hilbert space ℓ^2 is the collection of sequences $\{a_1, a_2, \dots\}$ with $\sum |a_i|^2 < \infty$.

[56] Continuity of a linear functional λ on a Banach space B is equivalent to an estimate $|\lambda(b)|_{\mathbb{C}} \leq C \cdot \|b\|_B$ for some constant C .

Thus, $\ell^{2,-s}$ is (isomorphic to) the dual of $\ell^{2,s}$. ///

[10.0.4] **Claim:** The map $u \rightarrow \{\widehat{u}(n)\}$ on $H^s(S^1)$ by taking Fourier coefficients is a Hilbert-space isomorphism

$$H^s(S^1) \approx \ell^{2,s}$$

Proof: That the two-sided sequence of Fourier coefficients $u(\psi_{-n})$ is in $\ell^{2,s}$ is part of the definition of $H^s(S^1)$. The more serious question is *surjectivity*.

Let $\{c_n\} \in \ell^{2,s}$. For $s \geq 0$, the s^{th} Levi-Sobolev norm dominates the 0^{th} , so distributions in $H^s(S^1)$ are at least $L^2(S^1)$ -functions. The definition of $H^s(S^1)$ in this case makes $H^s(S^1)$ a Hilbert space, and we directly invoke the Plancherel theorem, using the orthonormal basis $\frac{\psi_n}{\sqrt{2\pi}} \cdot (1+n^2)^{-s/2}$ for $H^s(S^1)$. This gives the surjectivity $H^s(S^1) \rightarrow \ell^{2,s}$ for $s \geq 0$.

For $s < 0$, to prove the surjectivity, for $\{c_n\}$ in $\ell^{2,s}$ we will define a distribution u lying in $H^s(S^1)$, by

$$u(f) = \sum_n \widehat{f}(n) \cdot c_{-n} \quad (f \in C^\infty(S^1))$$

By Cauchy-Schwarz-Bunyakovsky,

$$\begin{aligned} \left| \sum_n \widehat{f}(n) \cdot c_{-n} \right| &\leq \sum_n |\widehat{f}(n)| (1+n^2)^{-s/2} \cdot |c_n| (1+n^2)^{s/2} \\ &\leq \left(\sum_n |\widehat{f}(n)|^2 (1+n^2)^{-s} \right)^{1/2} \cdot \left(\sum_n |c_n|^2 (1+n^2)^s \right)^{1/2} = |f|_{H^{-s}} \cdot |\{c_n\}|_{\ell^{2,s}} \end{aligned}$$

This shows that u is a continuous linear functional on $H^{-s}(S^1)$. For $s < 0$, the test functions $C^\infty(S^1)$ imbed continuously into $H^{-s}(S^1)$, so u gives a continuous functional on $C^\infty(S^1)$, so is a distribution. This proves that the Fourier coefficient map is a surjection to $\ell^{2,s}$ for $s < 0$. ///

[10.0.5] **Remark:** After this preparation, the remainder of this section is completely unsurprising. The following corollary is the conceptual point of this story.

[10.0.6] **Corollary:** For any $s \in \mathbb{R}$, the complex bilinear pairing

$$\langle \cdot, \cdot \rangle : H^s \times H^{-s} \rightarrow \mathbb{C} \quad \text{by} \quad f \times u \rightarrow \langle f, u \rangle = \sum_n \widehat{f}(n) \cdot \widehat{u}(-n)$$

gives an isomorphism

$$H^{-s} \approx (H^s)^*$$

by sending $u \in H^{-s}$ to $\lambda_u \in (H^s)^*$ defined by

$$\lambda_u(f) = \langle f, u \rangle \quad (\text{for } f \in H^s(S^1))$$

[10.0.7] **Remark:** The pairing of this last claim is *unsymmetrical*: the left argument is from H^s while the right argument is from H^{-s} .

Proof: This pairing via Fourier coefficients is simply the composition of the maps $H^s(S^1) \approx \ell^{2,s}$ and $H^{-s}(S^1) \approx \ell^{2,-s}$ with the pairing of $\ell^{2,s}$ and $\ell^{2,-s}$ given just above. ///

[10.0.8] **Corollary:** The space of all distributions on S^1 is

$$\text{distributions} = C^\infty(S^1)^* = \bigcup_{s \geq 0} H^s(S^1)^* = \bigcup_{s \geq 0} H^{-s}(S^1) = \text{colim}_{s \geq 0} H^{-s}(S^1)$$

thus expressing $C^\infty(S^1)^*$ as an ascending union of Hilbert spaces. ///

[10.0.9] Corollary: A distribution $u \sim \sum_n c_n \psi_n$ can be evaluated on $f \in C^\infty(S^1)$ by

$$u(f) = \sum_n \widehat{f}(n) \cdot \widehat{u}(-n)$$

Proof: Since u lies in some $H^{-s}(S^1)$, it gives a continuous functional on $H^s(S^1)$, which contains $C^\infty(S^1)$. The Plancherel-like evaluation formula above gives the equality. ///

A collection of Fourier coefficients $\{c_n\}$ is of moderate growth when there is a constant C and an exponent N such that

$$|c_n| \leq C \cdot (1 + n^2)^N \quad (\text{for all } n \in \mathbb{Z})$$

[10.0.10] Corollary: Let $\{c_n\}$ be a collection of complex numbers of moderate growth. Then there is a distribution u with those as Fourier coefficients, that is, there is u with

$$u(\psi_{-n}) = c_n$$

Proof: For constant C and exponent N such that $|c_n| \leq C \cdot (1 + n^2)^N$,

$$\sum_n |c_n|^2 \cdot (1 + n^2)^{-(2N+1)} \leq \sum_n C^2 \cdot (1 + n^2)^{2N} \cdot (1 + n^2)^{-(2N+1)} = C^2 \cdot \sum_n (1 + n^2)^{-1} < \infty$$

That is, from the previous discussion, the sequence gives an element of $H^{-(2N+1)}(S^1) \subset C^\infty(S^1)^*$. ///

[10.0.11] Corollary: For $u \sim \sum_n c_n \psi_n \in H^s(S^1)$ the derivative (for any $s \in \mathbb{R}$) is

$$u' \sim \sum_n in \cdot c_n \cdot \psi_n \in H^{s-1}$$

Proof: Invoke the definition (compatible with integration by parts) of the derivative of distributions, and integrating by parts to see that $\widehat{f}'(n) = in \cdot \widehat{f}(n)$ for $f \in C^\infty(S^1) = H^\infty(S^1)$,

$$u'(f) = -u(f') = -\sum_n \widehat{f}'(n) \cdot \widehat{u}(-n) = -\sum_n in \widehat{f}(n) \cdot \widehat{u}(-n) = \sum_n \widehat{f}(n) \cdot -in \widehat{u}(-n)$$

as claimed. The Fourier coefficients $-in \cdot \widehat{u}(n)$ do satisfy

$$\sum_n |in \widehat{u}(n)|^2 \cdot (1 + n^2)^{s-1} \leq \sum_n (1 + n^2) |\widehat{u}(n)|^2 \cdot (1 + n^2)^{s-1} = \sum_n |\widehat{u}(n)|^2 \cdot (1 + n^2)^s = |u|_{H^s}^2 < \infty$$

which proves that the differentiation maps H^s to H^{s-1} continuously. ///

[10.0.12] Remark: In the latter proof the sign in the subscript in the definition of the pairing $\ell^{2,s} \times \ell^{2,-s}$ was essential.

[10.0.13] Corollary: The collection of finite linear combinations of exponentials ψ_n is dense in every $H^s(S^1)$, for $s \in \mathbb{R}$. In particular, $C^\infty(S^1)$ is dense in every $H^s(S^1)$, for $s \in \mathbb{R}$.

Proof: The exponentials are an orthogonal basis for every Levi-Sobolev space. ///

[10.0.14] Remark: The topology of colimit of Hilbert spaces is the *finest* of several reasonable topologies on distributions. Density in a finer topology is a stronger assertion than density in a coarser topology.

11. The provocative example explained

The confusing example of the sawtooth function is clarified in the context we've developed. By now, we know that Fourier series whose coefficients satisfy sufficient decay conditions *are* differentiable. Even when the coefficients do not decay, but only grow *moderately*, the Fourier series is that of a *generalized function*. In other words, *we can (nearly) always differentiate* Fourier series term by term, as long as we can tolerate the outcome being a *generalized function*, rather than necessarily a *classical function*.

Again, $s(x)$ is the *sawtooth function*

$$s(x) = x - \pi \quad (\text{for } 0 \leq x < 2\pi)$$

made *periodic* by demanding $s(x + 2\pi n) = s(x)$ for all $n \in \mathbb{Z}$, so

$$s(x) = x - 2\pi \cdot \left\lfloor \frac{x}{2\pi} \right\rfloor - \pi \quad (\text{for } x \in \mathbb{R})$$

where $\lfloor x \rfloor$ is the greatest integer less than or equal x . Away from $2\pi\mathbb{Z}$, this function is infinitely differentiable, with derivative 1. At integers it jumps down from value to π to value $-\pi$. We do not attempt to define a value *at* $2\pi\mathbb{Z}$.

We want to *differentiate* this function compatibly with integration by parts, and compatibly with term-by-term differentiation of Fourier series.

The sawtooth function *is* well-enough behaved to give a *distribution* by integrating against it. Therefore, as we saw above, it *can* be differentiated as a distribution, and be correctly differentiated as (as a distribution) by differentiating its Fourier expansion termwise.

A earlier, Fourier coefficients are computed by integrating against e^{-inx}

$$\frac{1}{2\pi} \int_0^{2\pi} s(x) \cdot e^{-inx} dx = \begin{cases} \frac{1}{-in} & (\text{for } n \neq 0) \\ 0 & (\text{for } n = 0) \end{cases}$$

Thus, *at least* as a distribution, its Fourier expansion is

$$s(x) = i \sum_{n \neq 0} \frac{1}{n} \cdot e^{inx}$$

The series *does* converge pointwise to $s(x)$ for x away from (images of) integers, as we proved happens at left and right differentiable points for piecewise C^o functions.

We are entitled to differentiate, at worst within the class of distributions, within which we are assured of a reasonable sense to our computations. *Further*, we are entitled (for any distribution) to differentiate the Fourier series term-by-term. That is, as distributions,

$$\begin{aligned} s'(x) &= - \sum_{n \neq 0} e^{inx} \\ s''(x) &= - \sum_{n \neq 0} in e^{inx} \\ &\dots \\ s^{(k)}(x) &= - \sum_{n \neq 0} (in)^{k-1} e^{inx} \end{aligned}$$

and so on, just as successive derivatives of smooth functions $f(x) = \sum_n c_n e^{inx}$ are obtained by termwise differentiation

$$f^{(k)}(x) = \sum_{n \neq 0} (in)^k c_n e^{inx}$$

The difficulty of interpreting the right-hand side of the Fourier series for $s^{(k)}$ as having pointwise values is irrelevant.

More to the point, these Fourier series are things to integrate smooth functions against, by an extension of the Plancherel formula for inner products of L^2 functions. Namely, for any smooth function $f(x) \sim \sum_n c_n e^{inx}$, the imagined integral of f against $s^{(k)}$ should be expressible as the sum of products of Fourier coefficients

$$\text{imagined } \langle f, s^{(k)} \rangle = \sum_{n \neq 0} c_n \cdot \left(\frac{(in)^k}{-in} \right)^{\text{conj}}$$

(where $\alpha \rightarrow \alpha^{\text{conj}}$ is complex conjugation) and the latter expression should behave well when rewritten in a form that refers to the literal function s . Indeed,

$$\sum_{n \neq 0} c_n \cdot \left(\frac{(in)^k}{-in} \right)^{\text{conj}} = (-1)^k \sum_{n \neq 0} (in)^k c_n \cdot \left(\frac{1}{-in} \right)^{\text{conj}} = (-1)^k \int_{S^1} f^{(k)}(x) \bar{s}(x) dx$$

by the Plancherel theorem applied to the L^2 functions $f^{(k)}$ and s . Let u be the distribution given by integration against s . Then, by the definition of differentiation of distributions, we have computed that

$$(-1)^k \int_{S^1} f^{(k)}(x) \bar{s}(x) dx = (-1)^k u(f^{(k)}) = u^{(k)}(f)$$

It is in this sense that the sum $\sum_{n \neq 0} c_n \cdot \frac{(in)^k}{-in}$ is integration of s against f .

Further, for f a smooth function with support away from the discontinuities of s , it is true that $u''(f) = 0$, giving s'' a vague pointwise sense of being 0 away from the discontinuities of s . This was clear at the outset, but now is given precise meaning.

Thus, as claimed at the outset of the discussion of functions on the circle, we can differentiate $s(x)$ legitimately, and the differentiation of the Fourier series of the sawtooth function $s(x)$ correctly represents this differentiation, viewing $s(x)$ and its derivatives as *distributions*.

12. Appendix: products and limits of topological vector spaces

Here we carry out the diagrammatic proof that products and limits of topological vector spaces *exist*, and are locally convex when the factors or limitands are locally convex. Nothing surprising happens.

[12.0.1] **Claim:** Products and limits of topological vector spaces exist. In particular, limits are *closed* (linear) subspaces of the corresponding products. When the factors or limitands are locally convex, so is the product or limit.

[12.0.2] **Remark:** Part of the point is that products and limits of locally convex topological vector spaces *in the larger category of not-necessarily locally convex topological vector spaces* are nevertheless locally convex. That is, enlarging the category in which we take test objects does not change the outcome, in this case. By contrast, coproducts and colimits in general are sensitive to local convexity of the test objects. [57]

Proof: After we construct products, limits are constructed as closed subspaces of them.

Let V_i be topological vector spaces. We claim that the topological-space product $V = \prod_i V_i$ (with projections p_i) (with the product topology) is a topological vector space product. Let $\alpha_i : V_i \times V_i \rightarrow V_i$ be the addition on V_i . The family of composites $\alpha_i \circ (p_i \times p_i) : V \times V \rightarrow V_i$ induces a map $\alpha : V \times V \rightarrow V$ as in

$$\begin{array}{ccc} V \times V & \xrightarrow{\alpha} & V \\ p_i \times p_i \downarrow & & \downarrow p_i \\ V_i \times V_i & \xrightarrow{\alpha_i} & V_i \end{array}$$

This defines what we will show to be a *vector addition* on V . Similarly, the scalar multiplications $s_i : \mathbb{C} \times V_i \rightarrow V_i$ composed with the projections $p_i : V \rightarrow V_i$ give a family of maps

$$s_i \circ (1 \times p_i) : \mathbb{C} \times V \rightarrow V_i$$

which induce a map $s : \mathbb{C} \times V \rightarrow V$ which we will show to be a *scalar multiplication* on V . That these maps are *continuous* is given us by starting with the topological-space product.

That is, we must prove that vector addition is commutative and associative, that scalar multiplication is associative, and that the two have the usual distributivity. All these proofs are the same in form. For commutativity of vector addition, consider the diagram

$$\begin{array}{ccccc} & & V_i \times V_i & \xrightarrow{v \times w \rightarrow v+w} & V_i \\ & p_i \times p_i \nearrow & & & \nearrow p_i \\ V \times V & & & \xrightarrow{v \times w \rightarrow v+w} & V \\ & \xrightarrow{v \times w \rightarrow w+v} & & & \searrow p_i \\ & & V_i \times V_i & \xrightarrow{v \times w \rightarrow w+v} & V_i \\ & p_i \times p_i \searrow & & & \searrow p_i \\ & & V \times V & \xrightarrow{v \times w \rightarrow w+v} & V \end{array}$$

The upper half of the diagram is the induced-map definition of vector addition on V , and the lower half is the induced map definition of the reversed-order vector addition. The commutativity of addition on each V_i implies that going around the top of the diagram from $V \times V$ to V_i yields the same as going around the bottom. Thus, the two induced maps $V \times V \rightarrow V$ must be the same, since induced maps are *unique*.

[57] For example, uncountable coproducts do not exist among not-necessarily locally convex topological vector spaces, essentially because the not-locally-convex spaces ℓ^p with $0 < p < 1$ exist.

The proofs of associativity of vector addition, associativity of scalar multiplication, and distributivity, use the same idea. Thus, *products* of topological vector spaces exist.

We should not forget to prove that the product is *Hausdorff*, since we implicitly require this of topological vector spaces. This is immediate, since a (topological space) product of Hausdorff spaces is readily shown to be Hausdorff.

Consider now the case that each V_i is locally convex. By definition of the product topology, every neighborhood of 0 in the product is of the form $\Pi_i U_i$ where U_i is a neighborhood of 0 in V_i , and all but finitely many of the U_i are the whole V_i . Since V_i is locally convex, we can shrink every U_i that is *not* V_i to be a convex open containing 0, while each *whole* V_i is convex. Thus, the product is locally convex when every factor is.

To construct limits, reduce to the product.

[12.0.3] **Claim:** Let V_i be topological vector spaces with transition maps $\varphi_i : V_i \rightarrow V_{i-1}$. The limit $V = \lim_i V_i$ *exists*, and, in particular, is the closed linear subspace (with subspace topology) of the product $\Pi_i V_i$ (with projections p_i) defined by the (closed) conditions

$$\lim_i V_i = \{v \in \Pi_i V_i : (\varphi_i \circ p_i)(v) = p_{i-1}(v), \text{ for all } i\}$$

Proof: (of claim) Constructing the alleged limit as a closed subspace of the product immediately yields the desired properties of vector addition and scalar multiplication, as well as the Hausdorff-ness. What we must show is that the construction does function as a limit.

Given a compatible family of continuous linear maps $f_i : Z \rightarrow V_i$, there is induced a unique continuous linear map $F : Z \rightarrow \Pi_i V_i$ to the product, such that $p_i \circ f = f_i$ for all i . The *compatibility* requirement on the f_i exactly asserts that $f(Z)$ sits inside the subspace of $\Pi_i V_i$ defined by the conditions $(\varphi_i \circ p_i)(v) = p_{i-1}(v)$. Thus, f maps to this subspace, as desired.

Further, for all limitands locally convex, we have shown that the product is locally convex. The local convexity of a linear subspace (such as the limit) follows immediately. ///

13. Appendix: Fréchet spaces and limits of Banach spaces

A larger class of topological vector spaces arising in practice is the class of *Fréchet spaces*. In the present context, we can give a nice definition: a *Fréchet space* is a *countable* limit of Banach spaces. [58] Thus, for example,

$$C^\infty(S^1) = \bigcap_k C^k(S^1) = \lim_k C^k(S^1)$$

is a Fréchet space, *by (this) definition*.

Despite its advantages, the present definition is not the usual one. [59] We make a comparison, and elaborate

[58] Of course, it suffices that a limit have a countable cofinal subfamily.

[59] A common definition, with superficial appeal, is that a Fréchet space is a complete (invariantly) metrized space that is locally convex. This has the usual disadvantage that there are many different metrics that can give the same topology. This also ignores the manner in which Fréchet spaces usually arise, as countable limits of Banach spaces. There is another common definition that does halfway acknowledge the latter construction, namely, that a Fréchet space is a *complete* topological vector space with topology given by a countable collection of *seminorms*. The latter definition is essentially equivalent to ours, but requires explanation of the suitable notion of *completeness* in a not-necessarily metric situation, as well as explanation of the notion of *seminorm* and how topologies are specified by seminorms. We skirt the latter issues for the moment.

on the features of Fréchet spaces.

A metric $d(\cdot, \cdot)$ on a vector space V is *invariant* (implicitly, under addition), when

$$d(x + z, y + z) = d(x, y) \quad (\text{for all } x, y, z \in V)$$

All metrics we'll care about on topological vector spaces will be invariant in this sense.

[13.0.1] **Claim:** A Fréchet space is locally convex and complete (invariantly) metrizable. [60]

Proof: Let $V = \lim_i B_i$ be a countable limit of Banach spaces B_i , where $\varphi_i : B_i \rightarrow B_{i-1}$ are the transition maps and $p_i : V \rightarrow B_i$ are the projections. From the appendix, the limit is a closed linear subspace of the product, and the product is the cartesian product with the product topology and component-wise vector addition. Recall that a product of a *countable* collection of metric spaces is metrizable, and is complete if each factor is complete. A closed subspace of a complete metric space is complete metric. Thus, $\lim_i B_i$ is complete metric.

As proven in the previous appendix, *any* product or limit of locally convex spaces is locally convex, whether or not it has a countable cofinal family. Thus, the limit is Fréchet. ///

Addressing the comparison between local convexity and limits of Banach spaces,

[13.0.2] **Theorem:** Every locally convex topological vector space is a *subspace* of a limit of Banach spaces (and vice-versa).

[13.0.3] **Remark:** This little theorem encapsulates the construction of *semi-norms* to give a locally convex topology. It can also be used to reduce the general Hahn-Banach theorem for locally convex spaces to the Hahn-Banach theorem for Banach spaces.

Proof: In one direction, we already know that a product or limit of Banach spaces is locally convex, since Banach spaces are locally convex.

In the Banach or normed-space situation, the topology comes from a metric $d(v, w) = |v - w|$ defined in terms of a *single* function $v \rightarrow |v|$ with the usual properties

$$|\alpha \cdot v| = |\alpha|_{\mathbb{C}} \cdot |v| \quad (\text{homogeneity})$$

$$|v + w| \leq |v| + |w| \quad (\text{triangle inequality})$$

$$|v| \geq 0, \quad (\text{equality only for } v = 0) \quad (\text{definiteness})$$

By contrast, for more general (but locally convex) situations, we consider a *family* Φ of functions $p(v)$ for which the definiteness condition is weakened slightly, so we require

$$p(\alpha \cdot v) = |\alpha|_{\mathbb{C}} \cdot p(v) \quad (\text{homogeneity})$$

$$p(v + w) \leq p(v) + p(w) \quad (\text{triangle inequality})$$

$$p(v) \geq 0 \quad (\text{semi-definiteness})$$

Such a function $p(\cdot)$ is a *semi-norm*. For Hausdorff-ness, we further require that the family Φ is *separating* in the sense that, given $v \neq 0$ in V , there is $p \in \Phi$ such that $p(v) > 0$.

[60] As is necessary to prove the *equivalence* of the various definitions of *Fréchet space*, the converse of this claim is true, namely, that every locally convex and complete (invariantly) metrizable topological vector space is a countable limit of Banach spaces. Proof of the converse requires work, namely, development of ideas about seminorms. Since we don't need this converse at the moment, we do not give the argument.

A separating family Φ of semi-norms on a complex vector space V gives a *locally convex* topology by taking as local sub-basis^[61] at 0 the sets

$$U_{p,\varepsilon} = \{v \in V : p(v) < \varepsilon\} \quad (\text{for } \varepsilon > 0 \text{ and } p \in \Phi)$$

Each of these is convex, because of the triangle inequality for the semi-norms.

[13.0.4] Remark: The topology obtained from a (separating) family of seminorms may appear to be a random or frivolous generalization of the notion of topology obtained from a *norm*. However, it is the correct extension to encompass *all* locally convex topological vector spaces, as we see now.^[62]

For a locally convex topological vector space V , for every open U in a local basis B at 0 of *convex* opens, try to define a *seminorm*

$$p_U(v) = \inf\{t > 0 : t \cdot U \ni v\}$$

We discover some necessary adjustments, and then verify the semi-norm properties.

First, we show that for any $v \in V$ the set over which the inf is taken is non-empty. Since scalar multiplication $\mathbb{C} \times V \rightarrow V$ is (jointly!) continuous, for given $v \in V$, given a neighborhood U of $0 \in V$, there are neighborhoods W of $0 \in \mathbb{C}$ and U' of v such that

$$\alpha \cdot w \in U \quad (\text{for all } \alpha \in W \text{ and } w \in U')$$

In particular, since W contains a disk $\{|\alpha| < \varepsilon\}$ for some $\varepsilon > 0$, we have $t \cdot v \in U$ for all $0 < t < \varepsilon$. That is,

$$v \in t \cdot U \quad (\text{for all } t > \varepsilon^{-1})$$

Semi-definiteness of p_U is built into the definition.

To avoid nagging problems, we should verify that, for convex U containing 0, when $v \in t \cdot U$ then $v \in s \cdot U$ for all $s \geq t$. This follows from the convexity, by

$$s^{-1} \cdot v = \frac{t}{s} \cdot (t^{-1} \cdot v) = \frac{t}{s} \cdot (t^{-1} \cdot v) + \frac{s-t}{s} \cdot 0 \in U$$

since $t^{-1} \cdot v$ and 0 are in U .

The homogeneity condition $p(\alpha v) = |\alpha| p(v)$ already presents a minor issue, since convex sets containing 0 need have no special properties regarding multiplication by complex numbers. That is, the problem is that, given $v \in t \cdot U$, while $\alpha v \in \alpha \cdot t \cdot U$, we do *not* know that this implies $\alpha v \in |\alpha| \cdot t \cdot U$. Indeed, in general, it will not. To repair this, to make semi-norms we must use only convex opens U which are *balanced* in the sense that

$$\alpha \cdot U = U \quad (\text{for } \alpha \in \mathbb{C} \text{ with } |\alpha| = 1)$$

Then, given $v \in V$, we have $v \in t \cdot U$ if and only if $\alpha v \in t \cdot \alpha U$, and now

$$t \alpha U = t |\alpha| \frac{\alpha}{|\alpha|} U = t |\alpha| U$$

by the balanced-ness.

[61] Again, a *sub-basis* for a topology is a set of opens such that finite intersections form a *basis*. In other words, arbitrary unions of finite intersections give all opens.

[62] The semi-norms we construct here are sometimes called *Minkowski functionals*, even though they are not functionals in the sense of being continuous linear maps.

Now we have an obligation to show that there is a local basis (at 0) of convex *balanced* opens. Fortunately, this is easy to see, as follows. Given a convex U containing 0, from the continuity of scalar multiplication, since $0 \cdot v = 0$, there is $\varepsilon > 0$ and a neighborhood W of 0 such that $\alpha \cdot w \in U$ for $|\alpha| < \varepsilon$ and $w \in W$. Let

$$U' = \{\alpha \cdot w : |\alpha| \leq \frac{\varepsilon}{2}, w \in W\} = \bigcup_{|\alpha| \leq \varepsilon/2} \alpha \cdot W$$

Being a union of the opens $\alpha \cdot W$, this U' is open. It is inside U by arrangement, and is *balanced* by construction. That is, there is indeed a local basis of convex *balanced* opens at 0.

For the *triangle inequality* for p_U , given $v, w \in V$, let t_1, t_2 be such that $v \in t_1 \cdot U$ for $t_1 \geq t_1$ and $w \in t_2 \cdot U$ for $t_2 \geq t_2$. Then, using the convexity,

$$v + w \in t_1 \cdot U + t_2 \cdot U = (t_1 + t_2) \cdot \left(\frac{t_1}{t_1 + t_2} \cdot U + \frac{t_2}{t_1 + t_2} \cdot U \right) \subset (t_1 + t_2) \cdot U$$

This gives the triangle inequality

$$p_U(v + w) \leq p_U(v) + p_U(w)$$

Finally, we check that the semi-norm topology is the original one. This is unsurprising. It suffices to check at 0. On one hand, given an open W containing 0 in V , there is a convex, balanced open U contained in W , and

$$\{v \in V : p_U(v) < 1\} \subset U \subset W$$

Thus, the semi-norm topology is at least as fine as the original topology. On the other hand, given convex balanced open U containing 0, and given $\varepsilon > 0$,

$$\{v \in V : p_U(v) < \varepsilon\} \supset \frac{\varepsilon}{2} \cdot U$$

Thus, each sub-basis open for the semi-norm topology contains an open in the original topology. We conclude that the two topologies are the same.

A summary so far: for a locally convex topological vector space, the semi-norms attached to convex balanced neighborhoods of 0 give a topology identical to the original, and *vice-versa*.

Before completing the proof of the theorem, recall that a *completion* of a set with respect to a *pseudo-metric* can be defined much as the completion with respect to a genuine metric. This is relevant because a semi-norm may only give a pseudo-metric, not a genuine metric.

Let Φ be a (separating) family of seminorms on a vector space V . For a *finite subset* i of Φ , let X_i be the *completion* of V with respect to the semi-norm

$$p_i(v) = \sum_{p \in i} p(v)$$

with natural map $f_i : V \rightarrow X_i$. Order subsets of Φ by $i \geq j$ when $i \supset j$. For $i > j$ we have

$$p_i(v) = \sum_{p \in i} p(v) \geq \sum_{p \in j} p(v) = p_j(v)$$

so we have natural continuous (transition) maps

$$\varphi_{ij} : X_i \longrightarrow X_j \quad (\text{for } i > j)$$

We claim that each X_i is a *Banach space*, that V with its semi-norm topology has a natural continuous *inclusion* to the limit $X = \lim_i X_i$, and that V has the topology given by the subspace topology inherited from the limit.

The maps f_i form a compatible family of maps to the X_i , so there is a unique compatible map $f : V \rightarrow X$. By the separating property, given $v \neq 0$, there is $p \in \Phi$ such that $p(v) \neq 0$. Then for all i containing p , we have $f_i(v) \neq 0 \in X_i$. The subsets i containing p are *cofinal* in this limit, so $f(v) \neq 0$. Thus, f is an inclusion.

Since the limit is a (closed) subspace of the *product* of the X_i , it suffices to prove that the topology on V (imbedded in $\Pi_i X_i$ via f) is the subspace topology from $\Pi_i X_i$. Since the topology on V is at *least* this fine (since f is continuous), we need only show that the *subspace* topology is at least as fine as the semi-norm topology. To this end, consider a semi-norm-topology sub-basis set

$$\{v \in V : p_U(v) < \varepsilon\} \quad (\text{for } \varepsilon > 0 \text{ and convex balanced open } U \text{ containing } 0)$$

This is simply the intersection of $f(V)$ with the sub-basis set

$$\Pi_{p \neq \{p_U\}} X_i \times \{v \in X_{\{p_U\}} : p_U(v) < \varepsilon\}$$

with the last factor inside $X_{\{p_U\}}$. Thus, by construction, the map $f : V \rightarrow X$ is a homeomorphism of V to its image. ///

[13.0.5] **Remark:** We do *not* assert that an arbitrary locally convex topological vector space is a limit of Banach spaces. Such an assertion is equivalent to various *completeness* hypotheses, which we will investigate later.

14. Appendix: Urysohn and density of C^0

Urysohn's lemma is the technical point that allows us to relate *measurable* functions to *continuous* functions. Then, from the Lebesgue definition of *integral*, we can prove the density of continuous functions in L^2 spaces, for example.

[14.0.1] **Theorem:** (*Urysohn*) In a locally compact Hausdorff topological space X , given a compact subset K contained in an open set U , there is a continuous function $0 \leq f \leq 1$ which is 1 on K and 0 off U .

Proof: First, we prove that there is an open set V such that

$$K \subset V \subset \bar{V} \subset U$$

For each $x \in K$ let V_x be an open neighborhood of x with compact closure. By compactness of K , some finite subcollection V_{x_1}, \dots, V_{x_n} of these V_x cover K , so K is contained in the open set $W = \bigcup_i V_{x_i}$ which has compact closure $\bigcup_i \bar{V}_{x_i}$ since the union is *finite*.

Using the compactness again in a similar fashion, for each x in the closed set $X - U$ there is an open W_x containing K and a neighborhood U_x of x such that $W_x \cap U_x = \emptyset$.

Then

$$\bigcap_{x \in X - U} (X - U) \cap \bar{W} \cap \bar{W}_x = \emptyset$$

These are compact subsets in a Hausdorff space, so (again from compactness) some *finite* subcollection has empty intersection, say

$$(X - U) \cap (\bar{W} \cap \bar{W}_{x_1} \cap \dots \cap \bar{W}_{x_n}) = \emptyset$$

That is,

$$\bar{W} \cap \bar{W}_{x_1} \cap \dots \cap \bar{W}_{x_n} \subset U$$

Thus, the open set

$$V = W \cap W_{x_1} \cap \dots \cap W_{x_n}$$

meets the requirements.

Using the possibility of inserting an open subset and its closure between any $K \subset U$ with K compact and U open, we inductively create opens V_r (with compact closures) indexed by rational numbers r in the interval $0 \leq r \leq 1$ such that, for $r > s$,

$$K \subset V_r \subset \bar{V}_r \subset V_s \subset \bar{V}_s \subset U$$

From any such configuration of opens we construct the desired continuous function f by

$$f(x) = \sup\{r \text{ rational in } [0, 1] : x \in V_r, \} = \inf\{r \text{ rational in } [0, 1] : x \in \bar{V}_r, \}$$

It is not immediate that this sup and inf are the same, but if we *grant* their equality then we can prove the *continuity* of this function $f(x)$. Indeed, the sup description expresses f as the supremum of characteristic functions of open sets, so f is at least *lower semi-continuous*.^[63] The inf description expresses f as an infimum of characteristic functions of closed sets so is *upper semi-continuous*. Thus, f would be continuous.

To finish the argument, we must construct the sets V_r and prove equality of the inf and sup descriptions of the function f .

To construct the sets V_i , start by finding V_0 and V_1 such that

$$K \subset V_1 \subset \bar{V}_1 \subset V_0 \subset \bar{V}_0 \subset U$$

Fix a well-ordering r_1, r_2, \dots of the rationals in the open interval $(0, 1)$. Supposing that V_{r_1}, \dots, V_{r_n} have been chosen. let i, j be indices in the range $1, \dots, n$ such that

$$r_j > r_{n+1} > r_i$$

and r_j is the *smallest* among r_1, \dots, r_n above r_{n+1} , while r_i is the *largest* among r_1, \dots, r_n below r_{n+1} . Using the first observation of this argument, find $V_{r_{n+1}}$ such that

$$V_{r_j} \subset \bar{V}_{r_j} \subset V_{r_{n+1}} \subset \bar{V}_{r_{n+1}} \subset V_{r_i} \subset \bar{V}_{r_i}$$

This constructs the nested family of opens.

Let $f(x)$ be the sup and $g(x)$ the inf of the characteristic functions above. If $f(x) > g(x)$ then there are $r > s$ such that $x \in V_r$ and $x \notin \bar{V}_s$. But $r > s$ implies that $V_r \subset \bar{V}_s$, so this cannot happen. If $g(x) > f(x)$, then there are rationals $r > s$ such that

$$g(x) > r > s > f(x)$$

Then $s > f(x)$ implies that $x \notin V_s$, and $r < g(x)$ implies $x \in \bar{V}_r$. But $V_r \subset \bar{V}_s$, contradiction. Thus, $f(x) = g(x)$. ///

[14.0.2] Corollary: For a topological space X with a regular Borel measure μ , $C_c^o(X)$ is dense in $L^2(X, \mu)$.

Proof: The *regularity* of the measure is the property that $\mu(E)$ is both the *sup* of $\mu(K)$ for compacts $K \subset E$, and is the *inf* of $\mu(U)$ for opens $U \supset E$. From Urysohn's lemma, we have a continuous $f_{K,U}(x)$ which is 1 on K and 0 off U . Let $K_1 \subset K_2 \subset \dots$ be a sequence of compacts inside E whose measure approaches that of E from *below*, and let $U_1 \supset U_2 \supset \dots$ be a sequence of opens containing E whose measures approach that of E

[63] A (real-valued) function f is *lower semi-continuous* when for all bounds B the set $\{x : f(x) > B\}$ is open. The function f is *upper semi-continuous* when for all bounds B the set $\{x : f(x) < B\}$ is open. It is easy to show that a sup of lower semi-continuous functions is lower semi-continuous, and an inf of upper semi-continuous functions is upper semi-continuous. As expected, a function both upper and lower semi-continuous is continuous.

from *above*. Let f_i be a function as in Urysohn's lemma made to be 1 on K_i and 0 off U_i . Then Lebesgue's Dominated convergence theorem implies that

$$f_i \longrightarrow (\text{characteristic function of } E) \quad (\text{in } L^2(X, \mu))$$

From the definition of integral of *measurable functions*, finite linear combinations of characteristic functions are dense in L^2 (or any other L^p with $1 \leq p < \infty$). Thus, continuous functions are dense. ///
