

(March 12, 2020)

04. Representations of GL_2 and SL_2 over finite fields

Paul Garrett garrett@math.umn.edu <http://www.math.umn.edu/~garrett/>

[This document is http://www.math.umn.edu/~garrett/m/mfms/toward_SSW/04_finite_GL2.pdf]

1. Principal series representations of GL_2
2. Whittaker functionals, Whittaker models
3. Uniqueness of Whittaker functionals/models
4. Summary for GL_2
5. Conjugacy classes in SL_2 , odd q
6. Irreducibles of SL_2 , odd q

Irreducible complex representations of GL_2 and SL_2 over a finite field can be studied by methods applicable to p -adic reductive groups and real Lie groups.

We mostly resist using techniques too-special to finite groups and finite-dimensional representations, to practice more broadly applicable techniques. *Dimension* and *cardinality* are invoked as seldom as possible, although used when convenient. *complete reducibility* for finite-dimensional representations of finite groups is used only to add clarity.

1. Principal series representations of GL_2

Let $k = \mathbb{F}_q$ be a finite field with q elements. Let $G = GL_2(k)$ or $G = SL_2(k)$. Let

$$P = \left\{ \begin{pmatrix} * & * \\ 0 & * \end{pmatrix} \in G \right\} \quad N = \left\{ \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \in G \right\} \quad M = \left\{ \begin{pmatrix} * & 0 \\ 0 & * \end{pmatrix} \in G \right\} \quad w_o = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

The subgroup P is the *standard parabolic* subgroup, N its *unipotent radical*, and M the standard *Levi component* of P . The subgroup P is the semidirect product of M and N , with M normalizing N . This w_o is the *longest Weyl element*.

The important family of representations of G is the *principal series* of representations I_χ of G parametrized by characters (meaning one-dimensional representations)

$$\chi : M \longrightarrow \mathbb{C}^\times$$

For $G = SL_2$, $M \approx k^\times$ so these characters are characters χ_1 of k^\times via

$$\chi \left(\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \right) = \chi_1(a)$$

For $G = GL_2$, $M \approx k^\times \times k^\times$ and these characters are pairs (χ_1, χ_2) of characters of k^\times via

$$\chi \left(\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix} \right) = \chi_1(a)\chi_2(d)$$

In either case, extend χ to P by being identically 1 on N . The χ^{th} *principal series* representation of G attached to χ is the \mathbb{C} -vectorspace of functions

$$I_\chi = \text{Ind}_P^G \chi = \{ \mathbb{C}\text{-valued functions } f \text{ on } G : f(pg) = \chi(p) f(g) \text{ for all } p \in P, g \in G \}$$

The *action* of G on $\text{Ind}_P^G \chi$ is by the *right regular* representation

$$(R_g f)(x) = f(xg)$$

[1.1] **Remark:** An important aspect of representations of G induced from representations of subgroups is that they are *constructed*, so *exist*. One would hope to construct many, if not all, *irreducibles* by this process. We see below that *most* principal series, namely, those with the *regularity* property $\chi(wmw^{-1}) \neq \chi(m)$ for $m \in M$, are *irreducible*, and these irreducibles are about half of all irreducibles of G .

Induced representations have a convenient feature, namely, [1]

[1.2] **Theorem:** (*Frobenius Reciprocity*) For a representation σ of a subgroup H of G , and for a representation V of G , there is a natural isomorphism

$$F : \text{Hom}_G(V, \text{Ind}_H^G \sigma) \approx \text{Hom}_H(\text{Res}_H^G V, \sigma)$$

of \mathbb{C} -vectorspaces, where Res_H^G is the forgetful functor. The isomorphism F is

$$F(\Phi)(v) = \Phi(v)(1_G) \quad (\text{for } v \in V)$$

with inverse

$$(F^{-1}(\varphi)(v))(g) = R_g(\varphi(v))$$

(Given the formulas, the proof is straightforward.)

///

For a *one-dimensional* (for simplicity) irreducible $\sigma : H \rightarrow \mathbb{C}^\times$ of a group H , a σ -*isotypic* representation V of H is a (possibly large) representation V of H on which H acts entirely by σ , in the sense that

$$h \cdot v = \sigma(h) \cdot v \quad (\text{for all } v \in V, h \in H)$$

For a representation V of H , the (H, σ) -*isotype* $V^{H, \sigma}$ of V is the smallest H -subrepresentation $i : V^{H, \sigma} \rightarrow V$ of V such that any H -morphism

$$\varphi : W \rightarrow V$$

of a σ -isotypic H -representation W uniquely factors through $V^{H, \sigma}$, namely there is a unique $\varphi_o : W \rightarrow V^{H, \sigma}$ such that

$$\varphi = i \circ \varphi_o : W \rightarrow V^{H, \sigma} \rightarrow V$$

Existence of the isotype is proven by a readily-verifiable *construction*:

$$V^{H, \sigma} = \sum_{\varphi: \sigma \rightarrow V} \text{Im } \varphi$$

Dually, the (H, σ) -*co-isotype* $V_{H, \sigma}$ of a representation V of H is the smallest H -quotient of V such that any H -homomorphism $\varphi : V \rightarrow W$ with W σ -isotypic factors through $V_{H, \sigma}$. A *construction* readily shown to meet the characterizing requirement is

$$V_{H, \sigma} = V / \bigcap_{\varphi: V \rightarrow \sigma} \ker \varphi$$

In the special case of the trivial representation $\sigma = 1$ of H , the $(H, 1)$ -isotype is the sub-module of *H -fixed vectors* in a G -representation V :

$$H\text{-fixed vectors in } V = V^H = V^{H, 1} = \{v \in V : h \cdot v = v, \text{ for all } h \in H\}$$

[1] Indeed, in the long run, it is better to *characterize* the induced representation as making Frobenius Reciprocity hold, rather than *constructing* it and then proving that it has the property. Frobenius Reciprocity is an instance of an *adjunction relation* for *adjoint functors*.

The $H, 1$ -co-isotype is the quotient of H -co-fixed vectors:

$$H\text{-co-fixed vectors} = V_H = V_{H,1}$$

[1.3] **Remark:** That isotype and co-isotype are *functors* is the assertion that, in addition to transforming *objects* (G -representations), the *morphisms* (G -intertwining operators) are transformed compatibly. For example, given G -intertwining operator $\varphi : V \rightarrow W$ of G -representations, there is an H -intertwining operator $\varphi^{H,\sigma} : V^{H,\sigma} \rightarrow W^{H,\sigma}$, and *composition* of intertwining operators is preserved. In the cases at hand, this is easy to check.

[1.4] **Remark:** In situations where complete reducibility holds, co-isotypes are subrepresentations, thus are also *isotypes*. Nevertheless, *in general*, co-isotypes are *not* isotypes, and the distinction is meaningful.

As usual, a representation V of G is a representation of the subgroup N , by the forgetful functor $V \rightarrow \text{Res}_N^G$. A terminological convention is that *the* trivial representation of N is a *one-dimensional* vector space on which N acts trivially. [2]

Changing the notation slightly from the previous paragraph, the *Jacquet module* of V is

$$\text{Jacquet module } J_N V \text{ of } V = \text{co-isotype for trivial representation of } N = V_N$$

The *Jacquet functor* J_N is

$$J_N : V \longrightarrow V_N$$

[1.5] **Proposition:** For a G -representation V , the kernel of the Jacquet map $J_N : V \rightarrow V_N$ is generated by all expressions

$$v - n \cdot v$$

for $v \in V$ and $n \in N$. Also,

$$\ker J_N = \{v \in V : \sum_{n \in N} n \cdot v = 0\}$$

Proof: Under *any* N -map $r : V \rightarrow W$ with N acting trivially on W ,

$$r(v - nv) = rv - r(nv) = rv - n(rv) = rv - rv = 0$$

so the elements $v - nv$ are in the kernel of the quotient map to the Jacquet module. On the other hand, the linear span of these elements is stable under N , so we may form the quotient of V by these elements. This proves that the first description of the kernel is correct.

To prove the second characterization, suppose that

$$\sum_{n \in N} n \cdot v = 0$$

Then

$$v = v - 0 = v - \frac{1}{\#N} \sum_{n \in N} n \cdot v = \frac{1}{\#N} \sum_{n \in N} (v - n \cdot v)$$

a finite sum, expressing v as a linear combination of the desired form. On the other hand,

$$\sum_{n \in N} n \cdot (v - n_o \cdot v) = \sum_{n \in N} n \cdot v - \sum_{n \in N} (nn_o \cdot v) = \sum_{n \in N} n \cdot v - \sum_{n \in N} n \cdot v = 0$$

[2] In contrast to *the* one-dimensional trivial representation, a trivial representation of N is an *arbitrary-dimension* space on which N acts trivially.

by changing variables in the second integral. ///

[1.6] **Corollary:** The Jacquet functor is a functor from G -representations to M -representations.

Proof: The kernel of $V \rightarrow V_N$ is M -stable, by direction computation

$$m \cdot (v - n \cdot v) = mv - mnmm^{-1} \cdot mv = mv - n' \cdot mv \quad (\text{with } n' = mnmm^{-1} \in N)$$

since M normalizes N . ///

We will suppress the notation Res_H^G for a forgetful restriction functor, when its application is clear from context.

[1.7] **Corollary:** For G representations V there is a natural \mathbb{C} -linear isomorphism

$$\text{Hom}_G(V, I_\chi) \approx \text{Hom}_M(V_N, \chi)$$

(Combine the characterization of the Jacquet module with Frobenius Reciprocity.) ///

[1.8] **Corollary:** A (non-zero) irreducible representation V of G with $V_N \neq \{0\}$ is isomorphic to a subrepresentation of a principal series I_χ for some χ . On the other hand, if $V_N = \{0\}$, then V is *not* isomorphic to a subrepresentation of *any* principal series.

Proof: The representation space V_N of the group M is finite-dimensional, so by induction on dimension has an *irreducible* quotient $\varphi : V_N \rightarrow \chi$ for some representation χ of M , one-dimensional because M is abelian. In particular, this map φ is not 0. Thus, via the inverse L^{-1} of the isomorphism

$$L : \text{Hom}_G(V, I_\chi) \approx \text{Hom}_M(V_N, \chi)$$

we obtain a non-zero $T^{-1}\varphi \in \text{Hom}_G(V, I_\chi)$. ///

[1.9] **Remark:** A representation V is *supercuspidal* when $V_N = \{0\}$. Thus, by the above corollary, irreducibles of G *either* imbed into a principal series *or* are supercuspidal. For larger groups such as $GL(3, k)$ and $SL(3, k)$ there are intermediate cases.

The next issue is assessment of the irreducibility of the principal series I_χ , proving below that I_χ is *irreducible* when χ is *regular*, meaning that $\chi^w \neq \chi$, where for $m \in M$ the character χ^w is

$$\chi^w(m) = \chi(wmw^{-1})$$

The connection between imbeddability into principal series and non-vanishing of the Jacquet functor continues to be relevant.

First, *if* V were a *proper* subrepresentation of a principal series representation I_χ , *then* the quotient I_χ/V would be non-zero, and by induction on dimension would have an irreducible quotient π . Showing $\pi_N \neq 0$ would show (from above) that π is a subrepresentation of some principal series I_β , giving a non-zero G -intertwining $I_\chi \rightarrow I_\beta$.

[1.10] **Theorem:** The Jacquet functor $J_N : V \rightarrow V_N$ is an *exact functor*, meaning that for $f : A \rightarrow B$ and $g : B \rightarrow C$ are maps such that

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

is a short exact sequence of G -representations, the induced maps on Jacquet modules give an exact sequence

$$0 \rightarrow A_N \rightarrow B_N \rightarrow C_N \rightarrow 0$$

[1.11] **Remark:** This theorem can be interpreted as asserting that the group homology of N is always trivial (above degree 0), in the following sense. Even in a somewhat larger context, it is true that co-isotype functors are *right* exact and isotype functors are *left* exact, for reasons noted in the proof below. Given a projective resolution

$$\dots \xrightarrow{d} F^2 \xrightarrow{d} F^1 \xrightarrow{d} F^0 \rightarrow V \rightarrow 0$$

of an N -representation V (by N -representations F^i) the *group homology* of V the homology

$$H_n(V) = \frac{\ker d \text{ on } F_N^n}{d(F_N^{n+1})}$$

of the sequence

$$\dots \xrightarrow{d} F_N^2 \xrightarrow{d} F_N^1 \xrightarrow{d} F_N^0 \rightarrow 0$$

where, in particular, $H_0(V) = V_N$. That is, the higher group homology modules are the *left derived functors* of the (trivial representation) co-isotype functor. The long exact sequence attached to a short exact sequence

$$0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$$

is

$$\dots \rightarrow H_2(A) \rightarrow H_2(B) \rightarrow H_2(C) \xrightarrow{\delta} H_1(A) \rightarrow H_1(B) \rightarrow H_1(C) \xrightarrow{\delta} H_0(A) \rightarrow H_0(B) \rightarrow H_0(C) \rightarrow 0$$

From this and $H_0(V) = V_N$ we have the universal result that

$$H_1(B) \rightarrow H_1(C) \xrightarrow{\delta} A_N \rightarrow B_N \rightarrow C_N \rightarrow 0$$

is exact. Similar remarks apply to isotypes and cohomology.

Proof: The right half-exactness is a general property of co-isotype functors, as special cases of *left adjoints* to *isotype* functors. [3] That is, the surjectivity of $g : B_N \rightarrow C_N$ follows from that of $q \circ g : B \rightarrow C_N$ by a very general mechanism. Likewise, since the composite $g \circ f : A \rightarrow C$ is 0, certainly

$$q \circ g \circ f : A \rightarrow C_N$$

is 0, so the composite $A_N \rightarrow B_N \rightarrow C_N$ is 0.

The injectivity of $A_N \rightarrow B_N$ and the fact that the image of A_N in B_N is the whole kernel of $B_N \rightarrow C_N$ are less general, using here the finiteness of the group N . Let $a \in A$ such that $q(fa) = 0 \in B_N$. Then

$$\sum_{n \in N} n \cdot fa = 0$$

Since f commutes with the action of N , this gives

$$f \left(\sum_{n \in N} n \cdot a \right) = 0$$

By the injectivity of f

$$\sum_{n \in N} n \cdot a = 0$$

[3] The fact that the right-exactness instantiates a general property of co-isotypes does not mean that the proof is trivial. The *left*-exactness of *isotypes* $V \rightarrow V^\sigma$ is easier to prove.

so $qa = 0 \in A_N$. This proves exactness at the left joint.

When $g(qb) = 0$, $q(gb) = 0$, so

$$\sum_{n \in N} n \cdot gb = 0$$

and then the N -homomorphism property of g , namely $ng = gn$, gives

$$g\left(\sum_{n \in N} n \cdot b\right) = 0$$

Thus, the integral is in the kernel of g , so is in the image of f . Let $a \in A$ be such that

$$fa = \sum_{n \in N} n \cdot b$$

Without loss of generality, $\text{meas}(N) = 1$. Then

$$\sum_{n \in N} n' \cdot fa' = \sum_{n \in N} \sum_{n \in N} n'n \cdot b' = \sum_{n \in N} \sum_{n \in N} n \cdot b'$$

by replacing n by $n'^{-1}n$. This gives

$$\sum_{n \in N} n \cdot (fa - b) = 0$$

Thus, $q(fa - b) = 0$ and $f(qa) = qb$. This finishes the proof of exactness at the middle joint. ///

As usual, the \mathbb{C} -linear *dual* or *contragredient* representation V^\vee of a G -representation V is the dual \mathbb{C} -vector space with the action

$$(g \cdot \lambda)(v) = \lambda(g^{-1} \cdot v)$$

for $\lambda \in V^\vee$, $v \in V$, and $g \in G$. Being more careful, since there are two different representations involved, let (π, V) be the given representation, and (π^\vee, V^\vee) the dual. The definition of π^\vee is

$$(\pi^\vee(g)(\lambda))(v) = \lambda(\pi(g^{-1})(v))$$

The mapping $v \times \lambda \rightarrow \lambda(v)$ will also be denoted by

$$v \times \lambda \rightarrow \lambda(v) = \langle v, \lambda \rangle \quad (\text{implicitly } \mathbb{C}\text{-bilinear})$$

Recall:

[1.12] Proposition: V is irreducible if and only if V^\vee is irreducible.

Proof: If V has a proper subrepresentation U , then the inclusion $U \rightarrow V$ yields a *surjection* $V^\vee \rightarrow U^\vee$. Since U is non-zero and is not all of V there is a functional identically 0 on U but not identically 0 on V . Thus, the latter surjection has a proper kernel, which is a proper subrepresentation of V^\vee . On the other hand, the same argument shows that for a proper subrepresentation Λ of V^\vee there is $x \in V^{\vee\vee}$ vanishing identically on Λ but not identically vanishing on V^\vee . The finite-dimensionality implies that the natural inclusion $V \subset V^{\vee\vee}$ is an isomorphism. ///

[1.13] Proposition: For a G -representation V , let $J_N^\vee : (V_N)^\vee \rightarrow V^\vee$ be the natural dual M -map $\mu \rightarrow J_N \circ \mu$ obtained from $J_N : V \rightarrow V_N$. Then we have an isomorphism

$$J_N \circ J_N^\vee : (V_N)^\vee \rightarrow (V^\vee)_N \quad (\text{the latter } J_N \text{ is } V^\vee \rightarrow (V^\vee)_N)$$

Proof: In fact, $\mu \rightarrow \mu \circ J_N$ injects $(V_N)^\vee$ to the subspace $(V^\vee)^N$ of N -fixed vectors in V^\vee , since for $n \in N$ and $v \in V$ we directly compute

$$(n \cdot (\mu \circ J_N))(v) = (\mu \circ J_N)(nv) = \mu(J_N(nv)) = \mu(J_N(v)) = (\mu \circ J_N)(v)$$

The N -fixed vectors $(V^\vee)^N$ inject to $(V^\vee)_N$, since for an N -fixed vector λ

$$\sum_{n \in N} n \cdot \lambda = \sum_{n \in N} \lambda = \text{meas}(N) \cdot \lambda$$

(invoking the description above of the kernel of the quotient map to the Jacquet module). Thus, $J_N \circ J_N^\vee$ is an injection.

At this point we use a special feature to prove that the map is an isomorphism. Since finite-dimensional spaces are *reflexive*, apply the previous argument to V^\vee in place of V to obtain

$$((V^\vee)_N)^\vee \rightarrow (V^{\vee\vee})_N \approx V_N$$

Generally, when $X \rightarrow Y$ is injective and $Y^\vee \rightarrow X^\vee$ is injective, both maps are isomorphisms, so we have the desired result. ///

This allows us to prove a result complementary to the earlier assertion that irreducibles V are subrepresentations of principal series if and only if $V_N \neq 0$. First, another useful property of induced representations:

[1.14] Proposition: For a finite-dimensional representation σ of a subgroup K of a finite group H , the \mathbb{C} -linear dual of the induced representation $\text{Ind}_K^H \sigma$ is

$$\left(\text{Ind}_K^H \sigma\right)^\vee \approx \text{Ind}_K^H (\sigma^\vee)$$

via the pairing

$$\langle f, \lambda \rangle = \sum_{h \in K \backslash H} \langle f(h), \lambda(h) \rangle_\sigma \quad (\langle \cdot, \cdot \rangle_\sigma \text{ the pairing on } \sigma \times \sigma^\vee)$$

Proof: By definition of the dual representation, the function

$$h \rightarrow \langle f(h), \lambda(h) \rangle_\sigma$$

is left K -invariant, so gives a function on the quotient $K \backslash H$. To complete the proof we must use special features, the finiteness of H and the reflexiveness of σ . Consider functions f and λ supported on single points in $K \backslash H$, with values in dual bases of σ and σ^\vee . These form dual bases for the indicated induced representations. ///

[1.15] Corollary: For V an irreducible quotient of a principal series, $V_N \neq 0$ and V imbeds into a principal series.

Proof: Consider a surjection

$$\text{Ind}_P^G \chi \xrightarrow{\varphi} V$$

By dualizing, and by the previous proposition, we have an injection

$$\varphi^\vee : V^\vee \rightarrow \left(\text{Ind}_P^G \chi\right)^\vee \approx \text{Ind}_P^G (\chi^\vee)$$

Thus, V^\vee imbeds into a principal series. From above, this implies that $(V^\vee)_N$ is non-trivial. Thus, by the isomorphism just above, $(V_N)^\vee$ is non-trivial. Thus, V_N must be non-trivial, so V imbeds to some principal series. ///

Thus, failure of irreducibility of I_χ gives rise to G -maps

$$I_\chi \rightarrow I_\beta$$

which are neither injections nor surjections. To study this, we have the following result, due to Mackey in the finite case, and extended by Bruhat to p -adic and Lie groups. For w in the Weyl group $W = \{1, w_o\}$ and for a character χ of M , let

$$\chi^w(m) = \chi(wmw^{-1})$$

The following result uses the finiteness of the group G .

[1.16] Theorem: The complex vectorspace $\text{Hom}_G(I_\chi, I_\beta)$ of G -maps from one principal series to another is

$$\text{Hom}_G(I_\chi, I_\beta) \approx \bigoplus_{w \in W} \text{Hom}_M(\chi^w, \beta)$$

Generally, for two subgroups A and B of a finite group H , and for one-dimensional representations α, β of them, we have a complex-linear isomorphism

$$\text{Hom}_H(\text{Ind}_A^H \alpha, \text{Ind}_B^H \beta) \approx \bigoplus_{w \in A \backslash H / B} \text{Hom}_{w^{-1}Aw \cap B}(\alpha^w, \beta)$$

[1.17] Remark: The decomposition over the double coset $A \backslash H / B$ is an *orbit decomposition* or *Mackey decomposition* or *Mackey-Bruhat decomposition* of the space of H -maps.

Proof: By Frobenius Reciprocity

$$\text{Hom}_H(\text{Ind}_A^H \alpha, \text{Ind}_B^H \beta) \approx \text{Hom}_B(\text{Ind}_A^H \alpha, \beta)$$

As a B -representation space, $\text{Ind}_A^H \alpha$ breaks up into a sum over B -orbits on $A \backslash H$, indexed by $w \in A \backslash H / B$. Via the natural bijection

$$A \backslash AwB \rightarrow (w^{-1}Aw \cap B) \backslash B \quad \text{by} \quad Awb \rightarrow (w^{-1}Aw \cap B)b$$

functions on AwB with the property

$$f(awb) = \alpha(a) f(wb)$$

for $a \in A$ and $b \in B$ become functions on B with

$$f(b_o b) = \alpha(wb_o w^{-1}) f(b)$$

for b_o in $w^{-1}Aw \cap B$. Thus,

$$\text{Hom}_H(\text{Ind}_A^H \alpha, \text{Ind}_B^H \beta) \approx \bigoplus_{w \in A \backslash H / B} \text{Hom}_B(\text{Ind}_{w^{-1}Aw \cap B}^B \alpha^w, \beta)$$

For two B -representations X and Y , there is a natural *dualization isomorphism*

$$\text{Hom}_B(X, Y^\vee) \approx \text{Hom}_B(X \otimes Y, \mathbb{C}) \approx \text{Hom}_B(Y, X^\vee)$$

Thus, since finite-dimensional spaces are reflexive, using formulas from above for duals of induced representations,

$$\begin{aligned} \mathrm{Hom}_H(\mathrm{Ind}_A^H \alpha, \mathrm{Ind}_B^H \beta) &\approx \bigoplus_{w \in A \backslash H/B} \mathrm{Hom}_B(\beta^{-1}, \mathrm{Ind}_{w^{-1}Aw \cap B}^B (\alpha^w)^{-1}) \\ &\approx \bigoplus_{w \in A \backslash H/B} \mathrm{Hom}_{w^{-1}Aw \cap B}(\beta^{-1}, (\alpha^w)^{-1}) \quad (\text{by Frobenius Reciprocity}) \end{aligned}$$

Dualizing once more,

$$\mathrm{Hom}_H(\mathrm{Ind}_A^H \alpha, \mathrm{Ind}_B^H \beta) \approx \bigoplus_{w \in A \backslash H/B} \mathrm{Hom}_{w^{-1}Aw \cap B}(\alpha^w, \beta)$$

as claimed. Since $P \cap w^{-1}Pw$ contains M for every Weyl element w , and since we do not care what fragment of N it may or may not contain since both α and β have been extended trivially to N , this gives the assertion for principal series. ///

[1.18] Corollary: For regular χ the only G -maps of the principal series I_χ to itself are scalars.

Proof: The property that χ be regular is exactly that $\chi^w \neq \chi$. Thus, from the theorem

$$\mathrm{Hom}_G(I_\chi, I_\chi) \approx \mathrm{Hom}_M(\chi, \chi) \approx \mathbb{C}$$

since χ is one-dimensional. ///

The proof of the following corollary is contrary to the spirit of our discussion, as it invokes Complete Reducibility, but it indicates facts which we will also verify by a more generally applicable method.

[1.19] Corollary: For regular χ the principal series I_χ is irreducible and G -isomorphic to I_{χ^w} .

Proof: (Again, this is a bad proof in the sense that it uses Complete Reducibility.) For any subrepresentation W of I_χ there is a subrepresentation U such that $I_\chi = U \oplus W$. The projection to U by $u \oplus w \rightarrow u$ is a G -representation. But by the corollary above for regular χ this map must be a scalar on I_χ so either U or W is 0, proving irreducibility. Then the non-zero intertwining from I_χ to I_{χ^w} cannot avoid being an isomorphism. ///

We'll give another proof of the irreducibility of regular I_χ shortly. But at the moment we cheat in another way to count irreducibles of $G = GL_2(k)$, comparing to the number we've constructed by regular principal series.

Recall that *the number of irreducible complex representations of a finite group is the same as the number of conjugacy classes in the group.*

In $G = GL_2(k)$ with k finite with q elements, by elementary linear algebra (Jordan form) there are conjugacy classes

central	$\begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix}$	$q - 1$	of them	
non-semi-simple	$\begin{pmatrix} x & 1 \\ 0 & x \end{pmatrix}$	$q - 1$	of them	$(x \neq 0)$
non-central split semi-simple	$\begin{pmatrix} x & 0 \\ 0 & y \end{pmatrix}$	$(q - 1)(q - 2)/2$	of them	$(x \neq y)$
anisotropic semi-simple	\dots	$(q^2 - q)/2$	of them	

where the anisotropic elements are conjugacy classes consisting of matrices with eigenvalues lying properly in the unique quadratic extension of k . Conjugation by the longest Weyl element accounts for the division

by 2 in the non-central split semi-simple case. The division by 2 in the non-split semisimple accounts for the Galois action being given by a conjugation within the group.

These conjugacy classes match in an *ad hoc* fashion with specific representations. Match the central conjugacy classes with the one-dimensional representations (composing determinant with characters $k^\times \rightarrow \mathbb{C}^\times$). Match the non-semi-simple classes with the *complements* (cheating here) to the determinant representations inside the irregular principal series, called *special* representations. Match the regular principal series with non-central split semi-simple classes. Thus, numerically, there are bijections

central	\longleftrightarrow	one-dimensional
non-semi-simple	\longleftrightarrow	special
non-central split semi-simple	\longleftrightarrow	regular principal series
anisotropic semi-simple	\longleftrightarrow	supercuspidal (!)

[1.20] **Remark:** Leftovers are assigned to supercuspidal irreducibles *by default*, lacking any immediate alternative for counting them. From the present viewpoint supercuspidals are defined in a negative sense, as things for lacking a construction. The Segal-Shale-Weil representation will give a sharply different construction of supercuspidal representations for SL_2 .

2. Whittaker functionals, Whittaker models

A more extensible approach to studying the irreducibility of regular principal series representations is by distinguishing a suitable one-dimensional subspace of representations and tracking its behavior under G -maps. In fact, it turns out to be better in general to do a slightly subtler thing and distinguish a one-dimensional space of *functionals*, as follows. For a *non-trivial* character (one-dimensional representation) ψ of N , identify its representation space with \mathbb{C} . For a representation V of G an N -map $V \rightarrow \psi$ is a *Whittaker functional*. A *Whittaker model* for V is a (not identically 0) element of $\text{Hom}_G(V, \text{Ind}_N^G \psi)$. When

$$\dim_{\mathbb{C}} \text{Hom}_N(V, \psi) = 1$$

(as in the following result) one speaks of the *uniqueness of Whittaker functionals*, or *uniqueness of Whittaker models*, since Frobenius reciprocity would then give

$$\dim_{\mathbb{C}} \text{Hom}_N(V, \psi) = \dim_{\mathbb{C}} \text{Hom}_G(V, \text{Ind}_N^G \psi)$$

[2.1] **Remark:** Emphasis on Whittaker functionals arose in part from consideration of Fourier expansions of modular forms.

[2.2] **Remark:** For GL_2 the choice of non-trivial ψ does not matter since M acts transitively on non-trivial ψ :

$$\psi\left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a^{-1} & 0 \\ 0 & 1 \end{pmatrix}\right) = \psi\left(\begin{pmatrix} 1 & ax \\ 0 & 1 \end{pmatrix}\right)$$

More precisely:

[2.3] **Proposition:** For GL_2 (not SL_2), with ψ and ψ' two non-trivial characters on N , there is a unique $m \in M/Z$ such that

$$\psi'(n) = \psi(mnm^{-1}) \quad (\text{for all } n \in N)$$

Therefore, there is a G -isomorphism

$$T : \text{Ind}_N^G \psi \approx \text{Ind}_N^G \psi'$$

given by

$$Tf(g) = f(mg)$$

Proof: The first assertion amounts to the fact that every $\psi : \mathbb{F}_q \rightarrow \mathbb{C}^\times$ is of the form

$$\psi(x) = \psi_o(\mathrm{tr}_{\mathbb{F}_q/\mathbb{F}_p} x)$$

where \mathbb{F}_p is the prime field under \mathbb{F}_q and tr is the Galois trace. ^[4] The formula written gives a G -map, because left multiplication by m commutes with right multiplication by g . The map is arranged to convert left equivariance by ψ into left equivariance by ψ' . ///

[2.4] **Remark:** For SL_2 the choice of ψ *does* matter, since the number of orbits of characters on N under the M -action in that case is the cardinality of $k^\times/(k^\times)^2$, which is 2.

[2.5] **Proposition:** For all χ on M ,

$$\dim_{\mathbb{C}} \mathrm{Hom}_N(\mathrm{Ind}_P^G \chi, \psi) = 1$$

Proof: By Frobenius Reciprocity and the (Mackey) orbit decomposition, as earlier,

$$\mathrm{Hom}_N(\mathrm{Ind}_P^G \chi, \psi) \approx \bigoplus_{w \in P \backslash G/N} \mathrm{Hom}_{w^{-1}Nw \cap N} \chi^w, \psi$$

For $w = 1$, since χ is trivial on N , the space of homomorphisms from χ^w to ψ is 0. Thus, there is only one non-zero summand, corresponding to the longest Weyl element $w = w_o$, and this summand gives a one-dimensional space of N -maps. ///

Given the uniqueness, an explicit formula for the Whittaker functional becomes all the more interesting, to allow normalization and comparison.

[2.6] **Proposition:** Let w be the longest Weyl element. For $f \in \mathrm{Ind}_P^G \chi$ the formula

$$\Lambda f = \sum_{n \in N} f(wn) \overline{\psi(n)} \in \mathbb{C}$$

defines a non-zero element Λ of $\mathrm{Hom}_N(\mathrm{Ind}_P^G \chi, \psi)$.

Proof: It is formal, by changing variables in the integral, that the indicated expression is an N -map to ψ . To see that it is not identically 0 it suffices to see that it is non-zero on a well-chosen f . In particular, exploit the finiteness and take f to be 1 at w and 0 otherwise. Then

$$\Lambda f = \sum_{n \in N} f(w\nu) \overline{\psi(\nu)} d\nu = \mathrm{meas}\{1\} \neq 0$$

as desired. ///

[2.7] **Proposition:** For finite-dimensional representations V of N

$$\mathrm{Hom}_N(V, \psi) = 0 \iff \mathrm{Hom}_N(V^\vee, \psi^\vee) = 0$$

[4] This classification of characters on \mathbb{F}_q is substantially a corollary of the larger fact that the *trace pairing on a finite separable extension is non-degenerate*: that is, for a finite separable field extension K/k , the symmetric k -bilinear k -valued form \langle, \rangle on $K \times K$ defined by $\langle x, y \rangle = \mathrm{tr}_{K/k}(xy)$ is non-degenerate, in the sense that for every $x \in K$ there is $y \in K$ such that $\langle x, y \rangle \neq 0$.

Proof: For a non-zero Whittaker functional $\Lambda \in \text{Hom}_N(V, \psi)$, pick x in the second dual $V^{\vee\vee}$ such that $x(\Lambda) \neq 0$. Then

$$\left(\sum_{n \in N} nx \psi(n) \right) (\Lambda) = \sum_{n \in N} (nx)(\Lambda) \psi(n) = \sum_{n \in N} x(n^{-1}\Lambda) \psi(n) = \sum_{n \in N} x(\Lambda) = x(\Lambda) \cdot \text{meas}(N)$$

which shows that $\sum_{n \in N} nx \psi(n)$ is not 0. For $\nu \in N$

$$\sum_{n \in N} n\nu \cdot x \psi(n) = \psi^\vee(\nu) \sum_{n \in N} n \cdot x \psi(n)$$

by replacing n by $n\nu^{-1}$. ///

We can use the Whittaker functionals Λ to redo our study of intertwinings $T : I_\chi \rightarrow I_{\chi^w}$ among principal series.

[2.8] **Proposition:** A finite-dimensional representation V of G with

$$\text{Hom}_N(V, \psi) = \mathbb{C}$$

and with

$$\text{Hom}_N(V^\vee, \psi^\vee) = \mathbb{C}$$

is irreducible if and only if the Whittaker functional in $\text{Hom}_N(V, \psi)$ generates the dual V^\vee and the Whittaker functional in $\text{Hom}_N(I_\chi^\vee, \psi^\vee)$ generates the second dual $V^{\vee\vee} \approx V$.

Proof: On one hand, if V is irreducible, then (from above) the dual is irreducible, so certainly is generated (under G) by the (non-zero) Whittaker functional. The same applies to the second dual. This is the easy part of the argument. On the other hand, suppose that the Whittaker functionals generate (under G) the dual and second dual. A proper subrepresentation Λ of V^\vee cannot contain the Whittaker functional, since the Whittaker functional generates the whole representation. Thus, the image of the Whittaker functional in the quotient $Q = V^\vee/\Lambda$ is not 0. From just above, since Q contains a non-zero Whittaker functional so must Q^\vee (for the character ψ^\vee). But then the natural inclusion

$$Q^\vee \subset V^{\vee\vee}$$

shows that the Whittaker vector generates a proper subrepresentation of $V^{\vee\vee}$, contradiction. ///

[2.9] **Proposition:** The dual I_χ^\vee of a principal series I_χ fails to be generated by a Whittaker functional Λ if and only if there is a non-zero intertwining $T : I_\chi^\vee \rightarrow I_{\chi^w}^\vee$ in which $T\Lambda = 0$, for some w in the Weyl group W .

Proof: If Λ generates a proper subrepresentation V of I_χ^\vee , then there is an irreducible non-zero quotient Q of I_χ^\vee/V . From above, Q again imbeds into some principal series I_ω . This yields a non-zero intertwining $I_\chi^\vee \rightarrow I_\omega$ in which the Whittaker functional is mapped to 0. ///

Recall that the only principal series representation admitting a non-zero intertwining from I_χ is I_{χ^w} , for w in the Weyl group. For a character $\omega : k^\times \rightarrow \mathbb{C}^\times$, define a Gauss sum

$$g(\omega, \psi) = \sum_{x \in k^\times} \omega(x) \bar{\psi} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$$

The normalized Whittaker functional in I_χ^\vee is

$$\Lambda_\chi f = \sum_{n \in N} f(won) \psi(n)$$

[2.10] **Proposition:** Let $w = w_o$ be the longest Weyl element. Under the intertwining

$$T : I_\chi \rightarrow I_{\chi^w} \quad \text{by} \quad Tv(g) = \sum_{n \in N} v(wng)$$

the normalized Whittaker functional Λ_{χ^w} in $(I_{\chi^w})^\vee$ is mapped by the adjoint T^\vee to

$$T^\vee(\Lambda_{\chi^w}) = g(\chi, \psi) \cdot \Lambda_\chi \in (I_\chi)^\vee$$

Proof: Using the uniqueness of the Whittaker functionals on principal series, it suffices to compute the values of the images on a well-chosen function f .

$$\Lambda^w T f = \sum_{n \in N} T f(wn) \overline{\psi(n)} = \sum_{n \in N} \sum_{n \in N} f(w\nu wn) \overline{\psi(n)} d\nu$$

To compare this to

$$\Lambda f = \sum_{n \in N} f(wn) \overline{\psi(n)}$$

take f to be

$$f(nmw) = \chi(m)$$

for $n \in N$ and $m \in M$, and 0 otherwise. That is, f is supported on Pw and is 1 at w . Then $wn \in Pw$ if and only if $n = 1$. Thus,

$$\Lambda f = \overline{\psi(1)} = 1$$

On the other hand, the condition $w\nu wn \in Pw$ is met in a more complicated manner. Indeed, letting $\nu = \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix}$ we have the awkward-but-important standard identity for $x \neq 0$:

$$w\nu w = \begin{pmatrix} -1 & 0 \\ x & -1 \end{pmatrix} = \begin{pmatrix} 1 & -1/x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1/x & 0 \\ 0 & x \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & -1/x \\ 0 & 1 \end{pmatrix}$$

Note that this identity works in both GL_2 and in SL_2 . Thus,

$$w\nu wn \in Pw \iff n = \begin{pmatrix} 1 & 1/x \\ 0 & 1 \end{pmatrix}$$

and in that case

$$w\nu wn = \begin{pmatrix} 1 & -1/x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1/x & 0 \\ 0 & x \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

Thus, for $G = GL_2$

$$\Lambda_{\chi^w} T f = \sum_{x \in k^\times} \frac{\chi_2}{\chi_1}(x) \psi^\vee \begin{pmatrix} 1 & 1/x \\ 0 & 1 \end{pmatrix}$$

Replacing x by $1/x$, we conclude that with the intertwining

$$Tv(g) = \sum_{n \in N} v(wng)$$

the normalized Whittaker functional Λ_{χ^w} is mapped to $g(\chi, \psi)$ times the Whittaker functional Λ_χ under the adjoint T^\vee . For $G = SL_2$ the conclusion is nearly identical, with χ_2 replaced by χ_1^{-1} , in effect. ///

[2.11] **Corollary:** For regular χ the corresponding Gauss sum is non-zero, hence the Whittaker functional is never annihilated by a non-zero intertwining, hence the Whittaker functional generates I_χ^\vee . Likewise the corresponding Whittaker functional generates $I_\chi^{\vee\vee}$. Thus, I_χ is irreducible.

Proof: We recall a computation that proves the Gauss sum is non-zero for $\chi_1 \neq \chi_2$.

$$|g(\chi, \psi)|^2 = \sum_{x \in k^\times} \sum_{y \in k^\times} \frac{\chi_1(x/y)}{\chi_2} \bar{\psi} \begin{pmatrix} 1 & x-y \\ 0 & 1 \end{pmatrix} = \sum_{x \in k^\times} \sum_{y \in k^\times} \frac{\chi_1(x)}{\chi_2} \bar{\psi} \begin{pmatrix} 1 & y(x-1) \\ 0 & 1 \end{pmatrix}$$

replacing x by xy . For fixed $x \neq 1$, the sum over y would be over k^\times if it were not missing the $y = 0$ term, so by the *cancellation lemma* (orthogonality of characters) it is

$$\sum_{y \in k} \bar{\psi} \begin{pmatrix} 1 & y(x-1) \\ 0 & 1 \end{pmatrix} - 1 = -1$$

For $x = 1$, the sum is $q - 1$, where $|k| = q$. Thus,

$$|g(\chi, \psi)|^2 = q - \sum_{x \in k^\times} \frac{\chi_1}{\chi_2}(x) = q - 0 \quad (\text{for } \chi_1 \neq \chi_2)$$

Thus, the Gauss sum is non-zero. Thus, the adjoint T^\vee of the intertwining $T : I_\chi \rightarrow I_{\chi^w}$ just above does not annihilate the Whittaker functional. Since χ is regular, χ^w is regular, and every non-zero intertwining of I_χ to itself is a non-zero multiple of the identity, so again the Whittaker functional is not annihilated by the adjoint T^\vee .

Thus, the Whittaker functional generates the dual I_χ^\vee . Similarly, the corresponding Whittaker functional generates the second dual, and from above we conclude that I_χ is irreducible. ///

[2.12] **Remark:** The previous discussion is a simple example illustrating the spirit of Casselman's 1980 use of *spherical* vectors to examine irreducibility of unramified principal series of p -adic reductive groups.

[2.13] **Remark:** For irregular χ we could invoke complete reducibility and the computation (above) that for irregular χ

$$\dim_{\mathbb{C}} \text{Hom}_G(I_\chi, I_\chi) = \text{card } P \backslash G/P = 2$$

to see that I_χ is a direct sum of two irreducibles. Further, we can immediately identify the one-dimensional subrepresentation $\chi_1 \circ \det$ of I_χ for irregular $\chi = (\chi_1, \chi_1)$ for GL_2 . It is immediate that $\chi_1 \circ \det$ has no Whittaker functional, so we can anticipate that (still using complete reducibility) the *other* irreducible in irregular I_χ has a Whittaker functional. This other irreducible is a *special* representation.

3. Uniqueness of Whittaker functionals/models

So far we have no tangible description for the supercuspidal irreducibles V , except that $V_N = 0$. In particular, we cannot address uniqueness of Whittaker functionals for supercuspidals by explicit computation since we have no tangible models, but their Whittaker models exist simply because the Jacquet modules are trivial (see just below). Note that, for a representation V of G ,

$$\text{Hom}_N(V, \psi) \approx \text{Hom}_G(V, \text{Ind}_N^G \psi) \quad (\text{by Frobenius Reciprocity})$$

That is, Whittaker functionals correspond to G -intertwinings to the *Whittaker space* $\text{Ind}_N^G \psi$.

[3.1] **Proposition:** A supercuspidal irreducible V of GL_2 has a Whittaker model.

Proof: As a representation of N (by restriction), V is a sum of irreducibles. Since V is supercuspidal its Jacquet module is trivial, so the trivial representation of N does not occur. Thus, a non-trivial representation ψ of N *does* occur. Since N is abelian, this irreducible is one-dimensional. Since V is stable under the action

of M , and (as observed earlier) M is transitive on non-trivial characters on N , every non-trivial ψ of N occurs in V . ///

[3.2] **Remark:** The analogous result about Whittaker models for supercuspidal representations is more complicated for SL_2 , since in that case M has *two* orbits on non-trivial characters of N .

[3.3] **Theorem:** Let ψ be a non-trivial character on N . The endomorphism algebra

$$\mathrm{Hom}_G(\mathrm{Ind}_N^G \psi, \mathrm{Ind}_N^G \psi)$$

is *commutative*. Thus, we have *Uniqueness of Whittaker functionals*: For an irreducible representation V of G

$$\dim_{\mathbb{C}} \mathrm{Hom}_N(V, \psi) \leq 1$$

Equivalently, we have *Uniqueness of Whittaker models*

$$\dim_{\mathbb{C}} \mathrm{Hom}_G(V, \mathrm{Ind}_N^G \psi) \leq 1$$

[3.4] **Remark:** For GL_2 , the only case where the dimension of intertwinings is 0 rather than 1 is for the one-dimensional representations, that is, for composition of determinant with characters of k^\times .

Proof: First, we see how commutativity of the endomorphism ring implies that multiplicities are ≤ 1 . Use complete reducibility, so

$$\mathrm{Ind}_N^G \psi \approx \bigoplus_V m_V \cdot V$$

where V runs through isomorphism classes of irreducibles and m_V is the multiplicity of V . Then

$$\mathrm{End}_G(\mathrm{Ind}_N^G \psi) \approx \prod_V M_{m_V}(\mathbb{C})$$

where $M_n(\mathbb{C})$ is the ring of n -by- n matrices with complex entries. Thus, this endomorphism ring is commutative if and only if all the multiplicities are 1.

To study the endomorphism ring, use the Mackey-Bruhat orbit decomposition of the space of intertwinings from one induced representation to another in the case that the two induced representations are the same. Thus, given

$$T \in \mathrm{Hom}_G(\mathrm{Ind}_A^G \alpha, \mathrm{Ind}_B^G \beta)$$

let K_T be a *kernel function*^[5] on $G \times G$ such that

$$Tf(g) = \sum_{h \in G} K_T(g, h) f(h)$$

The fact that T is a G -map gives, for all $x \in G$,

$$\begin{aligned} \sum_{h \in G} K_T(gx, h) f(h) &= Tf(gx) = (R_x T f)(g) \\ &= (TR_x f)(g) = \sum_{h \in G} K_T(g, h) f(hx) = \sum_{h \in G} K_T(g, hx^{-1}) f(h) \end{aligned}$$

[5] This use of *kernel* is incompatible with the use where x is in the kernel of a homomorphism f when $f(x) = 0$.

by replacing h by hx^{-1} , where R_x is the right regular representation. Thus, the kernel K_T arises from a function of a single variable, and the intertwining T can be rewritten as

$$Tf(g) = \sum_{h \in G} K_T(gh^{-1})$$

Since T maps to $\text{Ind}_B^G \beta$, and maps from $\text{Ind}_A^G \alpha$, it must be that

$$K_T(bxa) = \beta(b) \cdot K_T(x) \cdot \alpha(a)$$

for all $b \in B$, $g \in G$, $a \in A$. A direct computation shows that

$$K_{S \circ T} = K_S * K_T$$

for $S, T \in \text{Hom}_G(\text{Ind}_A^G \alpha, \text{Ind}_B^G \beta)$ with usual convolution

$$(f * \varphi)(g) = \sum_{x \in G} f(gx^{-1}) \varphi(x)$$

Thus, to prove commutativity of the endomorphism ring it is necessary and sufficient to prove commutativity of the convolution ring R of complex-valued functions u on G with the equivariance properties

$$u(bxa) = \beta(b) \cdot u(x) \cdot \alpha(a)$$

for $a \in A$, $b \in B$, $x \in G$.

Note that, for $A = B$ and $\alpha = \beta$, the convolution of two such functions falls back into the same class.

Following Gelfand-Graev and others, to prove commutativity of such a convolution ring, it suffices to find an involutive anti-automorphism σ of G such that for u on G with the property

$$u(bxa) = \psi(b) \cdot u(x) \cdot \psi(a)$$

we have

$$u(g^\sigma) = u(g) \quad (\text{for all } g \in G)$$

To verify that this criterion for commutativity really works, use notation

$$u^\sigma(g) = u(g^\sigma)$$

and let u, v be two such functions. Then

$$(u^\sigma * v^\sigma)(x) = \sum_{g \in G} u((xg^{-1})^\sigma) v(g^\sigma) = \sum_{g \in G} u(gx^\sigma) v(g^{-1})$$

by replacing g by $(g^\sigma)^{-1}$. Replacing g by $g(x^\sigma)^{-1}$ turns this into

$$\sum_{g \in G} u(g) v(x^\sigma g^{-1}) = (v * u)^\sigma(x)$$

That is,

$$(u * v)^\sigma = v^\sigma * u^\sigma$$

Therefore, if $u = u^\sigma$ and $v = v^\sigma$ then

$$u * v = (u * v)^\sigma = v^\sigma * u^\sigma = v * u$$

and the convolution ring is commutative.

To apply the Gelfand-Graev involution idea, we need to classify functions u such that, as above,

$$u(bxa) = \psi(b) \cdot u(x) \cdot \psi(a)$$

since these are the ones that could occur as Mackey-Bruhat kernels. We use the *Bruhat decomposition*, namely that

$$G = NM \cup NMwN \quad (\text{disjoint union, with } w = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix})$$

where M is diagonal matrices and w is a slightly different normalization of longest Weyl element. The group M normalizes N , but does not preserve ψ , since for $m \in M$

$$\psi(mnm^{-1}) \neq \psi(n)$$

for all $n \in N$ unless m is actually in the center Z of G , the scalar matrices. The two-sided equivariance condition entails

$$\psi(n)u(m) = u(nm) = u(m \cdot m^{-1}nm) = u(m)\psi(m^{-1}nm)$$

This does not hold for all $n \in N$ unless m is central. Thus, the left and right N, ψ -equivariant functions supported on NM are those whose support is NZ . For the equivariant functions supported on $NMwN$, there is no such issue, since $N \cap wNw^{-1} = \{1\}$. Thus, the $N \times N$ orbits which can support such equivariant functions are those with representatives $z \in Z$ and mw with $m \in M$. All such functions are linear combinations of functions

$$f(nz) = \psi(n) \quad (\text{for } n \in N, 0 \text{ otherwise})$$

for fixed $z \in Z$ and, for fixed $m \in M$,

$$f(nmw\nu) = \psi(n)\psi(\nu) \quad (\text{for } n, \nu \in N, 0 \text{ otherwise})$$

In this situation, with w the long Weyl element normalized as above, take involutive anti-automorphism

$$g^\sigma = wg^\top w^{-1}$$

This is the identity on the center Z , on N , and on elements mw with $m \in M$. Thus, it is the identity on all such equivariant functions. Thus, this convolution ring meets the Gelfand-Graev criterion for commutativity.

///

[3.5] **Remark:** The kernels K_T introduced in the proof have analogues in more complicated settings, and would more generally be called *Mackey-Bruhat distributions*. That is, the relevant kernel would not in general be given by a *function*, but by a Schwartz *distribution*.

[3.6] **Remark:** Without complete reducibility, the principle that commutativity of an endomorphism ring implies that multiplicities are all ≤ 1 acquires a more complicated form. One version is the *Gelfand-Kazhdan criterion*. The same approach, namely Mackey-Bruhat and Gelfand-Graev, yield further facts whose analogues are more complicated over non-finite fields.

[3.7] **Remark:** For *irregular* χ , we have already seen that I_χ decomposes as the direct sum of two non-isomorphic irreducibles. Thus, for given ψ , one of these subrepresentations has a Whittaker model and one does not. For GL_2 , the irregular principal series always have one-dimensional subrepresentation, which fails to have a Whittaker model. For SL_2 , it is less clear.

4. Summary for GL_2

There are $(q-1)(q-2)/2$ isomorphism classes of irreducible principal series (with $I_\chi \approx I_{\chi^w}$), namely the *regular* ones (i.e., with $\chi^w \neq \chi$). These all have Whittaker models. Their Jacquet modules are 2-dimensional. They are of dimension $|P \backslash G| = q+1$.

There are $q-1$ one-dimensional representations, obtained by composing characters with determinant. Their Jacquet modules are 1-dimensional, not surprisingly. These do not have Whittaker models.

There are $q-1$ *special* representations, subrepresentations of irregular principal series, with 1-dimensional Jacquet modules. They have Whittaker models (since every unramified principal series has a Whittaker functional and one-dimensional representations do not.) Special representations are of dimension q .

There are $q(q-1)/2$ supercuspidal irreducibles, by definition having 0-dimensional Jacquet module, all having a Whittaker model. Each has dimension $q-1$.

[4.1] Remark: Since M is transitive on non-trivial characters on N , there is (up to G -isomorphism) only *one* Whittaker space. This is not true for SL_2 .

[4.2] Remark: One numerical check for the above categorization is the fact (from decomposition of the biregular representation) that the sum of the squares of the dimensions of the irreducibles is the order of the group. Thus, we should have (in the same order that we reviewed them)

$$\begin{aligned} (q^2-1)(q^2-q) &= (\text{cardinality of } GL_2 \text{ over field with } q \text{ elements}) \\ &= (\text{irreducible principal series}) + (\text{one-dimensional}) + (\text{special}) + (\text{supercuspidal}) \\ &= \frac{(q-1)(q-2)}{2} \cdot (q+1)^2 + (q-1) \cdot 1^2 + (q-1) \cdot q^2 + \frac{q(q-1)}{2} \cdot (q-1)^2 \end{aligned}$$

Remove a factor of $q-1$ from both sides, leaving a supposed equality

$$(q^2-1)q = \frac{(q-2)}{2} \cdot (q+1)^2 + 1 + q^2 + \frac{q(q-1)}{2} \cdot (q-1)$$

Anticipating a factor of q throughout, combine the first two summands on the right-hand side to obtain (multiplying everything through by 2, as well)

$$2(q^2-1)q = (q^3-3q) + 2q^2 + q(q-1)^2$$

which allows removal of the common factor of q , to have the supposed equality

$$2(q^2-1) = q^2-3+2q+(q-1)^2$$

The degree is low enough to multiply out, giving an alleged equality

$$2q^2-2 = q^2-3+2q+q^2-2q+1$$

which is easy to verify. The reduction steps were reversible, so this counting check succeeds.

[4.3] Remark: Another numerical check would be by counting the irreducibles with Whittaker models, versus the dimension of the space of endomorphisms of the Whittaker space, since the latter is commutative (above). The number of irreducibles with Whittaker models is

$$\begin{aligned} \text{irreds with Whittaker models} &= (\text{irreducible principal series}) + (\text{special}) + (\text{supercuspidal}) \\ &= \frac{(q-1)(q-2)}{2} + (q-1) + \frac{q(q-1)}{2} = (q-1) \left[\frac{q-2}{2} + 1 + \frac{q}{2} \right] = q(q-1) \end{aligned}$$

On the other hand, the dimension of the space of endomorphisms of the Whittaker space (from the proof of commutativity of the endomorphism ring, above) is the cardinality

$$\begin{aligned} &(\text{number of left-and-right } N \times N \text{ orbits supporting left-and-right } \psi\text{-equivariant functions}) \\ &= \text{card}(N \backslash NZ / N \sqcup N \backslash Pw_oP / N) = \text{card}(Z) + \text{card}(M) = (q-1) + (q-1)^2 = q(q-1) \end{aligned}$$

where Z is the center of GL_2 . They match.

5. Conjugacy classes in SL_2 , odd q

Before pairing up conjugacy classes and irreducibles for SL_2 over a finite field with q elements, we must take greater pains to identify conjugacy classes. For SL_2 the parity of q matters, while it did not arise for GL_2 . In $G = SL_2(k)$ with k finite with q elements, the collection of conjugacy classes is more complicated than the pure linear algebra of $GL_2(k)$. The non-semisimple elements' conjugacy classes are most disturbed by the change from GL_2 to SL_2 . Let ϖ be a non-square in k^\times , and take q odd.

central	$\begin{pmatrix} x & 0 \\ 0 & x \end{pmatrix}$	2	of them	$(x = \pm 1)$
non-semisimple	$\begin{pmatrix} x & 1 \\ 0 & x \end{pmatrix}$	2	of them	$(x = \pm 1)$
non-semisimple	$\begin{pmatrix} x & \varpi \\ 0 & x \end{pmatrix}$	2	of them	$(x = \pm 1)$
non-central split semi-simple	$\begin{pmatrix} x & 0 \\ 0 & x^{-1} \end{pmatrix}$	$(q-3)/2$	of them	$(x \neq \pm 1)$
non-split semi-simple	...	$(q-1)/2$	of them	

where the anisotropic elements are conjugacy classes of matrices with eigenvalues lying properly in the (unique) quadratic extension of k , and with Galois norm 1. The division by 2 in the latter is because the Galois action is given by a conjugation in the group. In the case of split semi-simple elements the division by 2 reflects the fact that conjugation interchanges a and a^{-1} . Verification that these are exactly the SL_2 conjugacy classes is at least mildly interesting, and we carry out this exercise to have specifics used later.

Sketch the discussion for odd q . First, observe that if $g \in G$ has elements in its centralizer $C(g)$ in GL_2 having determinants running through all of k^\times , then

$$\{xgx^{-1} : x \in SL_2\} = SL_2 \cap \{xgx^{-1} : x \in GL_2\}$$

That is, with the hypothesis on the centralizer, the intersection with SL_2 of a GL_2 conjugacy class does not break into proper subsets under SL_2 conjugation. For g central or of the form $g = \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$ the element $\begin{pmatrix} d & 0 \\ 0 & 1 \end{pmatrix}$ in the centralizer has determinant $d \in k^\times$, meeting this hypothesis. Now consider non-split semi-simple elements g . It is elementary that such g lies in an imbedded copy of the norm-one elements K^1 in the unique quadratic extension K of k . The group K^\times imbeds compatibly in GL_2 , and determinant on the imbedded copy is the Galois norm. Since norm is surjective on finite fields, non-split semi-simple conjugacy classes also meet the hypothesis above, so there is no change from GL_2 to SL_2 .

The non-semi-simple classes are subtler. First, non-semisimple elements u must have rational eigenvalues, and the non-semi-simplicity then implies that such u stabilizes a unique line λ in k^2 . By the transitivity of

SL_2 on lines, all non-semi-simple conjugacy classes in SL_2 have representatives of the form $u = \begin{pmatrix} a & * \\ 0 & a^{-1} \end{pmatrix}$ with non-zero upper left entry with $a = \pm 1$, stabilizing the obvious line λ . If another such matrix $v = \begin{pmatrix} b & * \\ 0 & b^{-1} \end{pmatrix}$ with non-zero upper right entry is conjugate to u , say $x^{-1}vx = u$, then $vx = xu$ and

$$vx \cdot \lambda = xu \cdot \lambda$$

from which

$$vx \cdot \lambda = x \cdot \lambda$$

since u fixes λ . This implies that v fixes $x\lambda$, so $x\lambda = \lambda$ (since v fixes exactly one line), and necessarily x is of the form

$$x = \begin{pmatrix} b & * \\ 0 & b^{-1} \end{pmatrix}$$

for some $b \in k^\times$. By this point, the remaining computations are not hard. Specifically, conjugation by upper-triangular matrices in SL_2 acting on matrices $\begin{pmatrix} a & * \\ 0 & a^{-1} \end{pmatrix}$ adjusts the upper-right entry only by squares in k^\times . Since k^\times is cyclic, there are exactly two orbits. Thus, as asserted above, the non-semi-simple conjugacy classes have representatives

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix} \quad \begin{pmatrix} 1 & \varpi \\ 0 & 1 \end{pmatrix} \quad \begin{pmatrix} -1 & \varpi \\ 0 & -1 \end{pmatrix}$$

where ϖ is a non-square in k^\times .

6. Irreducibles of SL_2 , q odd

Now we classify irreducibles of $G = SL_2$ over a finite field with an *odd* number of elements q . Unlike the case of GL_2 , for SL_2 there are *two* inequivalent families of Whittaker models, as there are two characters ψ and ψ' on N , *not* related to each other by conjugation by M , unlike GL_2 . Fix two such SL_2 -unrelated ψ and ψ' , and refer to the ψ -Whittaker and ψ' -Whittaker models or functionals.

First, parallel to the discussion of principal series for GL_2 , the principal series

$$I_\chi = \text{Ind}_P^G \chi$$

for the $q-3$ regular χ 's on M are *irreducible*, and there is an isomorphism

$$I_\chi \rightarrow I_{\chi^{-1}}$$

so there are $(q-3)/2$ irreducibles occurring as principal series. There are exactly two *irregular* characters here, the trivial character and the (unique) other character that assumes values ± 1 . Let the corresponding principal series be denoted I_1 and I_{-1} . Just as for GL_2

$$I_1 = \mathbb{C} \oplus \text{special}$$

where \mathbb{C} is the trivial representation. The same techniques show that

$$I_{-1} = \text{direct sum of two irreducibles}$$

but neither of the two irreducibles is one-dimensional. Both of these have one-dimensional Jacquet modules, since they both imbed into a principal series.

[6.1] **Remark:** For $q \geq 3$ the derived group of $G = SL_2(\mathbb{F}_q)$ is G itself, so there are no non-trivial one-dimensional representations of G .

It remains true for SL_2 that for either Whittaker model, ψ or ψ' , there is a unique Whittaker functional on a (regular or not) principal series I_χ . The trivial representation has no Whittaker model of *either* type, so the *special* representation has a Whittaker model of *both* types. Irreducible principal series have Whittaker models of *both* types.

The nature of the Whittaker models (or lack thereof) is not clear yet for the irreducibles into which the irregular I_{-1} decomposes.

[6.2] **Proposition:** A supercuspidal irreducible for SL_2 has *either* a ψ -Whittaker model or a ψ' -Whittaker model.

Proof: A supercuspidal, which by definition has a trivial Jacquet module, must have a non-trivial ψ -isotype for N for *some* ψ . As observed in the discussion of the Whittaker spaces for GL_2 , conjugation by M gives G -isomorphic Whittaker spaces. Thus, if ψ and ψ' are representatives for the two M -orbits, a supercuspidal must have one or the other Whittaker model. ///

[6.3] **Proposition:** The number of irreducibles of SL_2 with ψ -Whittaker models is $q + 1$. The number of irreducibles with ψ' -Whittaker models is $q + 1$. The number of irreducibles which have *both* types of Whittaker models is $q - 1$.

Proof: The argument used in the GL_2 -case, following Mackey-Bruhat and Gelfand-Graev, succeeds here. The support of a left and right ψ -equivariant distribution on SL_2 must have support on

$$NZ \sqcup NMw_oN$$

and (keeping in mind that q is odd) the dimension of the space of all such is the cardinality

$$\text{card } N \backslash (NZ \sqcup NMw_oN) / N = 2 + (q - 1) = q + 1$$

The same conclusion works for any non-trivial character. If, instead, we require left ψ' -equivariance and right ψ -equivariance with M -inequivalent characters, we claim that only the larger Bruhat cell can support appropriate distributions, so the dimension is $q - 1$. Indeed, this is exactly the assumption that ψ and ψ' are not conjugated to each other by any element of the Levi component M in SL_2 . ///

Thus, the two non-isomorphic types of Whittaker models have exactly $q - 1$ isomorphism classes in common out of $q + 1$ in each. The $(q - 3)/2$ irreducible principal series account for some of these common ones. The *special* representation (in I_1) is another that lies in both, since the trivial representation lies in *neither*, and I_1 has a unique Whittaker vector (for either character).

[6.4] **Lemma:** One of the two irreducible summands of I_{-1} lies in one Whittaker space and the other lies in the other Whittaker space.

Proof: When an irreducible V has non-trivial

$$\psi : \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \rightarrow \psi_o(x)$$

isotype for N , under the action of M it also has a non-trivial

$$\psi_a : \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \rightarrow \psi_o(a^2x)$$

isotype for N . There are $(q - 1)/2$ characters in such an M -orbit. Thus, if ψ and ψ' are M -inequivalent and V has *both* ψ -Whittaker and ψ' -Whittaker models, V has a non-trivial isotype for *all* of the $q - 1$ non-trivial

characters on N . As remarked above, the two summands in I_{-1} both have one-dimensional Jacquet modules (trivial N -isotypes), and are not one-dimensional. Thus, the dimension of each summand in I_{-1} is at least

$$1 + (q - 1)/2 = (q + 1)/2$$

The dimension of the whole I_{-1} is $q + 1$, so it must be that each has dimension *exactly* $(q + 1)/2$. Thus, indeed, one has one type of Whittaker model, and the other has the other type. ///

So far, each Whittaker space has the unique special representation (from I_1), $(q - 3)/2$ irreducible principal series, and 1 from among the two summands of I_{-1} . Each supercuspidal irreducible has at least one Whittaker model from among the two. Only the trivial (one-dimensional) representation has *no* Whittaker model of either type.

The previous proposition shows that there are 4 irreducibles with *exactly one* Whittaker model, and that two of these have a ψ -model and two have a ψ' -model. The two irreducible summands of I_{-1} account for two of these. The remaining irreducibles are (by definition) supercuspidal. Thus, there are exactly 2 supercuspidal irreducibles of SL_2 having a *single type* of Whittaker model.

We can do a numerical check. Again, the number of conjugacy classes in SL_2 over a field with an odd number q of elements is

$$\begin{aligned} & (\text{central}) + (\text{non-semi-simple}) + (\text{new non-semi-simple}) \\ & + (\text{non-central split semisimple}) + (\text{non-split semisimple}) \\ & = 2 + 2 + 2 + \frac{(q - 3)}{2} + \frac{(q - 1)}{2} = q + 4 \end{aligned}$$

Thus, excluding the trivial representation, there are $q + 3$ irreducibles *with* at least one type of Whittaker model. There are $q - 1$ irreducibles in common between the two types of Whittaker models, and each model has dimension $q + 1$, so the total indeed is

$$2 \cdot (q + 1) - (q - 1) = q + 3$$

[6.5] Remark: Remarks just above also show that the supercuspidal irreducibles with *both* types of Whittaker models are of dimension $q - 1$ (the number of all non-trivial characters of N), while the 2 supercuspidal irreducibles with only *one* type of model are of dimension $(q - 1)/2$, distinguishing these two smaller supercuspidals among supercuspidals.
