

(February 15, 2019)

08d. Distributions supported at 0

Paul Garrett garrett@math.umn.edu <http://www.math.umn.edu/~garrett/>

[This document is

http://www.math.umn.edu/~garrett/m/real/notes_2018-19/08d_distns_at_zero.pdf]

[0.1] Theorem: A distribution u with *support* $\{0\}$ is a (finite) linear combination of Dirac's δ and its derivatives.

Recall the notion of *support* of a distribution: the *support* of a distribution u is the *complement* of the *union* of all open sets $U \in \mathbb{R}^n$ such that

$$u(f) = 0 \quad (\text{for } f \in \Delta_K \text{ with compact } K \subset U)$$

Proof: The space Δ of test functions on \mathbb{R}^n is $\Delta = \bigcup_K \Delta_K$, where Δ_K is test functions supported on compact K . The latter is a Fréchet space, with *norms*

$$\nu_{k,K}(f) = \sup_{i \leq k, x \in K} |f^{(i)}(x)|$$

Thus, it suffices to classify u in Δ_K^* with support $\{0\}$.

We have seen that a continuous linear map T from a *limit* of Banach spaces (such as Δ_K) to \mathbb{C} factors through a limitand. Thus, there is an *order* $k \geq 0$ such that u factors through

$$C_K^k = \{f \in C^k(K) : f^{(\alpha)} \text{ vanishes on } \partial K \text{ for all } \alpha \text{ with } |\alpha| \leq k\}$$

We need an auxiliary gadget. Fix a smooth compactly-supported function ψ identically 1 on a neighborhood of 0, bounded between 0 and 1, and (necessarily) identically 0 outside some (larger) neighborhood of 0. For $\varepsilon > 0$ let

$$\psi_\varepsilon(x) = \psi(\varepsilon^{-1}x)$$

Since the support of u is just $\{0\}$, for all $\varepsilon > 0$ and for all $f \in \mathcal{D}(\mathbb{R}^n)$ the support of $f - \psi_\varepsilon \cdot f$ does not include 0, so

$$u(\psi_\varepsilon \cdot f) = u(f)$$

Thus, for some constant C (depending on k and K , but not on f)

$$|\psi_\varepsilon f|_k = \sup_{x \in K} \sup_{|\alpha| \leq k} |(\psi_\varepsilon f)^{(\alpha)}(x)| \leq C \cdot \sup_{|\alpha| \leq k} \sup_x \sup_{0 \leq j \leq i} \varepsilon^{-|\alpha|} \left| \psi^{(j)}(\varepsilon^{-1}x) f^{(i-j)}(x) \right|$$

For f vanishing to order k at 0, that is, $f^{(\alpha)}(0) = 0$ for all multi-indices α with $|\alpha| \leq k$, on a fixed neighborhood of 0, by a Taylor-Maclaurin expansion, for some constant C

$$|f(x)| \leq C \cdot |x|^{k+1}$$

and, generally, for α^{th} derivatives with $|\alpha| \leq k$,

$$|f^{(\alpha)}(x)| \leq C \cdot |x|^{k+1-|\alpha|}$$

For some constant C

$$|\psi_\varepsilon f|_k \leq C \cdot \sup_{|\alpha| \leq k} \sup_{0 \leq j \leq i} \varepsilon^{-|\alpha|} \cdot \varepsilon^{k+1-|\alpha|+|\alpha|} \leq C \cdot \varepsilon^{k+1-|\alpha|} \leq C \cdot \varepsilon^{k+1-k} = C \cdot \varepsilon$$

Thus, for all $\varepsilon > 0$, for smooth f vanishing to order k at 0,

$$|u(f)| = |u(\psi_\varepsilon f)| \leq C \cdot \varepsilon$$

Thus, $u(f) = 0$ for such f .

That is, u is 0 on the intersection of the kernels of δ and its derivatives $\delta^{(\alpha)}$ for $|\alpha| \leq k$. Generally,

[0.2] Proposition: A continuous linear function $\lambda \in V^*$ vanishing on the intersection of the kernels of a finite collection $\lambda_1, \dots, \lambda_n$ of continuous linear functionals on V is a linear combination of the λ_i .

Proof: The linear map

$$q : V \longrightarrow \mathbb{C}^n \quad \text{by} \quad v \longrightarrow (\lambda_1 v, \dots, \lambda_n v)$$

is *continuous* since each λ_i is continuous, and λ factors through q , as $\lambda = L \circ q$ for some linear functional L on \mathbb{C}^n . We know all the linear functionals on \mathbb{C}^n , namely, L is of the form

$$L(z_1, \dots, z_n) = c_1 z_1 + \dots + c_n z_n \quad (\text{for some constants } c_i)$$

Thus,

$$\lambda(v) = (L \circ q)(v) = L(\lambda_1 v, \dots, \lambda_n v) = c_1 \lambda_1(v) + \dots + c_n \lambda_n(v)$$

expressing λ as a linear combination of the λ_i . ///

[0.3] Remark: The following lemma resolves a potential confusion.

[0.4] Lemma: For compact K inside the *complement* of the support of a distribution u ,

$$u(f) = 0 \quad (\text{for } f \in \Delta_K)$$

Proof: This is plausible, but not utterly trivial. Let $\{U_i : i \in I\}$ be open sets such that for compact K' inside any single U_i and $f \in \Delta_{K'}$ we have $u(f) = 0$. Let $\{\psi_i : i \in I\}$ be a smooth locally finite *partition of unity*^[1] subordinate to $\{U_i : i \in I\}$. Take $f \in \Delta_{K'}$ for K' compact inside $U = \bigcup_i U_i$. Then

$$f = f \cdot 1 = \sum_i f \cdot \psi_i$$

and the sum is *finite*. Then

$$u(f) = u\left(\sum_i f \cdot \psi_i\right) = \sum_i u(f \cdot \psi_i) = \sum_i 0 = 0$$

(The fact that the sum is finite allows interchange of summation and evaluation.) ///

[1] That is, the functions ψ_i are smooth, take values between 0 and 1, sum to 1 at all points, and on any compact there are only finitely-many which are non-zero. The existence of such partitions of unity is not completely trivial to prove.