

(January 5, 2020)

08b. $C^\infty(\mathbb{T})$ is not Banach

Paul Garrett garrett@math.umn.edu <http://www.math.umn.edu/~garrett/>

[This document is
http://www.math.umn.edu/~garrett/m/real/notes_2019-20/08b_Cinfy_is_not_Banach.pdf]

1. Countable limits of Banach spaces
2. Factoring through limitand topologies
3. $C^\infty(\mathbb{T})$ is not normable

Many natural function spaces, such as $C^\infty[a, b]$ and $C^\infty(\mathbb{T})$, are *not* Banach, nor even *norm-able* but still do have a metric topology and are complete: these are *Fréchet spaces*, appearing as countable (projective) *limits* of Banach spaces. It is reasonable to ask *why* these spaces are not Banach, and in fact not even *normable*, that is, their topologies cannot be given by a any norm, regardless of metric completeness.

In brief, in tangible terms, the root cause of this impossibility is that no estimates on the first k derivatives of a function on \mathbb{T} give an estimate on the $(k + 1)^{th}$ derivative, for any k . This is discussed precisely below, and abstracted somewhat.

1. Countable limits of Banach spaces

We could take *countable limit of Banach spaces* as the definition of *Fréchet space*.

As earlier, $C^\infty(\mathbb{T})$ is a countable nested intersection, which is a countable (projective) *limit*:

$$C^\infty(\mathbb{T}) = \bigcap_{k \geq 0} C^k(\mathbb{T}) = \lim_k C^k(\mathbb{T})$$

From very general category-theory arguments, *there is at most one projective-limit topology* on $C^\infty(\mathbb{T})$, up to unique isomorphism. Existence of the topology on X satisfying the limit condition can be proven by identifying X as the diagonal *closed subspace* of the *topological product* of the *limitands* X_k : letting $p_{k,k-1} : X_k \rightarrow X_{k-1}$ be the transition maps,

$$X = \{ \{x_k : x_k \in C^k[a, b]\} : p_{k,k-1}(x_k) = x_{k-1} \text{ for all } k \}$$

The subspace topology on X is the limit topology, seen as follows. The projection maps $p_k : \prod_j X_j \rightarrow X_k$ from the whole product to the factors X_k are continuous, so their restrictions to the diagonally imbedded X are continuous. Further, letting $i_k : X_k \rightarrow X_{k-1}$ be the transition map, on that diagonal copy of X we have $i_k \circ p_k = p_{k-1}$ as required.

On the other hand, *any* family of maps $\varphi_k : Z \rightarrow X_k$ induces a map $\tilde{\varphi} : Z \rightarrow \prod X_k$ such that $p_k \circ \tilde{\varphi} = \varphi_k$, by the property of the product. *Compatibility* $i_k \circ \varphi_k = \varphi_{k-1}$ implies that the image of $\tilde{\varphi}$ is inside the diagonal, that is, inside the copy of X . Thus, this construction does produce a limit.

A *countable* product of *metric* spaces X_k with metrics d_k has no canonical single metric, but is *metrizable*. One of many topologically equivalent metrics is the usual

$$d(\{x_k\}, \{y_k\}) = \sum_{k=0}^{\infty} 2^{-k} \frac{d_k(x_k - y_k)}{d_k(x_k - y_k) + 1}$$

When the metric spaces X_k are *complete*, the product is complete. A closed subspace of a complete metrizable space is complete metrizable, so the diagonal X is complete metric.

Even in general, the topologies on vector spaces V are required to be *translation invariant*, meaning that for an open neighborhood U of 0, for any $x \in V$, the set $x + U = \{x + u : u \in U\}$ is an open neighborhood of

x , and vice-versa. ^[1] Thus, to specify the topology on a limit X of Banach spaces X_k , we need only give a local basis at 0. From the construction above, a local basis is given by all sets

$$U_{k,\delta} = \{x \in X : |p_k(x)|_{X_k} < \delta\} \quad (\text{for } \delta > 0 \text{ and index } k)$$

2. Factoring through limitand topologies

[2.1] **Lemma:** Given a continuous linear map T from $C^\infty(\mathbb{T})$ to a *normed space* Y , there is an index k such that when $C^\infty(\mathbb{T})$ is given the (weaker) C^k topology, $T : C^\infty(\mathbb{T}) \rightarrow Y$ is still continuous.

[2.2] **Corollary:** Every continuous linear map T from $C^\infty(\mathbb{T})$ to a Banach space Y factors through some limitand $C^k(\mathbb{T})$. That is, there is $T_k : C^k(\mathbb{T}) \rightarrow Y$ such that $T = T_k \circ i_k$, where $i_k : C^\infty(\mathbb{T}) \rightarrow C^k(\mathbb{T})$ is the inclusion.

Proof: (of Corollary) After applying the lemma, since the target space of T is *complete*, we can extend $T : C^\infty(\mathbb{T}) \rightarrow Y$ by continuity (in the C^k topology) to the C^k -completion of C^∞ , which is C^k . ///

The lemma is a special case of the analogous lemma that has nothing to do with spaces of functions, but, rather, is true for more general reasons:

[2.3] **Lemma:** Let $X = \lim_k X_k$ be a limit of Banach spaces X_k , with projection maps $p_k : X \rightarrow X_k$. Suppose that $p_k(X)$ is *dense* in X_k . Then every continuous linear map $T : X \rightarrow Y$ to a *normed space* Y factors through some limitand X_k . That is, there is $T_k : X_k \rightarrow Y$ such that $T = T_k \circ p_k$.

Proof: Given $\varepsilon > 0$, by the description above of the topology on the limit, there are $\delta > 0$ and index k such that $T(U_{k,\delta})$ is inside the ε -ball at 0 in Y .

Then, given any other $\varepsilon' > 0$, we claim that T maps

$$\frac{\varepsilon'}{\varepsilon} \cdot U_{k,\delta} = U_{k,\delta\varepsilon'/\varepsilon}$$

to the open ε' -ball in Y . Indeed,

$$|T\left(\frac{\varepsilon'}{\varepsilon} \cdot U_{k,\delta}\right)|_Y = \frac{\varepsilon'}{\varepsilon} \cdot |T(U_{k,\delta})|_Y < \frac{\varepsilon'}{\varepsilon} \cdot \varepsilon = \varepsilon'$$

as claimed. Thus, $T : X \rightarrow Y$ is continuous when X is given the X_k topology, for the index k that makes this work. Thus, T extends by continuity to the $|\cdot|_{X_k}$ -completion of X . By the density assumption, this is X_k . ///

[2.4] **Remark:** Finite Fourier series, which are in $C^\infty(\mathbb{T})$, are dense in every $C^k(\mathbb{T})$, so $C^\infty(\mathbb{T})$ is dense in every $C^k(\mathbb{T})$.

[2.5] **Remark:** In the case that $Y = \mathbb{C}$, the density assumption is unnecessary, since Hahn-Banach gives an extension. But for general Banach Y , without the density assumption, we can only conclude that T factors through the $|\cdot|_{X_k}$ -completion of X , since not all closed subspaces of Banach spaces are *complemented*.

[1] For Hilbert and Banach spaces, this translation-invariance is clear, since the topology is metric, and comes from a norm.

3. $C^\infty(\mathbb{T})$ is not normable

If $C^\infty(\mathbb{T})$ were *normable*, then the identity map $j : C^\infty(\mathbb{T}) \rightarrow C^\infty(\mathbb{T})$ would be continuous when the source is given the C^k topology, for some k . In particular, for every $\varepsilon > 0$, there would be a sufficiently small $|\cdot|_{C^k(\mathbb{T})}$ -ball B whose image in $C^\infty(\mathbb{T})$ under the inclusion is inside the ε -ball in the $C^{k+1}(\mathbb{T})$ topology on $C^\infty(\mathbb{T})$. Specifically, for $\varepsilon = 1$, there should be a sufficiently small $\delta > 0$ such that the δ -ball in the C^k topology is inside the unit ball in the C^{k+1} topology.

However, there are C^∞ functions whose C^k norms are arbitrarily small, but whose C^{k+1} norm is 1, for example, $\sin(nx)/n^{k+1}$. Thus, we achieve a contradiction. ///