A HAMILTON-JACOBI EQUATION FOR THE CONTINUUM LIMIT OF NON-DOMINATED SORTING*

JEFF CALDER^{\dagger}, SELIM ESEDOGLU^{\dagger}, AND ALFRED O. HERO^{\ddagger}

Abstract. We show that non-dominated sorting of a sequence X_1, \ldots, X_n of *i.i.d.* random variables in \mathbb{R}^d has a continuum limit that corresponds to solving a Hamilton-Jacobi equation involving the probability density function f of X_i . Non-dominated sorting is a fundamental problem in multi-objective optimization, and is equivalent to finding the canonical antichain partition and to problems involving the longest chain among Euclidean points. As an application of this result, we show that non-dominated sorting is asymptotically stable under bounded random perturbations in X_1, \ldots, X_n . We give a numerical scheme for computing the viscosity solution of this Hamilton-Jacobi equation and present some numerical simulations for various density functions.

Key words. Hamilton-Jacobi equations, viscosity solutions, optimal control, longest chain in Euclidean space, non-dominated sorting, Pareto-optimality

AMS subject classifications. 35F21, 35D40, 60C05, 06A07, 90C27

1. Introduction. Let X_1, \ldots, X_n be *i.i.d.* random variables on \mathbb{R}^d with density function $f \in L^1(\mathbb{R}^d)$. The points form a partially ordered set $\mathscr{X}_n = \{X_1, \ldots, X_n\}$ under the partial order

(1.1)
$$x \leq y \iff x_i \leq y_i \text{ for } i = 1, \dots, d.$$

Let $\ell(n)$ denote the length of a longest chain¹ in \mathscr{X}_n , and for $x \in \mathbb{R}^d$ let $u_n(x)$ denote the length of a longest chain in \mathscr{X}_n consisting of points less than x. We are interested in the asymptotic properties of u_n as $n \to \infty$.

When f is a smooth density on $[0, 1]^d$, hence $\ell(n) = u_n(1, \ldots, 1)$, the problem of studying the asymptotics of $\ell(n)$ has a long history. It begins with Ulam's famous problem [44] of finding the length of a longest increasing subsequence of a random permutation. Hammersley [24] made some of the first breakthroughs in understanding Ulam's problem. He observed that the distribution of the length of a longest increasing subsequence among n numbers chosen uniformly at random is the same as the distribution of $\ell(n)$ for uniformly distributed points on $[0, 1]^2$. Using subadditive ergodic theory, Hammersley showed that $n^{-\frac{1}{2}}\ell(n)$ converges almost surely to a constant c as $n \to \infty$, and he conjectured that c = 2. In subsequent papers, Vershik and Kerov [45] and Logan and Shepp [31] showed that $c \le 2$ and $c \ge 2$, respectively. Hammersley's results were generalized by Bollobás and Winkler [6] to uniformly distributed points on $[0, 1]^d$; they showed that there exist positive constants c_d such that $n^{-\frac{1}{d}}\ell(n) \to c_d$ almost surely as $n \to \infty$, and $c_d \uparrow e$ as $d \to \infty$. The only known values of c_d are $c_1 = 1$ and $c_2 = 2$. Deuschel and Zeitouni [15] generalized Hammersley's results in another direction. For X_1, \ldots, X_n *i.i.d.* on $[0, 1]^2$ with C^1 density function $f : [0, 1]^2 \to \mathbb{R}$, bounded away from zero, they showed that $n^{-\frac{1}{2}}\ell(n) \to 2\overline{J}$ in probability, where \overline{J} is the supremum of the energy

$$J(\boldsymbol{\varphi}) = \int_0^1 \sqrt{\boldsymbol{\varphi}'(\boldsymbol{x}) f(\boldsymbol{x}, \boldsymbol{\varphi}(\boldsymbol{x}))} \, d\boldsymbol{x},$$

over all $\varphi : [0,1] \rightarrow [0,1]$ nondecreasing and right continuous.

^{*}The research described in this paper was partially supported by ARO grant W911NF-09-1-0310 and NSF grants CCF-1217880 and DMS-0914567.

[†]Department of Mathematics, University of Michigan. ({jcalder,esedoglu}@umich.edu)

[‡]Department of Electrical Engineering and Computer Science, University of Michigan. (hero@eecs.umich.edu)

¹A *chain* is a totally ordered subset of \mathscr{X}_n .



FIG. 1.1. Examples of Pareto fronts for $X_1, \ldots X_n$ chosen from the uniform distribution on $[0,1]^2$. In (b), 29 equally spaced fronts are depicted.

There is another motivation for studying the asymptotics of u_n that arises in multiobjective optimization problems. Such problems are of immense importance in many fields of science and engineering, including control theory and path planning [34, 30, 33], gene selection and ranking [41, 27, 26, 28, 17, 18, 19], data clustering [25], database systems [29, 37] and image processing and computer vision [35, 10]. In a discrete multi-objective optimization problem, one has several objective functions $g_i : S \rightarrow [0, \infty)$, where $i = 1, \ldots, d$ and $S = \{x^1, \ldots, x^n\}$ is a finite set, and is tasked with finding an element $x \in S$ that minimizes *all* of the functions simultaneously. This is generally an impossible task, and instead, a family of solutions are obtained based on the notion of Pareto-optimality. A feasible solution $x \in S$ is called *Pareto-optimal* if for every $y \in S$, we have $g_i(y) > g_i(x)$ for some *i*, or $g_i(y) = g_i(x)$ for all *i*; in other words, no other feasible solution is better in every objective. The collection of Pareto-optimal elements is denoted \mathscr{F}_1 and called the *first Pareto front*. It is the most natural notion of solution for a discrete multi-objective optimization problem. If we set $X_i = (g_1(x^i), \ldots, g_d(x^i)) \in \mathbb{R}^d$ for $i = 1, \ldots, n$, then assuming all X_i are distinct, it is not hard to see that

$$x^i \in \mathscr{F}_1 \iff u_n(X_i) = 1$$

The second Pareto front, \mathscr{F}_2 , consists of the Pareto-optimal elements of $S \setminus \mathscr{F}_1$, and in general

$$\mathscr{F}_k$$
 = Pareto optimal elements of $S \setminus \bigcup_{j < k} \mathscr{F}_j$.

The Pareto front that a particular feasible solution lies on is useful for ranking feasible solutions. As before, when the X_i are all distinct we have

$$x^i \in \mathscr{F}_k \iff u_n(X_i) = k.$$

This observation is essential. It says that studying the asymptotic shapes of the Pareto fronts $\mathscr{F}_1, \mathscr{F}_2, \ldots$ is equivalent to studying the longest chain function u_n . Figure 1.1(a) shows the Pareto fronts for n = 50 points uniformly distributed on $[0, 1]^2$, and Figure 1.1(b) shows the Pareto fronts for $n = 10^6$ points. The points X_i that are on the same Pareto front are connected by a continuous staircase curve that represents the jump set of u_n .

In the multi-objective optimization literature, the process of computing the Pareto fronts for a collection of points is called *non-dominated sorting* [13]. In the combinatorics literature,

the partition $S = \mathscr{F}_1 \cup \mathscr{F}_2 \cup \cdots$ is called the *canonical antichain partition* [16]. Although we have described non-dominated sorting in the context of a discrete optimization problem, it is a fundamental tool in continuous optimization as well. Many state of the art algorithms for continuous optimization involve a large number of discrete subproblems, each of which requires non-dominated sorting. The most common examples are the so-called genetic and evolutionary algorithms for continuous multi-objective optimization [13, 20, 21, 12, 42]. The applications of non-dominated sorting are not restricted to optimization; indeed, there are further striking applications in combinatorics [16, 32], molecular biology [38, 1], graph theory [32], Young Tableaux [46, 16] and even in physical layout problems in the design of integrated circuits [1].

The goal of this paper is to study the asymptotics of u_n , and hence the asymptotics of non-dominated sorting. Our main result, Theorem 1.2, states that $n^{-\frac{1}{d}}u_n$ converges almost surely to a continuous function U, which is the viscosity solution of a Hamilton-Jacobi equation. Our proof is based on linking the asymptotics of u_n to a variational problem, which is a generalization of the variational problem discovered by Deuschel and Zeitouni [15] to higher dimensions. The Hamilton-Jacobi equation satisfied by U is the Hamilton-Jacobi-Bellman equation [3] for the corresponding variational problem. We describe our main result in Section 1.1, and postpone the proofs to Sections 2 and 3. In Section 4, we give a numerical scheme for computing U, and show simulation results comparing the level sets of U to Pareto fronts for various density functions.

1.1. Main result. For $x, y \in \mathbb{R}^d$, we write $x \leq y$ if $x \leq y$ and $x \neq y$. When $x_i < y_i$ for i = 1, ..., d, we write x < y, and we set $\mathbb{R}^d_+ = \{x \in \mathbb{R}^d : x > 0\}$. We will always assume $d \geq 2$. For $s, t \in \mathbb{R}$, $s \leq t$ and s < t will retain their usual definitions. Let $\Omega \subset \mathbb{R}^d$ and let $f : \mathbb{R}^d \to [0, \infty)$. We place the following assumptions on f and Ω .

(H1) There exists a continuous nondecreasing function $m: [0, \infty) \to [0, \infty)$ satisfying m(0) = 0 such that

$$|f(x) - f(y)| \le m(|x - y|),$$

for $x, y \in \Omega$, and f(x) = 0 for $x \notin \Omega$,

(H2) $\Omega \subset \mathbb{R}^d_+$ is open and bounded with Lipschitz boundary. Set

(1.2)
$$\mathscr{A} = \left\{ \gamma \in C^1([0,1]; \mathbb{R}^d) : \gamma'(t) \ge 0 \text{ for all } t \in [0,1] \right\}.$$

Recall that $\gamma'(t) \ge 0$ means that $\gamma'_i(t) \ge 0$ for i = 1, ..., d and $\gamma'(t) \ne 0$. Define $J : \mathscr{A} \to [0, \infty)$ by

(1.3)
$$J(\gamma) = \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt,$$

and $U: \mathbb{R}^d \to \mathbb{R}$ by

(1.4)
$$U(x) = \sup_{\gamma \in \mathscr{A}: \gamma(1) \leq x} J(\gamma).$$

We make the following definition.

DEFINITION 1.1. *Given a domain* $\mathcal{O} \subset \mathbb{R}^d$ *, we say that a function* $u : \mathcal{O} \to \mathbb{R}$ *is* Paretomonotone *if* $x \leq y \implies u(x) \leq u(y)$ *for all* $x, y \in \mathcal{O}$.

In Section 2, we show that U is a Pareto-monotone viscosity solution of the Hamilton-Jacobi partial differential equation (PDE)

(1.5)
$$\begin{cases} U_{x_1} \cdots U_{x_d} = \frac{1}{d^d} f & \text{ on } \mathbb{R}^d_+ \\ U = 0 & \text{ on } \partial \mathbb{R}^d_+. \end{cases}$$

The PDE (1.5) should be interpreted as the Hamilton-Jacobi-Bellman equation for the value function U. We note that f need only be Borel-measurable, bounded and have compact support in \mathbb{R}^d_+ for U to be a viscosity solution of (1.5). The stronger assumptions (H1) and (H2) are needed to prove that U is the unique Pareto-monotone viscosity solution of (1.5) (see Theorem 2.10) under an appropriate boundary condition at infinity. Our main result is

THEOREM 1.2. Let f satisfy (H1), let Ω satisfy (H2), and let X_1, \ldots, X_n be i.i.d. with density f. Then there exists a positive constant c_d such that

$$n^{-\frac{1}{d}}u_n \longrightarrow c_d U$$
 in $L^{\infty}(\mathbb{R}^d)$ almost surely.

The constants c_d are the same as those given by Bollobás and Winkler [6]. In particular, $c_1 = 1, c_2 = 2$ and $c_d \uparrow e$ as $d \to \infty$. When f is a product density, i.e., $f(x) = f_1(x_1) \cdots f_d(x_d)$, the value function U is given by

(1.6)
$$U(x) = \left(\int_{0 \le y \le x} f(y) \, dy\right)^{\frac{1}{d}} = \left(\int_{0}^{x_{i}} f_{1}(t) \, dt\right)^{\frac{1}{d}} \cdots \left(\int_{0}^{x_{d}} f_{d}(t) \, dt\right)^{\frac{1}{d}}.$$

For the case f = 1 and d = 2, Aldous and Diaconis [2, p. 204] provided a non-rigorous derivation of (1.5) by viewing the problem as an interacting particle process. They used this to motivate their proof that c = 2 in Ulam's problem, but make no rigorous statements about the relationship between (1.5) and the longest chain problem. A similar, though tangentially related, PDE also appears in growth models in multiple dimensions that are defined through the height of a random partial order [39, p. 209].

Theorem 1.2 provides a new tool with which to study the asymptotics of non-dominated sorting and the longest chain problem. As an example of the applicability of this result, we show in Theorem 3.7 that non-dominated sorting is asymptotically stable under bounded random perturbations. Evidently, Theorem 1.2 reduces the problem of non-dominated sorting to solving a Hamilton-Jacobi equation. From an algorithmic perspective, this may be useful in designing fast approximate algorithms for non-dominated sorting, or finding lengths of longest chains. We study some of these applications in a subsequent paper [7].

1.2. Motivation. As motivation, let us give an informal derivation of the Hamilton-Jacobi PDE (1.5). Suppose $f : \mathbb{R}^d \to \mathbb{R}$ is continuous and $n^{-\frac{1}{d}}u_n \to U \in C^1(\mathbb{R}^d)$ uniformly. Fix *n* large enough so that $n^{-\frac{1}{d}}u_n \approx U$. Then the *k*th Pareto front should be well approximated by the level set $\{y : U(y) = n^{-\frac{1}{d}}k\}$. It is not hard to see that *U* should be Pareto-monotone (recall Definition 1.1), and hence it is reasonable to assume that $U_{x_i} > 0$ for all *i*. Fix $x, v \in \mathbb{R}^d$ with $\langle DU(x), v \rangle > 0$, where DU(x) denotes the gradient of *U* at *x*, and consider the quantity $n^{\frac{1}{d}}(U(x+v) - U(x))$. This is approximately the number of Pareto fronts passing between *x* and x + v. When counting these fronts, we may restrict ourselves to the region

$$A = \{ y : U(y) \ge U(x) \text{ and } y \le x + v \}.$$

This is because any samples in $\{y : U(y) < U(x)\}$ will be on a previous Pareto front and only samples that are less than x + v can influence the Pareto rank of x + v. See Figure 1.2 for a



FIG. 1.2. Some quantities from the informal derivation of the Hamilton-Jacobi PDE (1.5).

depiction of this region and some quantities from the derivation. Since $U_{x_i}(x) > 0$ for all *i*, and *U* is C^1 , *A* is well approximated by a simplex for small |v|, and furthermore, the samples within *A* are approximately uniformly distributed. Let *m* denote the number of samples falling in *A*. By scaling the simplex into a standard simplex, without disrupting the Pareto ordering within *A*, it is reasonable to conjecture that the number of Pareto fronts within *A* (or the length of a longest chain in *A*) is approximately $cm^{\frac{1}{d}}$ for some constant *c*, independent of *x*. For simplicity we take c = 1.

By the law of large numbers, we have $m \approx n \int_A f(y) dy$. Hence when |v| > 0 is small we have

(1.7)
$$n^{\frac{1}{d}}(U(x+v) - U(x)) \approx \left(n \int_{A} f(y) \, dy\right)^{\frac{1}{d}} \approx n^{\frac{1}{d}} |A|^{\frac{1}{d}} f(x)^{\frac{1}{d}},$$

where |A| denotes the Lebesgue measure of *A*. Let ℓ_1, \ldots, ℓ_d denote the side lengths of the simplex *A*. Then $|A| \approx c \ell_1 \cdots \ell_d$ for a constant *c* which we again take to be 1. Since $x + v - \ell_i e_i$ lies approximately on the tangent plane to the level set $\{y : U(y) = U(x)\}$, we see that

$$\langle DU(x), v - \ell_i e_i \rangle \approx 0$$

Rearranging the above we see that $\ell_i \approx U_{x_i}(x)^{-1} \langle DU(x), v \rangle$, and hence

(1.8)
$$|A| \approx U_{x_1}(x)^{-1} \cdots U_{x_d}(x)^{-1} \langle DU(x), v \rangle^d$$

For small |v|, we can combine (1.8) and (1.7) to obtain

$$\langle DU(x),v\rangle\approx U(x+v)-U(x)\approx f(x)^{\frac{1}{d}}U_{x_1}(x)^{-\frac{1}{d}}\cdots U_{x_d}(x)^{-\frac{1}{d}}\langle DU(x),v\rangle.$$

Simplifying, we see that U should satisfy

(1.9)
$$U_{x_1}\cdots U_{x_d} = f \quad \text{on} \quad \mathbb{R}^d,$$

up to scaling by a constant.

Although this derivation is informal, it is straightforward and conveys the essence of the result. It is difficult, however, to construct a rigorous proof based on these heuristics. There

are two main reasons for this. First, it supposes that $n^{-\frac{1}{d}}u_n$ converges to a limit U, which is not obvious. Second, it is essential that $U \in C^1$, as we require A to be an approximate simplex. Solutions of (1.9) are in general not smooth, and can have points of non-differentiability due to crossing characteristics. This is true even in the case that f is smooth, and is related to the geometry of Ω .

2. Analysis of variational problem. Before studying the variational problem (1.4), we recall some aspects of the theory of optimal control [3] that are relevant to our problem. We will describe the infinite horizon optimal control problem, but the discussion below applies with minor modifications to other variants of optimal control, such as finite horizon or undiscounted problems with exit times. The state of the control problem, y(t), is assumed to obey the dynamics

(2.1)
$$\begin{cases} y'(t) = g(y(t), \alpha(t)), & t > 0\\ y(0) = x, \end{cases}$$

where $\alpha : [0,\infty) \to A$ is the control, *A* is a topological space, and $g : \mathbb{R}^d \times A \to \mathbb{R}^d$. Given an initial condition $x \in \mathbb{R}^d$, the solution of (2.1) is denoted $y_x(\cdot)$. Let

$$\mathscr{A} := \{ \text{measurable functions } [0, \infty) \to A \}.$$

The goal in optimal control is to select the control $\alpha \in \mathscr{A}$ to minimize the cost functional

(2.2)
$$J(x,\alpha) := \int_0^\infty c(y_x(t),\alpha(t))e^{-\lambda t} dt$$

where $\lambda > 0$ and $c : \mathbb{R}^d \times A \to \mathbb{R}$. The value function for this problem is

(2.3)
$$v(x) := \inf_{\alpha \in \mathscr{A}} J(x, \alpha).$$

Under sufficient regularity assumptions on c and g (discussed below), the value function is a Hölder- (or Lipschitz) continuous viscosity solution of the Hamilton-Jacobi-Bellman equation

(2.4)
$$\lambda v + H(x, Dv) = 0 \quad \text{on} \quad \mathbb{R}^d,$$

where

(2.5)
$$H(x,p) = \sup_{a \in A} \{-\langle g(x,a), p \rangle - c(x,a)\}.$$

Although the variational problem (1.4) can be cast in this framework, the assumptions on the running cost $c(\cdot, \cdot)$ in the existing literature are too restrictive. For our variational problem, we have $\lambda = 0$, g(x, a) = a, $A = \mathbb{R}^d_+$,

(2.6)
$$c(x,a) = -f(x)^{\frac{1}{d}} (a_1 \cdots a_d)^{\frac{1}{d}},$$

and U(x) = -v(x). In the proofs of Theorems 3.5 and 3.6, we require the standard optimal control theory to hold for *f* piecewise constant on arbitrarily small grids. In the standard reference on optimal control [3], it is assumed that $x \mapsto c(x, a)$ is uniformly continuous. This assumption is then used to prove regularity of the value function *v*. There is relatively little research devoted to relaxing the regularity condition on *c*. There are some results for the optimal control problem associated with the Eikonal equation [36, 8, 14], which allow *c* to have

discontinuities. These results assume that $A = \mathbb{R}^d$ and make essential use of either Lipschitzness of v, or uniform continuity and/or coercivity of $p \mapsto H(x, p)$, none of which hold for the variational problem (1.4). Soravia [40] and Garavello and Soravia [22] considered a running cost of the form $c(x, a) = c_1(x, a) + c_2(x)$, where c_1 is continuous and c_2 is Borel-measurable, and showed that the standard optimal control results hold with minor modifications. This is incompatible with (2.6) when f is not continuous. A similar program is carried out for differential games here [23]. Barles et al. [4] study optimal control on multi-domains, where the discontinuity in c is assumed to lie in a half-space.

Under the assumption that f is compactly supported, bounded and Borel-measurable, the standard results on optimal control hold for the variational problem (1.4) with minor modifications to the proofs. In particular, in Lemma 2.1 we show that U is Hölder-continuous with exponent $\frac{1}{d}$, and in Theorem 2.5 we show that U is a viscosity solution of (1.5). The uniqueness of viscosity solutions of (1.5) under the assumption that f satisfies (H1) is a more delicate problem. This is addressed in Section 2.3.

In our main result, Theorem 1.2, we assume that f satisfies (H1), which is stronger than Borel-measurability. We assume Borel-measurability in much of this section so that our results apply to piecewise constant densities, which are used to approximate f in the proofs of Theorems 3.5 and 3.6. To be more precise, we set

(2.7)
$$\mathscr{B} = \{f : \mathbb{R}^d \to \mathbb{R} : f \text{ is bounded, Borel-measurable, and } \sup(f) \subset [0,1]^d\}$$

We note that the assumption $\operatorname{supp}(f) \subset [0,1]^d$ is not restrictive, as we can make a simple scaling argument to obtain the case where f has compact support in \mathbb{R}^d . We also note that Borel-measurability of f is necessary, as opposed to Lebesgue-measurability, to guarantee that the composition $t \mapsto f(\gamma(t))$ is Lebesgue measurable.

We now introduce some new notation. We will write $\gamma \leq x$ whenever $\gamma(t) \leq x$ for all $t \in [0, 1]$. We write $\gamma_1 \leq \gamma_2$ whenever $\gamma_1(1) \leq \gamma_2(0)$. The same definitions apply to $\leq, <, \geq, \geq$ and > with obvious modifications. For $y \in \mathbb{R}^d$ and r > 0 we set $B_r(y) = \{x \in \mathbb{R}^d : |x-y| < r\}$. For $x, y \in \mathbb{R}^d$ we set

(2.8)
$$w(x,y) = \begin{cases} \sup\{J(\gamma) : \gamma \in \mathscr{A} \text{ and } x \leq \gamma \leq y\} & \text{if } x \leq y\\ 0 & \text{otherwise.} \end{cases}$$

2.1. Basic properties of U. We establish here some basic properties of U. Namely, in Lemma 2.1 we establish Hölder-continuity of U, and in Lemma 2.2, we establish a dynamic programming principle for U.

LEMMA 2.1. Let $f \in \mathscr{B}$. Then U is Hölder-continuous with exponent $\frac{1}{d}$ and Hölder seminorm $[U]_{\frac{1}{d}} \leq \|f\|_{L^{\infty}(\mathbb{R}^d)}^{\frac{1}{d}}$.

Proof. Let $x, z \in \mathbb{R}^d$ and let $\varepsilon > 0$. Choose $\gamma \in \mathscr{A}$ with $\gamma \leq x$ and $J(\gamma) \geq U(x) - \varepsilon$. Since f(x) = 0 for $x \notin [0, 1]^d$, we may assume that $\gamma(t) \in [0, 1]^d$ for all $t \in [0, 1]$. Set

$$s = \sup\{t \in [0,1] : \gamma(t) \leq z\}.$$

If for all $t \in [0, 1]$ we have $\gamma(t) \nleq z$, then set s = 0. We claim that

(2.9)
$$U(z) \ge U(x) - \int_s^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt - \varepsilon.$$

To see this: In the case that s > 0, we have $\gamma(s) \leq z$ and hence

$$U(z) \ge \int_0^s f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt$$

= $J(\gamma) - \int_s^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt$
 $\ge U(x) - \int_s^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt - \varepsilon.$

In the case that s = 0, we have

$$U(z) \ge 0 = J(\gamma) - \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma'_1(t) \cdots \gamma'_d(t))^{\frac{1}{d}} dt$$
$$\ge U(x) - \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma'_1(t) \cdots \gamma'_d(t))^{\frac{1}{d}} dt - \varepsilon.$$

Hence (2.9) is established. Suppose s < 1. Then there must exist *i* such that $\gamma_i(s) \ge z_i$. It follows that

$$\int_s^1 \gamma_i'(t) dt = \gamma_i(1) - \gamma_i(s) \leq x_i - z_i = |x_i - z_i|.$$

Applying the generalized Hölder inequality we see that

$$\begin{split} \int_{s}^{1} f(\gamma(t))^{\frac{1}{d}} (\gamma_{1}'(t) \cdots \gamma_{d}'(t))^{\frac{1}{d}} dt &\leq \|f\|_{L^{\infty}(\mathbb{R}^{d})}^{\frac{1}{d}} \left(\int_{s}^{1} \gamma_{1}'(t) dt \right)^{\frac{1}{d}} \cdots \left(\int_{s}^{1} \gamma_{d}'(t) dt \right)^{\frac{1}{d}} \\ &\leq \|f\|_{L^{\infty}(\mathbb{R}^{d})}^{\frac{1}{d}} |x_{i} - z_{i}|^{\frac{1}{d}} \prod_{j \neq i} (\gamma_{j}(1) - \gamma_{j}(s))^{\frac{1}{d}} \\ &\leq \|f\|_{L^{\infty}(\mathbb{R}^{d})}^{\frac{1}{d}} |x_{i} - z_{i}|^{\frac{1}{d}}. \end{split}$$

Inserting this into 2.9 we obtain

(2.10)
$$U(x) - U(z) \le \|f\|_{L^{\infty}(\mathbb{R}^d)}^{\frac{1}{d}} |x_i - z_i|^{\frac{1}{d}} + \varepsilon \le \|f\|_{L^{\infty}(\mathbb{R}^d)}^{\frac{1}{d}} |x - z|^{\frac{1}{d}} + \varepsilon$$

If s = 1 then inspecting (2.9), we see that $U(x) - U(z) \le \varepsilon$, which implies (2.10). Sending $\varepsilon \to 0$ we find that $U(x) - U(z) \le ||f||_{L^{\infty}(\mathbb{R}^d)}^{\frac{1}{d}} |x - z|^{\frac{1}{d}}$. We can reverse the roles of x and z in the preceding argument to obtain the opposite inequality. \Box

REMARK 1. By a similar argument, we can show that $w : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is Höldercontinuous with exponent 1/d and $[w]_{\frac{1}{d}} \leq ||f||_{L^{\infty}(\mathbb{R}^d)}^{\frac{1}{d}}$. LEMMA 2.2 (Dynamic Programming Principle). Let $f \in \mathscr{B}$. Then for any r > 0 and $y \in \mathbb{R}^d$ we have

(2.11)
$$U(y) = \max_{x \in \partial B_r(y): x \le y} \{ U(x) + w(x, y) \}.$$

Proof. Let us denote the right hand side of (2.11) by v(y). We first show that $U(y) \le v(y)$. Let $\varepsilon > 0$ and let $\gamma \in \mathscr{A}$ such that $\gamma \leq y$ and $J(\gamma) \geq U(y) - \varepsilon$. Suppose that $|\gamma(1) - y| \geq r$. Then there exists $x \in \partial B_r(y)$ such that $\gamma(1) \leq x \leq y$ and hence

$$U(y) \leq J(\gamma) + \varepsilon \leq U(x) + \varepsilon \leq v(y) + \varepsilon.$$

If $|\gamma(0) - y| \le r$ then there exists $x \in \partial B_r(y)$ such that $x \le \gamma \le y$ and hence

$$v(y) \ge w(x,y) \ge J(\gamma) \ge U(y) - \varepsilon.$$

Finally, suppose that $|\gamma(1) - y| < r$ and $|\gamma(0) - y| > r$. Then there exists 0 < s < 1 such that $|\gamma(s) - y| = r$. Set $x = \gamma(s)$ and define $\gamma^1, \gamma^2 \in \mathscr{A}$ by

$$\gamma^1(t) = \gamma(st)$$
 and $\gamma^2(t) = \gamma(s+t(1-s))$ for $t \in [0,1]$.

Note that $\gamma^1 \leq x$ and $x \leq \gamma^2 \leq y$. Since *J* is invariant under a change of parametrization of γ , we see that

$$U(y) \leq J(\gamma) + \varepsilon = J(\gamma^1) + J(\gamma^2) + \varepsilon \leq U(x) + w(x,y) + \varepsilon \leq v(y) + \varepsilon.$$

Sending $\varepsilon \to 0$ we obtain $U(y) \le v(y)$.

We now show that $U(y) \ge v(y)$. By Lemma 2.1 and Remark 1, there exists $x \in \partial B_r(y)$ with $x \le y$ such that

$$v(y) = U(x) + w(x, y).$$

Let $\varepsilon > 0$ and let $\gamma^1, \gamma^2 \in \mathscr{A}$ with $\gamma^1 \leq x$ and $x \leq \gamma^2 \leq y$ such that

$$J(\gamma^1) \ge U(x) - \frac{\varepsilon}{2}$$
 and $J(\gamma^2) \ge w(x,y) - \frac{\varepsilon}{2}$

Since $\gamma^1 \leq \gamma^2 \leq y$, we can concatenate γ^1 and γ^2 to find that $U(y) \geq J(\gamma^1) + J(\gamma^2)$. Thus we have

$$U(y) \ge U(x) + w(x,y) - \varepsilon = v(y) - \varepsilon.$$

Sending $\varepsilon \to 0$ yields $U(y) \ge v(y)$. \Box

2.2. Hamilton-Jacobi-Bellman equation for *U***.** We digress momentarily to recall the definition of viscosity solution of

$$(2.12) H(Du) = f \text{ on } \mathcal{O},$$

where $\mathscr{O} \subset \mathbb{R}^d$ is open, $H : \mathbb{R}^d \to \mathbb{R}$ is continuous, $f : \mathscr{O} \to \mathbb{R}$ is bounded, and $u : \mathscr{O} \to \mathbb{R}$ is the unknown function. For more information on viscosity solutions of Hamilton-Jacobi equations, we refer the reader to [3]. The *superdifferential* of u at $x \in \mathscr{O}$, denoted $D^+u(x)$, is the set of all $p \in \mathbb{R}^d$ satisfying

(2.13)
$$u(y) \le u(x) + \langle p, y - x \rangle + o(|x - y|) \text{ as } \mathcal{O} \ni y \to x.$$

Similarly, the *subdifferential* of *u* at $x \in \mathcal{O}$, denoted $D^-u(x)$, is the set of all $p \in \mathbb{R}^d$ satisfying

(2.14)
$$u(y) \ge u(x) + \langle p, y - x \rangle + o(|x - y|) \text{ as } \mathcal{O} \ni y \to x.$$

Equivalently, we may set

$$D^+u(x) = \{D\varphi(x) : \varphi \in C^1(\mathscr{O}) \text{ and } u - \varphi \text{ has a local max at } x\},\$$

and

$$D^{-}u(x) = \{D\varphi(x) : \varphi \in C^{1}(\mathcal{O}) \text{ and } u - \varphi \text{ has a local min at } x\}.$$

DEFINITION 2.3. A viscosity subsolution of (2.12) is a continuous function $u : \mathcal{O} \to \mathbb{R}$ satisfying

(2.15)
$$H(p) \le f^*(x) \text{ for all } x \in \mathcal{O} \text{ and } p \in D^+u(x).$$

Similarly, a viscosity supersolution of (2.12) is a continuous function $u : \mathcal{O} \to \mathbb{R}$ satisfying

(2.16)
$$H(p) \ge f_*(x) \text{ for all } x \in \mathcal{O} \text{ and } p \in D^-u(x).$$

The functions f_* and f^* are the lower and upper semicontinuous envelopes of f, respectively, defined by

$$f^*(x) = \limsup_{r \searrow 0} \{ f(y) : y \in \mathscr{O} \text{ and } |x - y| \le r \},$$

and $f_* = -(-f)^*$. If *u* is a viscosity subsolution and supersolution of (2.12), then we say that *u* is a *viscosity solution* of (2.12).

After a basic proposition, we establish in Theorem 2.5 that U is a *Pareto-monotone* viscosity solution of (1.5).

PROPOSITION 2.4. Let $\mathcal{O} \subset \mathbb{R}^d$ be open and let $v : \mathcal{O} \to \mathbb{R}$ be continuous and Paretomonotone. Then

$$D^+v(x) \cup D^-v(x) \subset \mathbb{R}^d_+$$
 for all $x \in \mathcal{O}$.

Proof. Let $x \in \mathcal{O}$ and $p \in D^+v(x)$. For any index *i* and small enough t > 0, we have $x \leq x + te_i \in \mathcal{O}$. Since *v* is Pareto-monotone, we have

$$v(x) \leq v(x+te_i) \leq v(x) + p_i t + o(t)$$
 as $t \downarrow 0$.

Hence $p_i \ge o(t)/t$ as $t \searrow 0$ which implies that $p_i \ge 0$. The proof for $D^-v(x)$ is similar.

THEOREM 2.5. Let $f \in \mathcal{B}$. Then the value function U defined by (1.4) is a Paretomonotone viscosity solution of the Hamilton-Jacobi equation

(2.17)
$$U_{x_1}\cdots U_{x_d} = \frac{1}{d^d}f \quad on \quad \mathbb{R}^d.$$

Furthermore, U satisfies

(i) Whenever supp $(f) \subset \{x \in \mathbb{R}^d : 0 \leq x \leq z\}$, we have

$$U(x_1,\ldots,x_d) = U(\min(x_1,z_1),\cdots,\min(x_d,z_d)) \text{ for all } x \in \mathbb{R}^d_+$$

(*ii*) U(x) = 0 for every $x \in \mathbb{R}^d \setminus \mathbb{R}^d_+$.

Proof. It follows from the definition of U (1.4) that U is Pareto-monotone, and (ii) follows from the fact that $supp(f) \subset [0,1]^d$.

For (i), let $z \in \mathbb{R}^d$ such that $\operatorname{supp}(f) \subset \{x \in \mathbb{R}^d : 0 \leq x \leq z\}$ and let $x \in \mathbb{R}^d$ such that $x_i > z_i$ for some *i*. Set $\hat{x} = (\min(x_1, z_1), \dots, \min(x_d, z_d))$. Since *U* is Pareto-monotone we have $U(\hat{x}) \leq U(x)$. Let $\varepsilon > 0$ and $\gamma \in \mathscr{A}$ such that $\gamma \leq x$ and $U(x) \leq J(\gamma) + \varepsilon$. Let

$$s = \sup\{t : \gamma(t) \leq \hat{x}\}.$$

If for all $t \in [0,1]$ we have $\gamma(t) \not\leq \hat{x}$, then set s = 0. If s = 1, then $\gamma \leq \hat{x}$ and hence $U(x) \leq J(\gamma) + \varepsilon \leq U(\hat{x}) + \varepsilon$. If s = 0 then for every $t \in [0,1]$, $\gamma(t) \notin \operatorname{supp}(f)$, and hence $J(\gamma) = 0$. It follows that

$$U(x) \leq J(\gamma) + \varepsilon = \varepsilon \leq U(\hat{x}) + \varepsilon.$$

10

If 0 < s < 1, then for any t > s, $\gamma_i(t) > z_i$ for some *i*, and hence $f(\gamma(t)) = 0$. Set $\gamma^1(t) = \gamma(st)$ for $t \in [0,1]$. Then $\gamma^1 \leq \hat{x}$ and $J(\gamma) = J(\gamma^1)$, hence $U(x) \leq J(\gamma) + \varepsilon = J(\gamma^1) + \varepsilon \leq U(\hat{x}) + \varepsilon$. Sending $\varepsilon \to 0$ we see that $U(x) \leq U(\hat{x})$ and hence $U(x) = U(\hat{x})$.

We now show that U is a viscosity supersolution of (2.17). Let $y \in \mathbb{R}^d$, let $a \in \mathbb{R}^d_+$, and set $\gamma(t) = y - a(1-t)$. By Lemma 2.2 we have

(2.18)
$$U(y) \ge U(y - a(1 - t)) + \int_{t}^{1} f(y - a(1 - s))^{\frac{1}{d}} (a_{1} \cdots a_{d})^{\frac{1}{d}} ds$$
$$\ge U(y - a(1 - t)) + (1 - t) f_{*}(y) (a_{1} \cdots a_{d})^{\frac{1}{d}} + o(1 - t) \text{ as } t \nearrow 1.$$

Let $p \in D^-U(y)$. Since $y - a(1-t) \to y$ as $t \uparrow 1$, we have

$$\langle p, (1-t)a \rangle \stackrel{(2.14)}{\geq} U(y) - U(y - a(1-t)) + o(1-t)$$

 $\stackrel{(2.18)}{\geq} (1-t)f_*(y)(a_1 \cdots a_d)^{\frac{1}{d}} + o(1-t) \text{ as } t \uparrow 1.$

Sending t
eq 1 we obtain

$$\langle p,a\rangle \geq f_*(y)^{\frac{1}{d}}(a_1\cdots a_d)^{\frac{1}{d}}.$$

Since a > 0 was arbitrary, we obtain

(2.19)
$$\sup_{a>0} \left\{ -\langle p, a \rangle + f_*(y)^{\frac{1}{d}} (a_1 \cdots a_d)^{\frac{1}{d}} \right\} \le 0.$$

Since U is Pareto-monotone, Proposition 2.4 yields $p \ge 0$. Hence if $f_*(y) = 0$ then U is trivially a viscosity supersolution of (2.17) at y. We may therefore suppose that $f_*(y) > 0$. Fix i and set $a_j = 1$ for $j \ne i$. By (2.19) we have

$$\sup_{a_i>0} \left\{ -\sum_{j\neq i} p_j + a_i^{\frac{1}{d}} f_*(y)^{\frac{1}{d}} - a_i p_i \right\} \le 0.$$

Since a_i can be arbitrarily large, we must have $p_i > 0$ for the above to hold. Substituting $a_i = p_i^{-1}$ into (2.19) and simplifying we obtain

$$p_1\cdots p_d \geq \frac{1}{d^d}f_*(y).$$

Thus U is a viscosity supersolution of (2.17).

We now show that *U* is a viscosity subsolution of (2.17). Let $y \in \mathbb{R}^d$, let $\varepsilon > 0$, and let $p \in D^+U(y)$. By Lemmas 2.1 and 2.2 and Remark 1, for every r > 0 there exists $x \in \partial B_r(y)$ with $x \leq y$ such that U(y) = U(x) + w(x, y). Hence there exists $\gamma \in \mathscr{A}$ with $x \leq \gamma \leq y$ such that

$$U(y) \leq \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt + U(x) + \varepsilon r.$$

By Hölder's inequality

$$\begin{split} U(y) - U(x) &\leq \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt + \varepsilon r \\ &\leq (f^*(y)^{\frac{1}{d}} + o(1)) \left(\int_0^1 \gamma_1'(t) dt \right)^{\frac{1}{d}} \cdots \left(\int_0^1 \gamma_d'(t) dt \right)^{\frac{1}{d}} + \varepsilon r \\ &\leq f^*(y)^{\frac{1}{d}} |x_1 - y_1|^{\frac{1}{d}} \cdots |x_d - y_d|^{\frac{1}{d}} + o(r) + \varepsilon r, \end{split}$$

as $r \searrow 0$. Since $x \rightarrow y$ as $r \searrow 0$, we have

$$\langle p, y-x \rangle \stackrel{(2.13)}{\leq} U(y) - U(x) + o(r) \leq f^*(y)^{\frac{1}{d}} |x_1 - y_1|^{\frac{1}{d}} \cdots |x_d - y_d|^{\frac{1}{d}} + o(r) + \varepsilon r,$$

as $r \downarrow 0$. Choose r > 0 small enough so that $o(r)/r \le \varepsilon$, and set a = (y - x)/r. Then we have

$$-\langle p,a\rangle + f^*(y)^{\frac{1}{d}}(a_1\cdots a_d)^{\frac{1}{d}} \ge -2\varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, we see that

$$\sup_{a \ge 0: |a|=1} \left\{ -\langle p, a \rangle + f^*(y)^{\frac{1}{d}} (a_1 \cdots a_d)^{\frac{1}{d}} \right\} \ge 0.$$

Since U is Pareto-monotone, we have $p \ge 0$. If $p_i = 0$ for some *i*, then $p_1 \cdots p_d \le f^*(y)/d^d$. Thus we may assume that $p_i > 0$ for all *i*. Then the supremum above is attained at some a > 0 with |a| = 1. By scaling *a* so that $a_1 \cdots a_d = 1$, we see that

(2.20)
$$\sup_{a>0:a_1\cdots a_d=1}\left\{-\langle p,a\rangle+f^*(y)^{\frac{1}{d}}\right\}\geq 0.$$

Since $p_i > 0$ for all *i*, we have that

$$\limsup_{|a|\to\infty,\ a>0} -\langle p,a\rangle + f^*(y)^{\frac{1}{d}} = -\infty.$$

It follows that the supremum in (2.20) is attained at some $a^* > 0$. Introducing a Lagrange multiplier $\lambda > 0$, the necessary conditions for a^* to be a maximizer of the above constrained optimization problem are

$$p_i = \frac{\lambda}{a_i^*}$$
 for all $i \in \{1, \dots, d\}$ and $a_1^* \cdots a_d^* = 1$.

It follows that $\lambda = (p_1 \cdots p_d)^{\frac{1}{d}}$ and $a_i^* = p_i^{-1} (p_1 \cdots p_d)^{\frac{1}{d}}$. Substituting this into (2.20) we have

$$p_1 \cdots p_d \le \frac{1}{d^d} f^*(y),$$

which completes the proof. \Box

REMARK 2. We remark that U satisfies an important truncation property. Namely, if we fix $z \in \mathbb{R}^d$ and define $\widetilde{U}, \widehat{f} : \mathbb{R}^d \to \mathbb{R}$ by

$$\widetilde{U}(x_1, \dots, x_d) = U(\min(x_1, z_1), \dots, \min(x_d, z_d))$$
$$\widehat{f}(x) = \begin{cases} f^*(x) & \text{if } x \leq z \\ 0 & \text{otherwise,} \end{cases}$$

then we have

$$\widetilde{U}_{x_1}\cdots\widetilde{U}_{x_d}=\widehat{f}$$
 on \mathbb{R}^d

in the viscosity sense. Indeed, this follows directly from Theorem 2.5 by noting that supp $(\hat{f}) \subset \{x \in \mathbb{R}^d : 0 \leq x \leq z\}$ and

$$\widetilde{U}(x) = \sup_{\gamma \in \mathscr{A}: \gamma \leq x} \int_0^1 \widehat{f}(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt.$$

12

2.3. Comparison principle. We aim here to establish that U is the unique viscosity solution of (1.5) under hypotheses (H1) and (H2) on f and Ω , which in general allow f to be discontinuous. The standard results on uniqueness of viscosity solutions [3, 11] assume uniformly continuous dependence on spatial variables. There has been some recent work relaxing this condition, as it is important in many applications. Tourin [43] considered Hamilton-Jacobi equations of the form H(x,Du) = 0, where $x \mapsto H(x,p)$ is allowed to have a discontinuity along a smooth surface, and proved a comparison principle under the assumption that $p \mapsto H(x,p)$ is convex and uniformly continuous. Neither assumption holds for (1.5), although the non-convexity can be easily remedied. Deckelnick and Elliot [14] prove a comparison principle for Lipschitz viscosity solutions of Eikonal-type equations of the form H(Du) = f, where f satisfies a regularity condition similar to (H1), but slightly more general. As exhibited by the solution $U(x) = (x_1 \cdots x_d)^{\frac{1}{d}}$ of (1.5) for f = 1, solutions of (1.5) are not in general Lipschitz continuous. Camilli and Siconolfi [9] proposed a new notion of viscosity solution for Hamilton-Jacobi equations in which H has measurable dependence on the spatial variable x. They obtain general uniqueness results under the assumption that $p \mapsto H(x, p)$ is quasiconvex and coercive. Their results do not apply to (1.5) due to the coercivity assumption.

The main result in this section, Theorem 2.10, establishes uniqueness for (1.5) under hypotheses (H1) and (H2), and a boundary condition at infinity. Let us give a sketch of the proof now. Let u be a Pareto-monotone viscosity solution of (1.5). We first prove a standard comparison principle, in Theorem 2.6, for uniformly continuous f. We can then define the regularized value functions U_{ε} and U^{ε} by replacing f by its inf and sup convolutions f_{ε} and f^{ε} , respectively, in (1.4). Since f^{ε} and f_{ε} are Lipschitz continuous, the comparison principle from Theorem 2.6 yields $U_{\varepsilon} \le u \le U^{\varepsilon}$. The proof is completed by showing that $U_{\varepsilon}, U^{\varepsilon} \to U$ as $\varepsilon \to 0$, where U is the value function defined by (1.4). We establish a more general result in Lemma 2.9, the proof of which relies on the second comparison principle, Theorem 2.8. This comparison principle holds for f and Ω satisfying (H1) and (H2) under the additional assumption that the subsolution is truncatable, as per Definition 2.7. As pointed out in Remark 2, the value function U is truncatable, so Theorem 2.8 is applicable in the proof of Lemma 2.9.

THEOREM 2.6. Suppose $f : \mathbb{R}^d_+ \to \mathbb{R}$ is uniformly continuous with $\operatorname{supp}(f) \subset [0,1]^d$. Let *u* and *v* be viscosity sub- and supersolutions, respectively, of

$$(2.21) u_{x_1} \cdots u_{x_d} = f \quad on \quad \mathbb{R}^d_+,$$

and suppose that

(2.22)
$$u(x) = u(\min(x_1, 1), \dots, \min(x_d, 1)) \text{ for } x \in \mathbb{R}^d_+,$$

and v is Pareto-monotone. If $u \leq v$ on $\partial \mathbb{R}^d_+$ then $u \leq v$ on \mathbb{R}^d_+ .

The proof of Theorem 2.6 utilizes the method of doubling the variables, which is standard in the theory of viscosity solutions [3], with appropriate modifications for the boundary condition (2.22).

Proof. For $\theta > 0$, set $v_{\theta}(x) = v(x) + \theta^{\frac{1}{d}} \langle x, \mathbf{1}_d \rangle$, where $\mathbf{1}_d = (1, \dots, 1) \in \mathbb{R}^d$. Fix $x \in \mathbb{R}^d_+$ and $p \in D^- v_{\theta}(x)$. It is easy to see that $p - \theta^{\frac{1}{d}} \mathbf{1}_d \in D^- v(x)$. Hence we have

$$(p_1 - \boldsymbol{\theta}^{\frac{1}{d}}) \cdots (p_d - \boldsymbol{\theta}^{\frac{1}{d}}) - f(x) \ge 0.$$

Since *v* is Pareto-monotone, we have $p_i - \theta^{\frac{1}{d}} \ge 0$ for all *i*, and therefore

$$p_1 \cdots p_d = (p_1 - \theta^{\frac{1}{d}} + \theta^{\frac{1}{d}}) \cdots (p_d - \theta^{\frac{1}{d}} + \theta^{\frac{1}{d}}) \ge (p_1 - \theta^{\frac{1}{d}}) \cdots (p_d - \theta^{\frac{1}{d}}) + \theta$$
$$\ge f(x) + \theta.$$

Hence v_{θ} is a viscosity supersolution of

$$v_{\theta,x_1}\cdots v_{\theta,x_d} = f + \theta$$
 on \mathbb{R}^d_+ ,

and $u \leq v_{\theta}$ on $\partial \mathbb{R}^d_+$.

Suppose that $\delta := \sup_{\mathbb{R}^d_+} (u - v_{\theta}) > 0$. For $x, y \in \mathbb{R}^d_+$ and $\alpha > 0$, set

$$\Phi_{\alpha}(x,y) = u(x) - v_{\theta}(y) - \alpha |x-y|^2,$$

and $M_{\alpha} = \sup_{\mathbb{R}^{d}_{+} \times \mathbb{R}^{d}_{+}} \Phi_{\alpha}$. Setting x = y in $\Phi_{\alpha}(x, y)$ we see that $M_{\alpha} \ge \delta > 0$ for all α . Let

$$\hat{x} = (\min(x_1, 1), \dots, \min(x_d, 1))$$
 and $\hat{y} = (\min(y_1, 1), \dots, \min(y_d, 1))$

Note that $|\hat{x} - \hat{y}| \leq |x - y|$, $u(\hat{x}) = u(x)$ (by (2.22)) and $v_{\theta}(\hat{y}) \leq v_{\theta}(y)$ for all $x, y \in \mathbb{R}^{d}_{+}$. It follows that $\Phi_{\alpha}(\hat{x}, \hat{y}) \geq \Phi_{\alpha}(x, y)$, and hence Φ_{α} attains a maximum at some $(x_{\alpha}, y_{\alpha}) \in [0, 1]^{d} \times [0, 1]^{d}$. Since $M_{\alpha} \geq \delta$, we have

(2.23)
$$u(x_{\alpha}) - v_{\theta}(y_{\alpha}) \ge \delta + \alpha |x_{\alpha} - y_{\alpha}|^{2}.$$

It follows that

(2.24)
$$|x_{\alpha} - y_{\alpha}| \leq \frac{1}{\sqrt{\alpha}} \left(\|u\|_{L^{\infty}((0,1)^{d})} - v_{\theta}(0) \right)^{\frac{1}{2}}.$$

Since $u \leq v_{\theta}$ on $\partial \mathbb{R}^{d}_{+}$ and $(x, y) \mapsto u(x) - v_{\theta}(y)$ is continuous, it follows from (2.23) and (2.24) that $(x_{\alpha}, y_{\alpha}) \in (0, 1]^{d} \times (0, 1]^{d}$ for α large enough. For such α we have $p := 2\alpha(x_{\alpha} - y_{\alpha}) \in D^{+}u(x_{\alpha}) \cap D^{-}v_{\theta}(y_{\alpha})$, and hence

$$p_1 \cdots p_d \leq f(x_\alpha)$$
 and $p_1 \cdots p_d \geq f(y_\alpha) + \theta$.

Subtracting the above equations yields $f(x_{\alpha}) - f(y_{\alpha}) \ge \theta > 0$, which contradicts the uniform continuity of f and (2.24) as $\alpha \to \infty$. Therefore $u \le v_{\theta}$ on \mathbb{R}^d_+ . Sending $\theta \to 0$ completes the proof. \Box

We can prove a comparison principle for discontinuous f by assuming that the subsolution satisfies the truncation property described in Remark 2. For this, we make the following definition.

DEFINITION 2.7. Let u be a viscosity subsolution of

$$(2.25) u_{x_1} \cdots u_{x_d} = f \quad on \quad \mathbb{R}^d_+.$$

We say that u is truncatable if for every $z \in \mathbb{R}^d_+$, \widetilde{u} is a viscosity subsolution of

$$\widetilde{u}_{x_1}\cdots\widetilde{u}_{x_d}=\widehat{f} \quad on \quad \mathbb{R}^d_+,$$

where \tilde{u} and \hat{f} are defined in Remark 2.

We note that that truncatability is well-defined, i.e., it depends only on f^* . By Remark 2, the value function U is truncatable. It is easy to see that every C^1 Pareto-monotone subsolution of (2.25) is truncatable. It turns out, thanks to Theorem 2.10, that every Pareto-monotone viscosity solution of (2.25) satisfying (2.22) is truncatable.

We place the following assumptions on $f : \mathbb{R}^d_+ \to [0, \infty)$ and Ω .

(H1) There exists a continuous nondecreasing function $m: [0, \infty) \to [0, \infty)$ satisfying m(0) = 0 such that

$$|f(x) - f(y)| \le m(|x - y|),$$

for $x, y \in \Omega$, and f(x) = 0 for $x \notin \Omega$.

(H2) $\Omega \subset \mathbb{R}^d_+$ is open and bounded with Lipschitz boundary. In particular, since Ω is open (H1) implies that $f = f_*$ on \mathbb{R}^d_+ . The assumptions on Ω imply that the following cone condition is satisfied.

(H2^{*}) For every $x \in \partial \Omega$, there exists a cone \mathscr{K}_x with nonempty interior and a neighborhood V_x of x such that

$$y \in V_x \setminus \Omega \implies (y + \mathscr{K}_x) \cap \overline{\Omega} \cap V_x \subset \{y\}$$

To see this: For any $x \in \partial \Omega$ there exists, by Lipchitzness of $\partial \Omega$, a real number r > 0 and a Lipschitz continuous function $\Psi : \mathbb{R}^{d-1} \to \mathbb{R}$ such that, upon relabelling and reorienting the coordinate axes if necessary, we have

$$\Omega \cap B_r(x) = \{ y \in B_r(x) : y_d < \Psi(y_1, \dots, y_{d-1}) \}.$$

One can check that the cone

$$\mathscr{K}_x = \left\{ y \in \mathbb{R}^d : y_d \ge 2 \mathrm{Lip}(\Psi) \sqrt{y_1^2 + \cdots + y_{d-1}^2} \right\}$$

satisfies (H2^{*}). As it is more useful in the comparison principle proof, we will assume that (H2^{*}) holds instead of Lipschitzness of the boundary. We note that the cone condition (H2^{*}) is similar to the one used by Deckelnick and Elliot [14, p. 331].

THEOREM 2.8. Suppose that Ω satisfies (H2^{*}) and f satisfies (H1). Let u and v be viscosity sub- and supersolutions, respectively, of

$$(2.26) u_{x_1} \cdots u_{x_d} = f \quad on \quad \mathbb{R}^d_+,$$

and assume that u is truncatable and v is Pareto-monotone. Then $u \leq v$ on $\partial \mathbb{R}^d_+$ implies that $u \leq v$ on \mathbb{R}^d_+ .

As in the proof of Theorem 2.6, the proof below is based on the standard technique of doubling the variables [11]. The proof is similar to [14, Theorem 2.3] in the way that $(H2^*)$ is used, however, we cannot assume Lipschitzness of *v*. The truncatability condition on *u* in a sense replaces the Lipschitz condition on *v* in [14, Theorem 2.3].

Proof. For $\theta > 0$, set $v_{\theta}(x) = v(x) + \theta^{\frac{1}{d}} \langle x, \mathbf{1}_{d} \rangle + \theta$. Then $u < v_{\theta}$ on $\partial \mathbb{R}^{d}_{+}$. As in the proof of Theorem 2.6, v_{θ} is a viscosity supersolution of

(2.27)
$$v_{\theta,x_1} \cdots v_{\theta,x_d} = f + \theta \quad \text{on} \quad \mathbb{R}^d_+.$$

Now suppose that $\sup_{\mathbb{R}^d_+} (u - v_{\theta}) > 0$ and let

$$R = \sup\left\{r > 0 : u \le v_{\theta} \text{ on } \mathbb{R}^d_+ \cap B_r(0)\right\}.$$

Since $u < v_{\theta}$ on $\partial \mathbb{R}^d_+$ and $u - v_{\theta}$ is continuous, we see that $0 < R < \infty$. By the definition of R, there exists $z_0 \in \partial B_R(0) \cap \mathbb{R}^d_+$ such that $u(z_0) = v_{\theta}(z_0)$ and every neighborhood of z_0 contains a point y such that $u(y) > v_{\theta}(y)$. For r > 0 set

$$H = \{ x \in \mathbb{R}^d : z_0 - r \mathbf{1}_d < x < z_0 + r^2 \mathbf{1}_d \}.$$

and note that $\sup_{H}(u - v_{\theta}) > 0$ for any r > 0.

Notice that we may assume (H2^{*}) holds at any $x \in \mathbb{R}^d_+$. Indeed, if $x \notin \partial \Omega$ then we may set $V_x = B_{\sigma}(x)$ and choose $\sigma > 0$ small enough so that $\partial \Omega \cap V_x = \emptyset$. Then any cone \mathscr{K}_x will

suffice as either $V_x \setminus \Omega = \emptyset$ or $V_x \cap \overline{\Omega} = \emptyset$. Let $\eta \in S^{d-1}$ be in the interior of \mathscr{K}_{z_0} . For $x \in \mathbb{R}^d_+$ set

$$\widetilde{u}(x_1,...,x_d) = u(\min(x_1,z_{0,1}+r^2),...,\min(x_d,z_{0,d}+r^2))$$

For $\alpha > 0$ and $(x, y) \in \mathbb{R}^d_+ \times \mathbb{R}^d_+$, set

(2.28)
$$\Phi_{\alpha}(x,y) = \widetilde{u}(x) - v_{\theta}(y) - \alpha \left| x - y - \frac{1}{\sqrt{\alpha}} \eta \right|^{2},$$

and $M_{\alpha} = \sup_{\mathbb{R}^d_+ \times \mathbb{R}^d_+} \Phi_{\alpha}$. By continuity of \widetilde{u} and v_{θ} we have

(2.29)
$$\varepsilon := \liminf_{\alpha \to \infty} M_{\alpha} \ge \sup_{\mathbb{R}^d_+} (\widetilde{u} - v_{\theta}) > 0.$$

Set

$$D = \{x \in \mathbb{R}^d_+ : x_i \le z_{0,i} - r \text{ for some } i\}.$$

Let $x \in D$ such that $x \leq z_0 + r^2 \mathbf{1}_d$. Then there exists *i* such that $x_i \leq z_{0,i} - r$ and we have

$$|x|^2 \le (z_{0,i}-r)^2 + \sum_{j \ne i} (z_{0,j}+r^2)^2 = |z|^2 - 2z_{0,i}r + O(r^2).$$

Since |z| = R and $z_{0,i} > 0$, we can choose r > 0 small enough so that $x \in B_R(0)$. Fixing such an r > 0 we have

(2.30)
$$\widetilde{u} \le v_{\theta} \text{ on } D \cap \{x \in \mathbb{R}^d_+ : x \le z_0 + r^2 \mathbf{1}_d\}.$$

Since \tilde{u} and v_{θ} are uniformly continuous on compact sets, it follows from (2.30) that there exists $\delta > 0$ such that

$$D \ni x \leq z_0 + r^2 \mathbf{1}_d$$
 and $|x - y| \leq \delta \implies \widetilde{u}(x) - v_\theta(y) \leq \frac{\varepsilon}{2}$

for every $x, y \in \mathbb{R}^d_+$. Now let $x \in D$ and $y \in \mathbb{R}^d_+$ with $|x - y| \le \delta$, and set

$$\hat{x} = (\min(x_1, z_{0,1} + r^2), \dots, \min(x_d, z_{0,d} + r^2))$$
$$\hat{y} = (\min(y_1, z_{0,1} + r^2), \dots, \min(y_d, z_{0,d} + r^2))$$

Then $\hat{x} \in D$, $\hat{x} \leq z_0 + r^2 \mathbf{1}_d$, $|\hat{x} - \hat{y}| \leq |x - y| \leq \delta$, $\tilde{u}(x) = \tilde{u}(\hat{x})$ and $v_{\theta}(y) \geq v_{\theta}(\hat{y})$. We conclude that

$$\widetilde{u}(x) - v_{\theta}(y) \leq \widetilde{u}(\hat{x}) - v_{\theta}(\hat{y}) \leq \frac{\varepsilon}{2}.$$

Hence we have shown that for any $x, y \in \mathbb{R}^d_+$

(2.31)
$$x \in D \text{ and } |x-y| \le \delta \implies \widetilde{u}(x) - v_{\theta}(y) \le \frac{\varepsilon}{2}.$$

Fix $\alpha > 0$ large enough so that $M_{\alpha} > \varepsilon/2$ and let $(x, y) \in \mathbb{R}^d_+ \times \mathbb{R}^d_+$ such that $\Phi_{\alpha}(x, y) > \varepsilon/2$. Then we have

$$\widetilde{u}(x) - v(y) - \alpha \left| x - y - \frac{1}{\sqrt{\alpha}} \eta \right|^2 > \frac{\varepsilon}{2}.$$

In particular, we have $\widetilde{u}(x) - v_{\theta}(y) > \frac{\varepsilon}{2}$, and

$$|x-y| \leq \frac{1}{\sqrt{\alpha}} \left(\left| \|\widetilde{u}\|_{L^{\infty}(\mathbb{R}^d_+)} - v_{\theta}(0) \right|^{\frac{1}{2}} + 1 \right).$$

Taking α large enough so that $|x - y| \le \delta$, we see from (2.31) that $x \ge z_0 - r\mathbf{1}_d$. Now set

$$\hat{x} = (\min(x_1, z_{0,1} + r^2), \dots, \min(x_d, z_{0,d} + r^2)),$$

and $\hat{y} = y + \hat{x} - x$. Then we have $\hat{x} \in \overline{H}$, $\hat{x} - \hat{y} = x - y$, $\widetilde{u}(\hat{x}) = \widetilde{u}(x)$, and $v_{\theta}(\hat{y}) \le v_{\theta}(y)$. It follows that $\Phi_{\alpha}(\hat{x}, \hat{y}) \ge \Phi_{\alpha}(x, y)$. Hence for $\alpha > 0$ large enough, Φ_{α} attains a global maximum at $(x_{\alpha}, y_{\alpha}) \in \mathbb{R}^{d}_{+} \times \mathbb{R}^{d}_{+}$ satisfying $x_{\alpha} \in \overline{H}$ and $|x_{\alpha} - y_{\alpha}| \le C/\sqrt{\alpha}$.

Sending $\alpha \rightarrow \infty$ and extracting a subsequence, if necessary, we may assume that

$$x_{\alpha} \rightarrow x_0$$
 and $y_{\alpha} \rightarrow x_0$ as $\alpha \rightarrow \infty$,

for some $x_0 \in \overline{H}$. Since

$$\widetilde{u}(x_0) - v_{\theta}\left(x_0 - \frac{1}{\sqrt{\alpha}}\eta\right) \leq M_{\alpha} \leq \widetilde{u}(x_{\alpha}) - v_{\theta}(y_{\alpha}),$$

we see, by the continuity of \tilde{u} and v_{θ} , that

$$\lim_{\alpha\to\infty}M_{\alpha}=\widetilde{u}(x_0)-v_{\theta}(x_0).$$

It follows that

(2.32)
$$\alpha \left| x_{\alpha} - y_{\alpha} - \frac{1}{\sqrt{\alpha}} \eta \right|^{2} \to 0 \text{ as } \alpha \to \infty.$$

Let

$$p=2\alpha\left(x_{\alpha}-y_{\alpha}-\frac{1}{\sqrt{\alpha}}\eta\right).$$

Since $(x_{\alpha}, y_{\alpha}) \in \mathbb{R}^d_+ \times \mathbb{R}^d_+$ is a local max of Φ_{α} , we have $p \in D^+ \widetilde{u}(x_{\alpha}) \cap D^- v_{\theta}(y_{\alpha})$. Since \widetilde{u} is a truncatable, we have

$$p_1 \cdots p_d \leq (f)^*(x_\alpha) \leq f^*(x_\alpha)$$

where $\hat{f}(x) = f^*(x)$ for $x \leq z_0 + r^2 \mathbf{1}_d$ and $\hat{f}(x) = 0$ otherwise. By (2.27) we have

$$p_1 \cdots p_d \geq f_*(y_\alpha) + \boldsymbol{\theta}$$

Combining these we see that

(2.33)
$$f^*(x_{\alpha}) - f_*(y_{\alpha}) \ge \theta.$$

If $x_0 \notin \partial \Omega$, then for α large enough we have either $x_{\alpha}, y_{\alpha} \in \Omega$ or $x_{\alpha}, y_{\alpha} \in \mathbb{R}^d_+ \setminus \overline{\Omega}$. Hence by (H1) we have

(2.34)
$$f^*(x_{\alpha}) - f_*(y_{\alpha}) \le m(|x_{\alpha} - y_{\alpha}|).$$

Suppose now that $x_0 \in \partial \Omega$. We have two cases; either (1) $y_{\alpha} \in \Omega$ or (2) $y_{\alpha} \in \mathbb{R}^d_+ \setminus \Omega$.

Case 1. In this case we have $f_*(y_\alpha) = f(y_\alpha)$. If $x_\alpha \in \overline{\Omega}$ then $f^*(x_\alpha) = f(x_\alpha)$ and hence (2.34) holds. If $x_{\alpha} \in \mathbb{R}^{d}_{+} \setminus \overline{\Omega}$ then $f^{*}(x_{\alpha}) = 0$ and (2.34) holds trivially. Case 2. Set $w_{\alpha} = x_{\alpha} - y_{\alpha} - \frac{1}{\sqrt{\alpha}}\eta$ and note that

$$x_{\alpha} = y_{\alpha} + \frac{1}{\sqrt{\alpha}}(\eta + \sqrt{\alpha}w_{\alpha}).$$

By (2.32) we have $\sqrt{\alpha}w_{\alpha} \to 0$ as $\alpha \to \infty$. Since η is in the interior of \mathscr{K}_{z_0} , we see that $\eta + \sqrt{\alpha}w_{\alpha} \in \mathscr{K}_{z_0}$ for α large enough, and hence $x_{\alpha} \in y_{\alpha} + \mathscr{K}_{z_0}$. We can take r > 0 smaller, if necessary, so that $\overline{H} \in V_{z_0}$. Since $x_{\alpha} \in \overline{H}$, we can choose α large enough so that $y_{\alpha} \in V_{z_0}$. Since $x_{\alpha} \neq y_{\alpha}$ for α large enough, we have by (H2^{*}) that $x_{\alpha} \in \mathbb{R}^{d}_{+} \setminus \overline{\Omega}$ and hence $f^{*}(x_{\alpha}) = 0$ and (2.34) holds. Sending $\alpha \to 0$ in (2.34) contradicts (2.33), hence $u \le v_{\theta}$ on \mathbb{R}^{d}_{+} . Sending $\theta \to 0$ we find that $u \leq v$ on \mathbb{R}^d_+ .

In order to prove a general uniqueness result, Theorem 2.10, we require a perturbation result for the value function U with respect to sup and inf convolutions of the density f. Since a similar result, for a different type of perturbation, is required in the proof of our main result, Theorem 1.2, we state a more general result in Lemma 2.9. We first recall some notation standard in the theory of viscosity solutions. For a sequence of bounded functions $f_n: \mathbb{R}^d_+ \to \mathbb{R}$, the upper and lower limits are defined by

$$\limsup_{n \to \infty} f_n(x) := \lim_{j \to \infty} \sup \left\{ f_n(y) : n \ge j, \ y \in \mathbb{R}^d, \ \text{ and } \ |x - y| \le \frac{1}{j} \right\},$$

and

$$\liminf_{n\to\infty} f_n(x) := \liminf_{j\to\infty} \left\{ f_n(y) : n \ge j, \ y \in \mathbb{R}^d, \ \text{ and } \ |x-y| \le \frac{1}{j} \right\}.$$

LEMMA 2.9. Let $\{f_n\}_{n=1}^{\infty} \subset \mathscr{B}$ and suppose that Ω satisfies (H2^{*}), f satisfies (H1), and

(2.35)
$$f_* \leq \liminf_{n \to \infty} f_n \text{ and } \limsup_{n \to \infty} f_n \leq f^*.$$

For each n. set

$$v_n(x) = \sup_{\gamma \in \mathscr{A} : \gamma \leq x} \int_0^1 f_n(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt.$$

Then $v_n \rightarrow U$ uniformly where U is the value function given by (1.4).

Proof. We claim that $\{f_n\}_{n=1}^{\infty}$ is a uniformly bounded sequence. To see this, suppose to the contrary that there exists a sequence x_n in $[0,1]^d$ such that $f_n(x_n) \to \infty$ as $n \to \infty$. By passing to a subsequence, if necessary, we may assume that $x_n \to x_0 \in [0,1]^2$ as $n \to \infty$. By the definition of the upper limit and (2.35), we have

$$f^*(x_0) \ge \limsup_{n \to \infty} f_n(x_0) = \infty,$$

which contradicts the assumption that f satisfies (H1) and establishes the claim.

Since ${f_n}_{n=1}^{\infty}$ is uniformly bounded, there exists (by Lemma 2.1) a constant C such that $[v_n]_{\frac{1}{2}} \leq C$ for all *n*. The sequence v_n is therefore bounded and equicontinuous, and by the Arzela-Ascoli theorem there exists a subsequence v_{n_k} and a Hölder-continuous function $v: \mathbb{R}^d \to \mathbb{R}$ such that $v_{n_k} \to v$ uniformly on compact sets in \mathbb{R}^d as $k \to \infty$. By Theorem 2.5 (i), (ii), we conclude that the convergence is actually uniform on \mathbb{R}^d . By Theorem 2.5, each v_n is a Pareto-montone truncatable viscosity solution of

$$v_{n,x_1}\cdots v_{n,x_d} = \frac{1}{d^d}f_n$$
 on \mathbb{R}^d_+

By standard results on viscosity solutions (see [11, Remark 6.3]) and (2.35), we have that v is a Pareto-monotone viscosity solution of

$$v_{x_1}\cdots v_{x_d}=\frac{1}{d^d}f$$
 on \mathbb{R}^d_+ .

By the assumption that $\operatorname{supp}(f_n) \subset [0,1]^d$, we have that $v_n(x) = 0$ for all $x \notin \mathbb{R}^d_+$, hence v(x) = 0 for all $x \notin \mathbb{R}^d_+$.

We claim that v is truncatable. To see this, fix $z \in \mathbb{R}^d_+$ and define \tilde{v} , \tilde{v}_n , \hat{f} and \hat{f}_n as in Remark 2. Since v_n is truncatable, \tilde{v}_n is a viscosity solution of

$$\widetilde{v}_{n,x_1}\cdots\widetilde{v}_{n,x_d}=\frac{1}{d^d}\widehat{f}_n$$
 on \mathbb{R}^d_+

By the definition of \hat{f}_n , we have $\hat{f}_n \leq f_n^*$, with $\hat{f}_n(x) = f_n^*(x)$ for $x \leq z$. It follows that

$$\limsup_{n \to \infty} \widehat{f}_n(x) \le \limsup_{n \to \infty} f_n^*(x) \le f^*(x) = \widehat{f}(x) = (\widehat{f})^*(x) \text{ for } x \le z.$$

For $x \leq z$, there exists a neighborhood V of x on which \hat{f}_n is identically zero for all n. It follows that

$$\limsup_{n\to\infty}^* \widehat{f}_n(x) = 0 = (\widehat{f})^*(x),$$

and therefore $\limsup_{n\to\infty}^* \widehat{f_n} \leq (\widehat{f})^*$. Since $\widetilde{v}_{n_k} \to \widetilde{v}$ uniformly, we can again apply standard results on viscosity solutions [11] to find that \widetilde{v} is a viscosity subsolution of

$$\widetilde{v}_{x_1}\cdots\widetilde{v}_{x_d}=\frac{1}{d^d}\widehat{f}$$
 on $\mathbb{R}^d_+,$

which proves the claim.

By Theorem 2.8 we have v = U on \mathbb{R}^d_+ . Since U(x) = v(x) = 0 for $x \notin \mathbb{R}^d_+$ we have v = U on \mathbb{R}^d . The above argument can be used to show that every subsequence of v_n contains a uniformly convergent subsequence converging to U. It follows that $v_n \to U$ uniformly in \mathbb{R}^d as $n \to \infty$. \Box

We now establish uniqueness of viscosity solutions of (1.5).

THEOREM 2.10. Suppose that Ω satisfies (H2^{*}) and f satisfies (H1). Then there exists a unique Pareto-monotone viscosity solution u of

(2.36)
$$\begin{cases} u_{x_1} \cdots u_{x_d} = f & \text{on } \mathbb{R}^d_+ \\ u = 0 & \text{on } \partial \mathbb{R}^d_+, \end{cases}$$

satisfying the additional boundary condition

(2.37)
$$u(x) = u(\min(x_1, 1), \dots, \min(x_d, 1)) \text{ for } x \in \mathbb{R}^d_+.$$

Proof. By Theorem 2.5 there exists a Hölder-continuous Pareto-monotone viscosity solution u of (2.36). To prove uniqueness, we will show that $u = d \cdot U$, where U is the value function defined by (1.4). Let $\varepsilon > 0$ and consider the inf and sup convolutions of f, defined for $x \in \mathbb{R}^d_+$ by

$$f_{\varepsilon}(x) = \inf_{y \in \mathbb{R}^d_+} \left\{ f(y) + \frac{1}{\varepsilon} |x - y| \right\} \text{ and } f^{\varepsilon}(x) = \sup_{y \in \mathbb{R}^d_+} \left\{ f(y) - \frac{1}{\varepsilon} |x - y| \right\}.$$

Recall that f_{ε} and f^{ε} are Lipschitz continuous with constant $1/\varepsilon$ and $f_{\varepsilon} \leq f \leq f^{\varepsilon}$. Without loss of generality, we may assume that $\overline{\Omega} \subset (0,1)^d$, and hence for $\varepsilon > 0$ small enough, we have $\operatorname{supp}(f_{\varepsilon}), \operatorname{supp}(f^{\varepsilon}) \subset [0,1]^d$. For $x \in \mathbb{R}^d_+$, set

$$U^{\varepsilon}(x) = \sup_{\gamma \in \mathscr{A} : \gamma \leq x} \int_0^1 f^{\varepsilon}(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt,$$

and

$$U_{\varepsilon}(x) = \sup_{\gamma \in \mathscr{A} : \gamma \leq x} \int_0^1 f_{\varepsilon}(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt.$$

By Theorem 2.5, $d \cdot U^{\varepsilon}$ is a viscosity solution of

(2.38)
$$v_{x_1} \cdots v_{x_d} = f^{\varepsilon} \quad \text{on} \quad \mathbb{R}^d_+,$$

and satisfies the boundary condition (2.37). Since $f \leq f^{\varepsilon}$ and u is a viscosity solution of (2.36), we see that u is a viscosity subsolution of (2.38). Since u = U = 0 on $\partial \mathbb{R}^d_+$ and u satisfies (2.37), we can apply Theorem 2.6 to find that $u \leq d \cdot U^{\varepsilon}$. By a similar argument, we have that $u \geq d \cdot U_{\varepsilon}$. Since $f_{\varepsilon}, f^{\varepsilon} \in \mathscr{B}$ and (2.35) is satisfied for the sequences $\{f^{\varepsilon}\}_{\varepsilon>0}$ and $\{f_{\varepsilon}\}_{\varepsilon>0}$, we have by Lemma 2.9 that $U_{\varepsilon}, U^{\varepsilon} \to U$ uniformly in \mathbb{R}^d as $\varepsilon \to 0$, and hence $u = d \cdot U$. \Box

3. Large sample asymptotics of u_n . The proof of Theorem 1.2 is split into several steps. In Section 3.1, we prove a basic convergence result for piecewise constant density functions, which is a generalization of the results of Deuschel and Zeitouni [15]. In Section 3.2, we extend the convergence result to densities that are continuous on Ω and vanish on $\mathbb{R}^d \setminus \Omega$ by considering a sequence of piecewise constant approximations to f, applying the results from Section 3.1, and passing to the limit. This requires a perturbation result for the energy J, which we obtained from the comparison principle for the associated Hamilton-Jacobi PDE (1.5) in Lemma 2.9.

3.1. Piecewise constant densities. We aim to prove a basic convergence result for piecewise constant densities here. The proof is split into a lower bound, Theorem 3.1, and an upper bound, Theorem 3.4. We should note that the techniques used here are similar to those used by Deuschel and Zeitouni [15], who showed the same convergence result for C^1 densities on the unit hypercube in dimension d = 2.

Let us introduce some notation. For a finite set $S \subset \mathbb{R}^d$, let $\ell(S)$ denote the length of a longest increasing chain in *S*. The set function ℓ has an important invariance. If $\Psi : \mathbb{R}^d \to \mathbb{R}^d$ is a mapping that preserves the partial order \leq , i.e., $x \leq y \iff \Psi(x) \leq \Psi(y)$, then

(3.1)
$$\ell(S) = \ell(\Psi(S)) \text{ for any } S \subset \mathbb{R}^d.$$

For $A \subset \mathbb{R}^d$ we denote by $\chi_A : \mathbb{R}^d \to \mathbb{R}$ the characteristic function of the set *A*, which takes the value 1 on *A* and 0 on $\mathbb{R}^d \setminus A$. When *A* is Lebesgue measurable, we denote by |A| the



FIG. 3.1. An illustration of some quantities from the proof of Theorem 3.1.

Lebesgue measure of *A*. We set $\mathbf{0}_d = (0, \dots, 0) \in \mathbb{R}^d$ and $\mathbf{1}_d = (1, \dots, 1) \in \mathbb{R}^d$. Given an integer *L*, we partition $[0, 1)^d$ into L^d hypercubes of side length 1/L. More precisely, for a multiindex $\alpha \in \mathbb{N}^L$ with $\|\alpha\|_{\infty} \leq L$, where $\|\alpha\|_{\infty} = \max(\alpha_1, \dots, \alpha_L)$, we set

$$(3.2) Q_{L,\alpha} = \{ x \in [0,1)^d : \alpha - \mathbf{1}_d \leq Lx < \alpha \}.$$

We say that $f : [0,1)^d \to [0,\infty)$ is *L*-piecewise constant if f is constant on $Q_{L,\alpha}$ for all α . If f is *L*-piecewise constant then f is *kL*-piecewise constant for all $k \in \mathbb{N}$. For convenience, we also set

$$\overline{J} = \sup_{\gamma \in \mathscr{A}} J(\gamma) = \sup_{\gamma \in \mathscr{A}} \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt.$$

We now establish an asymptotic lower bound on $\ell(\{X_1, \ldots, X_n\})$.

THEOREM 3.1. Let $f : [0,1)^d \to [0,\infty)$ be L-piecewise constant, and let X_1, \ldots, X_n be i.i.d. with density f. Then

$$\liminf_{n\to\infty} n^{-\frac{1}{d}}\ell\left(\{X_1,\ldots,X_n\}\right) \ge c_d \overline{J} \quad \text{a.s.}$$

Proof. Let $\varepsilon > 0$ and select $\gamma \in \mathscr{A}$ with $J(\gamma) \ge \overline{J} - \frac{\varepsilon}{c_d}$. Without loss of generality, we may assume that $\gamma'(t) > 0$ for all $t \in [0, 1]$. Let s_1, \ldots, s_k denote the $k \le dL$ times at which γ intersects the set

$$\bigcup_{\alpha} \partial Q_{L,\alpha} \cap (0,1)^d.$$

Set $s_0 = 0$ and $s_{k+1} = 1$. For j = 0, ..., k set $I_j = [s_j, s_{j+1})$ and

(3.3)
$$R_j = \{ x \in [0,1)^d : \gamma(s_j) \leq x < \gamma(s_{j+1}) \}.$$

For every *j* we have $R_j \subset Q_{L,\alpha}$ for some α . Recalling the definition of *J* (1.3) we have

(3.4)
$$J(\gamma) = \sum_{j=0}^{k} \int_{I_j} f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt = \sum_{j=0}^{k} f(\gamma(s_j))^{\frac{1}{d}} \int_{I_j} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt,$$

where the second equality follows from the fact that f is constant on $R_j \subset Q_{L,\alpha}$. Applying the generalized Hölder inequality to (3.4) we have

$$J(\gamma) \leq \sum_{j=0}^{k} f(\gamma(s_j))^{\frac{1}{d}} \prod_{i=1}^{d} \left(\int_{I_j} \gamma'_i(t) dt \right)^{\frac{1}{d}} = \sum_{j=0}^{k} f(\gamma(s_j))^{\frac{1}{d}} |R_j|^{\frac{1}{d}}.$$

Setting $p_j = \int_{R_j} f(x) dx = f(\gamma(s_j)) |R_j|$ we have

(3.5)
$$\overline{J} \leq J(\gamma) + \frac{\varepsilon}{c_d} \leq \sum_{j=0}^k p_j^{\frac{1}{d}} + \frac{\varepsilon}{c_d}.$$

Fix $j \in \{0, ..., k\}$. Let n_j denote the number of points from $X_1, ..., X_n$ falling in R_j and set

(3.6)
$$\ell_j(n) = \ell(\{X_1, \dots, X_n\} \cap R_j).$$

Then n_j is Binomially distributed with parameters n and p_j . If f is identically zero on R_j then $\ell_j(n) = 0$ with probability one for all n, and $p_j = 0$, hence

$$(3.7) n^{-\frac{1}{d}}\ell_j(n) = c_d p_j^{\frac{1}{d}} \quad \text{a.s.}$$

If *f* is not identically zero on R_j , then since $\gamma'(t) > 0$ for all *t*, we have $|R_j| > 0$ and hence $p_j > 0$. The conditional law $\rho_j := p_j^{-1} \cdot f \cdot \chi_{R_j}$ is then uniform on R_j . Let i_1, \ldots, i_{n_j} be the indices of the n_j random variables out of X_1, \ldots, X_n that belong to R_j . Let $\Psi : R_j \to [0, 1)^d$ be the injective affine transformation mapping R_j onto $[0, 1)^d$. Then $\Psi(X_{i_1}), \ldots, \Psi(X_{i_{n_j}})$ are independent and uniformly distributed on $[0, 1)^d$. By [6, Remark 1], we have

$$n_j^{-\frac{1}{d}}\ell\left(\{\Psi(X_{i_1}),\ldots,\Psi(X_{i_n})\}\right)\to c_d$$
 a.s.

Since Ψ preserves the partial order \leq , we have by (3.1) that

$$\ell_j(n) = \ell\left(\{X_{i_1},\ldots,X_{i_{n_j}}\}\right) = \ell\left(\{\Psi(X_{i_1}),\ldots,\Psi(X_{i_{n_j}})\}\right).$$

Since $n^{-1}n_i \rightarrow p_i$ almost surely we have

(3.8)
$$n^{-\frac{1}{d}}\ell_j(n) = n_j^{-\frac{1}{d}}(n^{-1}n_j)^{\frac{1}{d}}\ell_j(n) \to c_d p_j^{\frac{1}{d}} \quad \text{a.s.}$$

Combining this (3.5), (3.7) and (3.8), we see that

(3.9)
$$n^{-\frac{1}{d}} \sum_{j=0}^{k} \ell_j(n) \to c_d \sum_{j=0}^{k} p_j^{\frac{1}{d}} \ge c_d \overline{J} - \varepsilon \quad \text{a.s.}$$

Since γ is a monotone curve (i.e., $\gamma'(t) \ge 0$), we can connect longest chains from each rectangle R_j together to form a chain in $[0, 1)^d$. It follows that

(3.10)
$$\ell(\{X_1, \dots, X_n\}) \ge \sum_{j=0}^k \ell_j(n).$$

Combining this with (3.9) we have

$$\liminf_{n\to\infty} n^{-\frac{1}{d}} \ell\left(\{X_1,\ldots,X_n\}\right) \ge c_d \overline{J} - \varepsilon \quad \text{a.s.},$$

which completes the proof. \Box

For the proof of the upper bound, we need to introduce some new notation. Let k_1 be an integer and set $\Delta x = 1/k_1$. Let k_2 be another integer and set $\Delta y = \Delta x/k_2$. For given k_1, k_2 , we say that a sequence of multiindices $\mathbf{b} = (b_j)_{j=1}^{k_1} \subset \mathbb{N}^{d-1}$ is *admissible* if $b_1 \leq \cdots \leq b_{k_1}$ and $||b_j||_{\infty} \leq k_1k_2$ for all *j*. We denote the set of admissible multiindices by $\Phi(k_1, k_2)$. For $\mathbf{b} \in \Phi(k_1, k_2)$, define $z_{\mathbf{b},0}, z_{\mathbf{b},1}, \dots, z_{\mathbf{b},k_1}$ in $[0, 1]^d$ by $z_{\mathbf{b},0} = \mathbf{0}_d$ and $z_{\mathbf{b},j} = (b_j \Delta y, j \Delta x)$ for $j \geq 1$. Since **b** is admissible, $z_{\mathbf{b},0}, \dots, z_{\mathbf{b},k_1}$ defines a chain in $[0, 1]^d$. Define $\gamma_{\mathbf{b}} : [0, 1] \to [0, 1]^d$ to be the polygonal curve connecting the points $z_{\mathbf{b},0}, \dots, z_{\mathbf{b},k_1}$, i.e.,

$$\gamma_{\mathbf{b}}(t) = z_{\mathbf{b},j-1} + \frac{1}{\Delta x} (z_{\mathbf{b},j} - z_{\mathbf{b},j-1})(t - (j-1)\Delta x)$$

for $t \in [(j-1)\Delta x, j\Delta x]$. For $\mathbf{b} \in \Phi(k_1, k_2)$ and $1 \le j \le k_1$, set

$$R_{\mathbf{b},j} = \left\{ x \in [0,1)^d : z_{\mathbf{b},j-1} - (\mathbf{1}_{d-1},0) \Delta y \leq x < z_{\mathbf{b},j} \right\}.$$

For each rectangle $R_{\mathbf{b},j}$, we set $p_{\mathbf{b},j} = \int_{R_{\mathbf{b},j}} f(x) dx$. We say that a chain $x_1 \leq x_2 \leq \cdots \leq x_m$ in $[0,1)^d$ is **b**-increasing if

$$\{x_1,\ldots,x_m\}\subset \bigcup_{j=1}^{k_1}R_{\mathbf{b},j}$$

It is not hard to see that for any k_1, k_2 , every chain in $[0,1)^d$ is **b**-increasing for some $\mathbf{b} \in \Phi(k_1, k_2)$. See Figure 3.1 for an illustration of the above definitions.

We first need a preliminary lemma which bounds the length of a longest chain within the narrow strip

(3.11)
$$T_j := [0,1]^{d-1} \times [(j-1)\Delta x, j\Delta x),$$

for any $j \in \{1, ..., k_1\}$. We note that the following lemma is a generalization of [15, Lemma 7]. The proof is based on the same idea of using a mixing process to embed $X_1, ..., X_n$ into another set of *i.i.d.* random variables that are uniform when restricted to the strip T_j .

LEMMA 3.2. Let $f : [0,1)^d \to [0,\infty)$ be L-piecewise constant, and let X_1, \ldots, X_n be i.i.d. with density f. Fix an integer $j \in \{1, \ldots, k_1\}$ and let $0 < \Delta x \le ||f||_{L^{\infty}((0,1)^d)}^{-1}$. Then

(3.12)
$$\limsup_{n \to \infty} n^{-\frac{1}{d}} \ell(\{X_1, \dots, X_n\} \cap T_j) \le c_d \left(2\Delta x \|f\|_{L^{\infty}((0,1)^d)}\right)^{\frac{1}{d}} \text{ a.s}$$

Proof. Set $M = ||f||_{L^{\infty}((0,1)^d)}$ and let $g = f + (M - f) \cdot \chi_{T_j}$. Let Y_1, \ldots, Y_n be *i.i.d.* according to the conditional density $\beta^{-1}(M - f) \cdot \chi_{T_j}$ where $\beta = \int_{T_j} M - f(x) dx$. Let m_1, \ldots, m_n be Bernoulli zero-one random variables with parameter $(1 + \beta)^{-1}$ and set

Define Z_1, \ldots, Z_n through the mixture process

$$Z_k = m_k X_{i_k} + (1 - m_k) Y_k$$



FIG. 3.2. An illustration of the quantities $\mathbf{R}_{\mathbf{b},j}$, $z_{\mathbf{b},j}$, $Q_{L,\alpha}$ and $\gamma_{\mathbf{b}}$ in two dimensions with $\mathbf{b} = (b_1, b_2, b_3, b_4) = (7,7,10,11)$. In this case, the unit square is partitioned into four squares, $Q_{L,(1,1)}$, $Q_{L,(1,2)}$, $Q_{L,(2,1)}$ and $Q_{L,(2,2)}$, which are separated by dotted lines in the figure.

Then Z_1, \ldots, Z_n are *i.i.d.* with density $(1 + \beta)^{-1}g$. Let *W* denote the cardinality of the set $\{Z_1, \ldots, Z_n\} \cap T_j$. Then *W* is binomially distributed with parameters *n* and $p := (1 + \beta)^{-1}\Delta xM$. Since *g* is constant on T_j , we can use a similar argument to that in Theorem 3.1 to show that

(3.14)
$$n^{-\frac{1}{d}}\ell(\{Z_1,\ldots,Z_n\}\cap T_j)\to c_d p^{\frac{1}{d}} \quad \text{a.s.}$$

Let $m = i_n$ and note that

$$\ell(\{X_1,...,X_m\}) = \ell(\{Z_k : m_k = 1\} \cap T_j) \le \ell(\{Z_1,...,Z_n\} \cap T_j),$$

and that $p \leq \Delta x M$. Combining this with (3.14) we have

(3.15)
$$\limsup_{n \to \infty} n^{-\frac{1}{d}} \ell\left(\{X_1, \dots, X_m\}\right) \le c_d (\Delta x M)^{\frac{1}{d}} \quad \text{a.s}$$

Since *m* is Binomially distributed with parameters *n* and $(1 + \beta)^{-1}$, we have $nm^{-1} \rightarrow 1 + \beta$ almost surely and hence

$$\limsup_{n \to \infty} m^{-\frac{1}{d}} \ell(\{X_1, \dots, X_m\}) = \limsup_{n \to \infty} (nm^{-1})^{\frac{1}{d}} n^{-\frac{1}{d}} \ell(\{X_1, \dots, X_m\})$$
$$\leq (1+\beta)^{\frac{1}{d}} c_d(\Delta x M)^{\frac{1}{d}} \quad \text{a.s.}$$

Since $\beta \leq \Delta x M \leq 1$ we have

$$\limsup_{n\to\infty} m^{-\frac{1}{d}}\ell\left(\{X_1,\ldots,X_m\}\right) \le c_d(2\Delta x M)^{\frac{1}{d}} \quad \text{a.s.}$$

The desired result (3.12) follows from noting that $n \mapsto m(n)$ is monotone nondecreasing along every sample path and $m \to \infty$ as $n \to \infty$ with probability one. \Box

The following short technical lemma is essential in the proof of Theorem 3.4

LEMMA 3.3. Let $f: [0,1)^d \to [0,\infty)$ be L-piecewise constant. For every $\varepsilon > 0$ and $k_1 \ge L$ we have

(3.16)
$$\sum_{j \in \mathscr{H}_{\mathbf{b}}} p_{\mathbf{b},j}^{\frac{1}{d}} \leq \overline{J} + \varepsilon.$$

for all $\mathbf{b} \in \Phi(k_1, k_2)$, the admissible multiindices, and $k_2 \ge C \|f\|_{L^{\infty}((0,1)^d)} k_1^{d-1} / \varepsilon^d$, where

(3.17)
$$\mathscr{H}_{\mathbf{b}} = \{ j : R_{\mathbf{b},j} \subset Q_{L,\alpha} \text{ for some } \alpha \}.$$

Proof. Let $k_1, k_2, \varepsilon > 0$, and $\mathbf{b} \in \Phi(k_1, k_2)$. Set $I_j = [(j-1)\Delta x, j\Delta x)$ and fix $j \in \{1, \dots, k_1\}$ and $t \in I_i$. Note that

$$|R_{\mathbf{b},j}| = \begin{cases} \Delta y^{d-1} \Delta x \prod_{i=1}^{d-1} (b_{j,i} - b_{j-1,i} + 1) & \text{if } j \ge 2\\ \Delta y^{d-1} \Delta x \prod_{i=1}^{d-1} (b_{j,i} - b_{j-1,i}) & \text{if } j = 1, \end{cases}$$

and

$$\Delta x^d \gamma_{\mathbf{b},1}'(t) \cdots \gamma_{\mathbf{b},d}'(t) = \Delta y^{d-1} \Delta x \prod_{i=1}^{d-1} (b_{j,i} - b_{j-1,i}),$$

where we set $\mathbf{b}_0 = 0$ for convenience. A short computation shows that

(3.18)
$$|\Delta x(\gamma'_{\mathbf{b},1}(t)\cdots\gamma'_{\mathbf{b},d}(t))^{\frac{1}{d}}-|R_{\mathbf{b},j}|^{\frac{1}{d}}| \leq C\Delta x^{\frac{1}{d}}\Delta y^{\frac{1}{d}},$$

where $C = (d-1)^{\frac{1}{d}}$. Since *f* is *L*-piecewise constant we have

(3.19)
$$f(\boldsymbol{\gamma}_{\mathbf{b}}(t)) = f(z_{\mathbf{b},j-1}) = \frac{p_{\mathbf{b},j}}{|R_{\mathbf{b},j}|},$$

for all $j \in \mathscr{H}_{\mathbf{b}}$ and $t \in I_j$. Noting that $\Delta x = |I_j|$ and recalling the definition of J (1.3) we have

$$\begin{split} \bar{J} &\geq J(\gamma_{\mathbf{b}}) \stackrel{(3.18)}{\geq} \sum_{j \in \mathscr{H}_{\mathbf{b}}} \frac{1}{|I_{j}|} \int_{I_{j}} f(\gamma_{\mathbf{b}}(t))^{\frac{1}{d}} (|R_{\mathbf{b},j}|^{\frac{1}{d}} - C\Delta x^{\frac{1}{d}} \Delta y^{\frac{1}{d}}) dt \\ &= \sum_{j \in \mathscr{H}_{\mathbf{b}}} \frac{|R_{\mathbf{b},j}|^{\frac{1}{d}}}{|I_{j}|} \int_{I_{j}} f(\gamma_{\mathbf{b}}(t))^{\frac{1}{d}} dt - C \sum_{j \in H_{\mathbf{b}}} \frac{1}{|I_{j}|} \int_{I_{j}} f(\gamma_{\mathbf{b}}(t))^{\frac{1}{d}} \Delta x^{\frac{1}{d}} \Delta y^{\frac{1}{d}} dt \\ 0) \qquad \stackrel{(3.19)}{\geq} \sum_{j \in \mathscr{H}_{\mathbf{b}}} p_{\mathbf{b},j}^{\frac{1}{d}} - ||f||_{L^{\infty}((0,1)^{d})}^{\frac{1}{d}} k_{1}^{\frac{d-1}{d}} k_{2}^{-\frac{1}{d}}. \end{split}$$

(3.2

Taking $k_2 \ge (C/\varepsilon)^d ||f||_{L^{\infty}((0,1)^d)} k_1^{d-1}$ completes the proof. \Box We now establish an asymptotic upper bound on $\ell(\{X_1,\ldots,X_n\})$.

THEOREM 3.4. Let $f : [0,1)^d \to [0,\infty)$ be L-piecewise constant, and let X_1, \ldots, X_n be i.i.d. with density f. Then

$$\limsup_{n\to\infty} n^{-\frac{1}{d}}\ell(\{X_1,\ldots,X_n\}) \leq c_d \overline{J} \quad \text{a.s.}$$

Proof. Let $k_2 > 0, k_1 \ge L, \varepsilon > 0$, and $\mathbf{b} \in \Phi(k_1, k_2)$. We suppose that $k_1 \ge L$ is a multiple of *L* so that *f* is k_1 -piecewise constant. Let $\ell_{\mathbf{b}}(n)$ denote the length of a longest **b**-increasing chain. Let n_j denote the number of X_1, \ldots, X_n that belong $R_{\mathbf{b},j}$ and set

(3.21)
$$\ell_{\mathbf{b},j}(n) = \ell\left(\{X_1,\ldots,X_n\} \cap R_{\mathbf{b},j}\right).$$

Due to the monotonicity of $z_{\mathbf{b},0}, \ldots, z_{\mathbf{b},k_1}$, at most (d-1)L of $R_{\mathbf{b},1}, \ldots, R_{\mathbf{b},k_1}$ can have a nonempty intersection with more than one hypercube $Q_{L,\alpha}$. It follows that $|\mathscr{H}^c_{\mathbf{b}}| \leq (d-1)L$, where $\mathscr{H}^c_{\mathbf{b}} = \{1, \ldots, k_1\} \setminus \mathscr{H}_{\mathbf{b}}$.

Since each **b**-increasing chain is the union of chains in $R_{\mathbf{b},1}, \ldots, R_{\mathbf{b},k_1}$, we have

(3.22)
$$\ell_{\mathbf{b}}(n) \leq \sum_{j=1}^{k_1} \ell_{\mathbf{b},j}(n) = \sum_{j \in \mathscr{H}_{\mathbf{b}}^c} \ell_{\mathbf{b},j}(n) + \sum_{j \in \mathscr{H}_{\mathbf{b}}} \ell_{\mathbf{b},j}(n).$$

We will deal with each of the above sums separately. For the first term, set $M = ||f||_{L^{\infty}((0,1)^d)}$ and let k_1 be large enough so that $\Delta x \leq 1/M$. Since $R_{\mathbf{b},j} \subset T_j$ for each j, we have by Lemma 3.2 that

$$\limsup_{n \to \infty} n^{-\frac{1}{d}} \sum_{j \in \mathscr{H}_{\mathbf{b}}^{c}} \ell_{\mathbf{b},j}(n) \le c_{d} |\mathscr{H}_{\mathbf{b}}^{c}| (2\Delta x M)^{\frac{1}{d}} \le c_{d} (d-1) L (2M)^{\frac{1}{d}} k_{1}^{-\frac{1}{d}} \quad \text{a.s.}$$

Choose k_1 large enough so that

(3.23)
$$\limsup_{n \to \infty} n^{-\frac{1}{d}} \sum_{j \in \mathscr{H}_{\mathbf{b}}^{c}} \ell_{\mathbf{b},j}(n) \leq \frac{\varepsilon}{2} \quad \text{a.s}$$

We now bound the second sum in (3.22). By Lemma 3.3, choose $k_2 = k(M, k_1, \varepsilon)$ so that

(3.24)
$$\sum_{j \in \mathscr{H}_{\mathbf{b}}} p_{\mathbf{b},j}^{\frac{1}{d}} \leq \overline{J} + \frac{\varepsilon}{2c_d},$$

for all $\mathbf{b} \in \Phi(k_1, k_2)$. For any $j \in \mathscr{H}_{\mathbf{b}}$, the conditional density ρ_j on $R_{\mathbf{b},j}$ is uniform. By a similar argument as in the proof of Theorem 3.1, we have that

$$n^{-\frac{1}{d}}\ell_{\mathbf{b},j}(n) \to c_d p_{\mathbf{b},j}^{\frac{1}{d}}$$
 a.s.

Combining this with (3.22), (3.23), and (3.24) we have

(3.25)
$$\limsup_{n \to \infty} n^{-\frac{1}{d}} \ell_{\mathbf{b}}(n) \le c_d \overline{J} + \varepsilon \quad \text{a.s.}$$

Since every chain in $X_1, ..., X_n$ is **b**-increasing for some $\mathbf{b} \in \Phi(k_1, k_2)$, we have

$$\ell(\{X_1,\ldots,X_n\}) \leq \max_{\mathbf{b}\in\Phi(k_1,k_2)}\ell_{\mathbf{b}}(n),$$

for every *n*. It follows that

$$\limsup_{n\to\infty} \ell(\{X_1,\ldots,X_n\}) \le c_d \overline{J} + \varepsilon \quad \text{a.s.},$$

which completes the proof. \Box

3.2. Continuous densities on Ω . We now generalize the convergence results on piecewise constant densities, Theorems 3.1 and 3.4, to continuous densities on Ω . Our main result, Theorem 1.2, is proved at the end of the section. The idea of our approach is to divide $[0,1)^d$ into a large number of hypercubes, and to flatten f on each sub-cube. We can then apply the results from Section 3.1 and take the limit as the size of the sub-cubes tends to zero. In order to pass to the limit, we apply the perturbation result given in Lemma 2.9.

Let X_1, \ldots, X_n be *i.i.d.* with density f. We recall that $u_n(x)$ denotes the length of a longest chain among X_1, \ldots, X_n consisting of points less than or equal to x under the partial order \leq . In other words

$$u_n(x) = \ell\left(\{X_i : X_i \leq x\}\right)$$

We also recall the definition of the value function U, defined in (1.4) by

$$U(x) = \sup_{\gamma \in \mathscr{A}: \gamma \leq x} \int_0^1 f(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt$$

We now establish pointwise asymptotic upper and lower bounds on u_n . THEOREM 3.5. Let $f : \mathbb{R}^d \to \mathbb{R}$ satisfy (H1) and let $\Omega \subset \mathbb{R}^d_+$ satisfy (H2). Then for every $z \in \mathbb{R}^d$ we have

(3.26)
$$\limsup_{n \to \infty} n^{-\frac{1}{d}} u_n(z) \le c_d U(z) \quad \text{a.s.}$$

Proof. Set $D = \{x \in \mathbb{R}^d : 0 \leq x < z\}$ and $p = \int_D f(x) dx$. Suppose that p = 0. It follows from (H1) that f is lower semicontinuous, and hence f(x) = 0 for $x \leq z$. Thus $u_n(z) = 0 =$ U(z) almost surely.

Suppose that p > 0 and let $\varepsilon > 0$. Let $k \in \mathbb{N}$ and partition D into k^d hypercubes $Q_{k,\alpha}$ for multiindices α with $\|\alpha\|_{\infty} \leq k$. Define $f_k : \mathbb{R}^d \to [0, \infty)$ by

(3.27)
$$f_k(x) = \sum_{\alpha} \left(\sup_{Q_{k,\alpha}} f \right) \chi_{Q_{k,\alpha}}(x) + f(x) \chi_{\mathbb{R}^d \setminus D}(x),$$

and set $p_k = \int_D f_k(x) dx$. For every integer k, f_k is k-piecewise constant on D and $f \leq f_k$. Define $v_k : \mathbb{R}^d \to \mathbb{R}$ by

(3.28)
$$v_k(x) = \sup_{\gamma \in \mathscr{A} : \gamma \leq x} \int_0^1 f_k(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt.$$

Note that the sequence f_k is uniformly bounded, Borel-measurable, and has compact support in $[0,1]^d$. Furthermore, it follows from (3.27) that (2.35) holds for the sequence f_k . Hence by Lemma 2.9 we have that $v_k \rightarrow U$ uniformly as $k \rightarrow \infty$. Now fix k large enough so that

$$(3.29) |v_k(z) - U(z)| \le \frac{\varepsilon}{c_d}.$$

Set

(3.30)
$$\lambda = \left(\int_{\mathbb{R}^d} f_k(x) \, dx\right)^{-1},$$

and define $g = \lambda f_k$. Then $\lambda f \leq g$ and we can write g as a convex combination of two distributions as follows:

$$g = \lambda f + (g - \lambda f)$$

Let Y_1, \ldots, Y_n be *i.i.d.* with density $(1 - \lambda)^{-1}(g - \lambda f)$, let m_1, \ldots, m_n be Bernoulli random variables with parameter λ , and set

$$i_j = m_1 + \cdots + m_j$$
.

Define

$$Z_j = m_j X_{i_j} + (1 - m_j) Y_j$$

Then a simple computation shows that Z_1, \ldots, Z_n are *i.i.d.* with density g. Let W denote the cardinality of $\{Z_1, \ldots, Z_n\} \cap D$. Since g is k-piecewise constant on D, we can apply Theorems 3.1 and 3.4 to obtain

(3.31)
$$\lim_{n \to \infty} W^{-\frac{1}{d}} \ell\left(\{Z_1, \dots, Z_n\} \cap D\}\right) = c_d p_k^{-\frac{1}{d}} v_k(z) \quad \text{a.s.}$$

Note that *W* is Binomially distributed with parameters *n* and λp_k , hence $n^{-1}W \rightarrow \lambda p_k$ almost surely. Applying this to (3.31) we have

(3.32)
$$\lim_{n \to \infty} n^{-\frac{1}{d}} \ell(\{Z_1, \dots, Z_n\} \cap D) = \lim_{n \to \infty} (n^{-1}W)^{\frac{1}{d}} W^{-\frac{1}{d}} \ell(\{Z_1, \dots, Z_n\} \cap D) = c_d \lambda^{\frac{1}{d}} v_k(z) \quad \text{a.s.}$$

Set $m = i_n$. Note that m is Binomially distributed with parameters n and λ , and

(3.33)
$$u_m(z) = \ell(\{X_1, \dots, X_m\} \cap D) \le \ell(\{Z_1, \dots, Z_n\} \cap D).$$

Combining (3.33) with (3.32) and the fact that $n^{-1}m \rightarrow \lambda$ as $n \rightarrow \infty$ we have

(3.34)
$$\limsup_{n \to \infty} m^{-\frac{1}{d}} u_m(z) \stackrel{(3.33)}{\leq} \lim_{n \to \infty} (m^{-1}n)^{\frac{1}{d}} n^{-\frac{1}{d}} \ell\left(\{Z_1, \dots, Z_n\} \cap D\right) \stackrel{(3.32)}{=} c_d v_k(z) \quad \text{a.s.}$$

Recalling (3.29) we have

$$\limsup_{n\to\infty} m^{-\frac{1}{d}}u_m(z) \le c_d U(z) + \varepsilon \quad \text{a.s.}$$

As in the proof of Lemma 3.2, the proof is completed by noting that $n \mapsto m(n)$ is monotone nondecreasing along every sample path and $m \to \infty$ as $n \to \infty$ with probability one. \Box

THEOREM 3.6. Let $f : \mathbb{R}^d \to \mathbb{R}$ satisfy (H1) and let $\Omega \subset \mathbb{R}^d_+$ satisfy (H2). Then for every $z \in \mathbb{R}^d$ we have

(3.35)
$$\liminf_{n \to \infty} n^{-\frac{1}{d}} u_n(z) \ge c_d U(z) \quad \text{a.s.}$$

Proof. Let $\varepsilon > 0$. As in the proof of Theorem 3.5, we set $D := \{x \in \mathbb{R}^d : 0 \leq x < z\}$ and we may suppose that $p := \int_D f(x) dx > 0$. As before, let $k \in \mathbb{N}$ and partition D into k^d hypercubes $Q_{k,\alpha}$ for multiindices α with $\|\alpha\|_{\infty} \leq k$. Define $f_k : \mathbb{R}^d \to [0,\infty)$ by

(3.36)
$$f_k(x) = \sum_{\alpha} \left(\inf_{\mathcal{Q}_{k,\alpha}} f \right) \chi_{\mathcal{Q}_{k,\alpha}}(x) + f(x) \chi_{\mathbb{R}^d \setminus D}(x),$$

28

and set $p_k = \int_D f_k(x) dx$. Define

(3.37)
$$q(x) = \begin{cases} \frac{f_k(x)}{f(x)}, & \text{if } f(x) > 0\\ 0, & \text{otherwise.} \end{cases}$$

For any α such that $Q_{k,\alpha} \subset \Omega$, we have by (H1) and (3.36) that

$$f_k(x) \ge f(x) - m\left(\frac{\sqrt{d}}{k}\right)$$
 for $x \in Q_{k,\alpha}$

It follows that $||q||_{L^{\infty}(\mathbb{R}^d)} \nearrow 1$ as $k \to \infty$. As in the proof of Theorem 3.5, we have that $v_k \to U$ uniformly as $k \to \infty$, where v_k is defined by (3.28). We can therefore fix k large enough so that

(3.38)
$$c_d \|q\|_{L^{\infty}(\mathbb{R}^d)}^{-\frac{1}{d}} v_k(z) \ge c_d U(z) - \varepsilon.$$

For i = 1, ..., n, let m_i be a Bernoulli zero-one random variable with parameter $||q||_{L^{\infty}(\mathbb{R}^d)}^{-1}q(X_i)$. Let $m = m_1 + \cdots + m_n$ and let $i_1, ..., i_m$ denote the indices for which $m_i = 1$. We claim that $X_{i_1}, ..., X_{i_m}$ are *i.i.d.* with density $g := \lambda f_k$ where λ is defined in (3.30). To see this, first note since f(x) = 0 implies $f_k(x) = 0$, we have $q(x)f(x) = f_k(x)$ for all $x \in \mathbb{R}^d$. Thus

(3.39)
$$P(m_i = 1) = \int_{\mathbb{R}^d} P(m_i = 1 | X_i = x) f(x) \, dx = \int_{\mathbb{R}^d} \frac{q(x)}{\|q\|} f(x) \, dx = \frac{1}{\lambda \|q\|},$$

where $||q|| = ||q||_{L^{\infty}(\mathbb{R}^d)}$. Let $j \ge 1$ and let $A \subset \mathbb{R}^d$ be measurable. We have

$$P(X_{i_j} \in A) = P(X_i \in A \mid m_i = 1)$$

= $\frac{P(X_i \in A \text{ and } m_i = 1)}{P(m_i = 1)}$
 $\stackrel{(3.39)}{=} \lambda ||q|| \int_A P(m_i = 1 \mid X_i = x) f(x) dx$
= $\lambda ||q|| \int_A \frac{q(x)}{||q||} f(x) dx$
= $\int_A \lambda f_k(x) dx.$

By the construction of X_{i_1}, \ldots, X_{i_m} , they are independent random variables, hence the claim is established.

Let *W* denote the cardinality of $\{X_{i_1}, \ldots, X_{i_m}\} \cap D$. By Theorems 3.1 and 3.4, we have

(3.40)
$$\lim_{n \to \infty} W^{-\frac{1}{d}} \ell\left(\{X_{i_1}, \dots, X_{i_m}\} \cap D\right) = c_d p_k^{-\frac{1}{d}} v_k(z) \quad \text{a.s.}$$

Define

$$w_i = \begin{cases} 1, & \text{if } m_i = 1 \text{ and } X_i \in D \\ 0, & \text{otherwise.} \end{cases}$$

Then $W = w_1 + \cdots + w_n$. Each w_i is a Bernoulli zero-one random variable with parameter

$$P(w_i = 1) = P(m_i = 1 \text{ and } X_i \in D) = \int_D P(m_i = 1 | X_i = x) f(x) dx = \frac{p_k}{\|q\|}.$$

It follows that *W* is Binomially distributed with parameters *n* and $||q||^{-1}p_k$, and hence $n^{-1}W \rightarrow ||q||^{-1}p_k$ almost surely. Combining this with (3.40) yields

(3.41)
$$\lim_{n \to \infty} n^{-\frac{1}{d}} \ell\left(\{X_{i_1}, \dots, X_{i_m}\} \cap D\right) = \lim_{n \to \infty} (n^{-1}W)^{\frac{1}{d}} W^{-\frac{1}{d}} \ell\left(\{X_{i_1}, \dots, X_{i_m}\} \cap D\right) = c_d \|q\|^{-\frac{1}{d}} v_k(z) \quad \text{a.s.}$$

Noting that

$$u_n(z) = \ell(\{X_1, \ldots, X_n\} \cap D) \ge \ell(\{X_{i_1}, \ldots, X_{i_m}\} \cap D),$$

we have

(3.42)
$$\liminf_{n \to \infty} n^{-\frac{1}{d}} u_n(z) \ge \lim_{n \to \infty} n^{-\frac{1}{d}} \ell\left(\{X_{i_1}, \dots, X_{i_m}\} \cap D\right)$$
$$\overset{(3.41)}{=} c_d \|q\|^{-\frac{1}{d}} v_k(z) \quad \text{a.s.}$$

Recalling (3.38) we have

$$\liminf_{n\to\infty} n^{-\frac{1}{d}} u_n(z) \ge c_d U(z) - \varepsilon \quad \text{a.s.},$$

which completes the proof. \Box

We now have the proof of Theorem 1.2.

Proof. Let $\varepsilon > 0$. Let $k \in \mathbb{N}$ and for a multiindex $\alpha \in \mathbb{Z}^d$, set $x_{\alpha} = \alpha/k$. Since U is uniformly continuous (by Lemma 2.1) we can choose k large enough so that

$$|U(x_{\alpha+\mathbf{1}_d}) - U(x_\alpha)| \le \frac{\varepsilon}{c_d}$$

for all $\alpha \in \mathbb{Z}^d$. Let *I* be the set of multiindices α for which $x_{\alpha} \in [0,1]^d$. Note that the cardinality of *I* is $(k+1)^d$. Since *I* is finite with cardinality independent of *n*, Theorems 3.5 and 3.6 yield

(3.44)
$$\lim_{n\to\infty}\sup_{\alpha\in I}|n^{-\frac{1}{d}}u_n(x_\alpha)-c_dU(x_\alpha)|=0 \quad \text{a.s.}$$

Let $z \in (0,1]^d$. Then there exists $\alpha \in I$ such that $x_{\alpha} < z \leq x_{\alpha+1_d}$. By the Pareto-monotonicity of u_n and (3.43) we have

$$n^{-\frac{1}{d}}u_{n}(z) - c_{d}U(z) \le n^{-\frac{1}{d}}u_{n}(x_{\alpha+1_{d}}) - c_{d}U(z) \stackrel{(3.43)}{\le} n^{-\frac{1}{d}}u_{n}(x_{\alpha+1_{d}}) - c_{d}U(x_{\alpha+1_{d}}) + \varepsilon$$

By a similar argument, we have

$$n^{-\frac{1}{d}}u_n(z)-c_dU(z)\geq n^{-\frac{1}{d}}u_n(x_\alpha)-c_dU(x_\alpha)-\varepsilon,$$

and hence

(3.45)
$$\|n^{-\frac{1}{d}}u_n - c_d U\|_{L^{\infty}((0,1)^d)} \leq \sup_{\alpha \in I} |n^{-\frac{1}{d}}u_n(x_{\alpha}) - c_d U(x_{\alpha})| + \varepsilon.$$

Combining (3.44) and (3.45) we have

$$\limsup_{n\to\infty} \|n^{-\frac{1}{d}}u_n - c_d U\|_{L^{\infty}((0,1)^d)} \leq \varepsilon \quad \text{a.s.},$$

and hence $\lim_{n\to\infty} \|n^{-\frac{1}{d}}u_n - c_d U\|_{L^{\infty}((0,1)^d)} = 0$ almost surely. The desired result now follows immediately from the boundary conditions on *U* proved in Theorem 2.5 (i), (ii) and the fact that there are almost surely no samples in $\mathbb{R}^d \setminus (0,1)^d$. \Box

As a straightforward application of Theorem 1.2, we can show that non-dominated sorting is stable under bounded random perturbations in the samples X_1, \ldots, X_n . For $\delta > 0$, we set

$$Z_i = X_i + Y_i \delta$$
 for $i = 1, \ldots, n$,

where Y_1, \ldots, Y_n are *i.i.d.* with a continuous compactly supported density function $g : \mathbb{R}^d \to \mathbb{R}$. For $x \in \mathbb{R}^d$, set

$$u_n^{\delta}(x) = \ell\left(\{Z_i : Z_i \leq x\}\right).$$

THEOREM 3.7 (Stability of non-dominated sorting). Let $f : \mathbb{R}^d \to \mathbb{R}$ satisfy (H1) and let $\Omega \subset \mathbb{R}^d_+$ satisfy (H2). There exist constants C_{δ} , depending only on δ , f, and g, such that

$$\limsup_{n\to\infty} n^{-\frac{1}{d}} \|u_n^{\delta} - u_n\|_{L^{\infty}(\mathbb{R}^d)} \le C_{\delta} \quad \text{a.s}$$

and $C_{\delta} \rightarrow 0$ as $\delta \rightarrow 0$.

Proof. Set $g_{\delta}(x) = \frac{1}{\delta^d} g\left(\frac{x}{\delta}\right)$. Then Z_1, \ldots, Z_n are *i.i.d.* with density $f_{\delta} := g_{\delta} * f$. Set

$$U^{\delta}(x) = \sup_{\gamma \in \mathscr{A} : \gamma \leq x} \int_0^1 f_{\delta}(\gamma(t))^{\frac{1}{d}} (\gamma_1'(t) \cdots \gamma_d'(t))^{\frac{1}{d}} dt$$

Without loss of generality, we may suppose that $\overline{\Omega} \subset (0,1)^d$. Since $\operatorname{supp}(f) \subset \overline{\Omega}$ and g has compact support, we can take $\delta > 0$ small enough so that $\operatorname{supp}(f_{\delta}) \subset [0,1]^d$. It is not hard to see that (2.35) holds for the sequence f_{δ} . Since each f_{δ} is continuous and bounded with compact support in $[0,1]^d$, it follows from Lemma 2.9 that $U^{\delta} \to U$ uniformly on \mathbb{R}^d . Note that

$$\begin{split} n^{-\frac{1}{d}} \| u_n^{\delta} - u_n \|_{L^{\infty}(\mathbb{R}^d)} &\leq \| n^{-\frac{1}{d}} u_n^{\delta} - c_d U^{\delta} \|_{L^{\infty}(\mathbb{R}^d)} + \| n^{-\frac{1}{d}} u_n - c_d U \|_{L^{\infty}(\mathbb{R}^d)} \\ &+ c_d \| U^{\delta} - U \|_{L^{\infty}(\mathbb{R}^d)}, \end{split}$$

for every *n*. Since f_{δ} is continuous on $(0,1)^d$ and $f_{\delta}(x) = 0$ for $x \notin (0,1)^d$, (H1) is satisfied for f_{δ} by taking $\Omega' = (0,1)^d$. We can therefore apply Theorem 1.2 to obtain

$$\|n^{-\frac{1}{d}}u_n^{\delta} - c_d U^{\delta}\|_{L^{\infty}(\mathbb{R}^d)} + \|n^{-\frac{1}{d}}u_n - c_d U\|_{L^{\infty}(\mathbb{R}^d)} \to 0$$
 a.s.

The proof is completed by setting $C_{\delta} = c_d \| U^{\delta} - U \|_{L^{\infty}(\mathbb{R}^d)}$.

4. Numerical demonstrations. Theorem 1.2 guarantees that the level sets of $c_d U$ will provide good approximations to the Pareto fronts for large *n*. In this section, we present a numerical scheme for computing *U* and show examples comparing the level sets of $c_d U$ to the Pareto fronts for various density functions. To compute *U*, we need to compute the Pareto-monotone viscosity solution of

(4.1)
$$U_{x_1}\cdots U_{x_d} = \frac{1}{d^d}f \text{ on } \mathbb{R}^d_+,$$

that satisfies U = 0 on $\partial \mathbb{R}^d_+$. Let U_α and f_α denote the values of U and f on a grid with spacing Δx , where α is a multi-index. For a given grid point α , the domain of dependence

for (4.1) is $\{\beta : \beta \leq \alpha\}$. Hence an upwind scheme will use backward difference quotients. Now consider substituting backward difference quotients into (4.1). We have

(4.2)
$$\prod_{i=1}^{d} (U_{\alpha} - U_{\alpha - e_i}) = \frac{\Delta x^d}{d^d} f_{\alpha}$$

We intend to solve the above equation for U_{α} in terms of $U_{\alpha-e_i}$. Since we intend to compute the Pareto-monotone solution of (4.1), we should look for a solution with $U_{\alpha} \ge U_{\alpha-e_i}$ for all *i*. Consider the mapping

$$p\mapsto \prod_{i=1}^d (p-a_i)-a_0,$$

where $a_i \ge 0$ for all *i*. Note that this mapping is strictly increasing for $p > \max(a_1, \ldots, a_d)$ and $\prod_{i=1}^d (p-a_i) = 0$ for $p = \max(a_1, \ldots, a_d)$. Hence, for any non-negative a_0, \ldots, a_d , there exists a unique $p \ge \max(a_1, \ldots, a_d)$ satisfying

$$\prod_{i=1}^d (p-a_i) = a_0.$$

We denote this solution *p* by $P(a_0, \ldots, a_d)$ and define our numerical scheme by

(4.3)
$$U_{\alpha} = P(\Delta x^{d} f_{\alpha}/d^{d}, U_{\alpha-e_{1}}, \dots, U_{\alpha-e_{d}})$$

with the boundary condition $U_{\alpha} = 0$ for α with $\min(\alpha_1, \ldots, \alpha_d) = 0$. The numerical solution can be computed by any sweeping pattern respecting the partial order \leq . This requires visiting each grid point exactly once, and hence has linear complexity. The value of $P(a_0, \ldots, a_d)$ can be computed numerically by either a binary search and/or Newton's method restricted to the interval $[\max(a_1, \ldots, a_d), \max(a_1, \ldots, a_d) + a_0^{1/d}]$. In the case of d = 2, we have a closed form expression for P

$$P(a_0, a_1, a_2) = \frac{1}{2}(a_1 + a_2) + \frac{1}{2}\sqrt{(a_1 - a_2)^2 + 4a_0}.$$

Note that we have chosen the positive square root to obtain the Pareto-monotone solution. We prove in a subsequent paper [7], that the numerical solutions, defined as above, converge to the unique Pareto-monotone viscosity solution of (4.1) as $\Delta x \rightarrow 0$.

For d = 2, we have $c_2 = 2$, hence the level sets of U will approximate the Pareto fronts, where $U_x U_y = f$. We now show examples of Pareto fronts alongside the level sets of U for X_1, \ldots, X_n sampled according to different density functions. In Figure 4.1, we consider a uniform density on a portion of the unit square and show the Pareto fronts for $n = 10^4$, $n = 10^5$ and $n = 10^6$ independent samples alongside the corresponding level sets of U. Observing the Figure, we see that the Pareto fronts are well approximated by the level sets of U for large n. We also notice that the level sets of U appear to yield a consistent underestimate of the Pareto fronts. Bollobás and Brightwell [5] showed that the normalized expectation of the longest increasing subsequence among n points chosen independently from the uniform distribution on $[0,1]^d$ is always bounded above by c_d , which is the limit of these normalized expectations as $n \to \infty$. In light of this result, our observation is not surprising and merely confirms the results in [5]. We also observe that although the boundary of Ω is smooth, the solution U develops shocks, or kinks, which are visible in the level sets of U. In Figure 4.2, we show the same comparison for a multi-modal density function on $[0,1]^2$. The density function is depicted by the plot in Figure 4.2 and we have the same expected underestimation present here as well.



FIG. 4.1. Comparison of the Pareto fronts and the level sets of U, where $U_xU_y = f$ and f is the uniform density on the gray region in the top left. The plots correspond to the Pareto fronts computed with $n = 10^4$, $n = 10^5$ and $n = 10^6$ independent samples from f. In each case, we show 15 equally spaced Pareto fronts and the corresponding level sets of U.

Acknowledgments. We thank John Duchi for pointing out an error in the proof of Theorem 1.2 in an earlier version of the manuscript

REFERENCES

- G.S. ADHAR, Parallel algorithms for chains and anti-chains of points on a plane, in International Conference on Parallel and Distributed Systems (ICPADS), vol. 2, Dec. 2007, pp. 1–7.
- [2] D. ALDOUS AND P. DIACONIS, Hammersley's interacting particle process and longest increasing subsequences, Probability Theory and Related Fields, 103 (1995), pp. 199–213.
- [3] M. BARDI AND I.C. DOLCETTA, Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations, Springer, 1997.
- [4] G. BARLES, A. BRIANI, AND E. CHASSEIGNE, A Bellman approach for two-domains optimal control problems in Rn, arXiv preprint:1112.3727, (2011).
- [5] BÉLA BOLLOBÁS AND GRAHAM BRIGHTWELL, The height of a random partial order: concentration of measure, The Annals of Applied Probability, 2 (1992), pp. 1009–1018.
- [6] BÉLA BOLLOBÁS AND PETER WINKLER, The longest chain among random points in Euclidean space, Proceedings of the American Mathematical Society, 103 (1988), pp. 347–353.
- [7] JEFF CALDER, SELIM ESEDOĞLU, AND ALFRED O HERO, A PDE-based approach to non-dominated sorting, arXiv preprint:1310.2498, (2013).
- [8] F. CAMILLI AND E. PRADOS, Shape-from-shading with discontinuous image brightness, Applied Numerical Mathematics, 56 (2006), pp. 1225–1237.
- [9] F. CAMILLI AND A. SICONOLFI, Hamilton-Jacobi equations with measurable dependence on the state vari-



FIG. 4.2. Comparison of the Pareto fronts and the level sets of U, where $U_x U_y = f$ and f is the density depicted by the plot in the top left. The plots correspond to the Pareto fronts computed with $n = 10^4$, $n = 10^5$ and $n = 10^6$ independent samples from f. In each case, we show 15 equally spaced Pareto fronts and the corresponding level sets of U.

able, Advances in Differential Equations, 8 (2003), pp. 733-768.

- [10] T.F. CHAN AND L.A. VESE, Active contours without edges, IEEE Transactions on Image Processing, 10 (2001), pp. 266–277.
- [11] M.G. CRANDALL, H. ISHII, AND P.L. LIONS, User's guide to viscosity solutions of second order partial differential equations, Bulletin of the American Mathematical Society, 27 (1992), pp. 1–67.
- [12] K. DEB, Multi-objective optimization using evolutionary algorithms, Wiley, Chichester, UK, 2001.
- [13] K. DEB, A. PRATAP, S. AGARWAL, AND T. MEYARIVAN, A fast and elitist multiobjective genetic algorithm: NSGA-II, IEEE Transactions on Evolutionary Computation, 6 (2002), pp. 182–197.
- [14] KLAUS DECKELNICK AND CHARLES M. ELLIOTT, Uniqueness and error analysis for Hamilton-Jacobi equations with discontinuities, Interfaces and Free Boundaries, 6 (2004), pp. 329–349.
- [15] JEAN-DOMINIQUE DEUSCHEL AND OFER ZEITOUNI, Limiting curves for i.i.d. records, The Annals of Probability, 23 (1995), pp. 852–878.
- [16] S. FELSNER AND L. WERNISCH, Maximum k-chains in planar point sets: Combinatorial structure and algorithms, SIAM Journal on Computing, 28 (1999), pp. 192–209.
- [17] G. FLEURY, A. HERO, S. YOSHIDA, T. CARTER, C. BARLOW, AND A. SWAROOP, Pareto analysis for gene filtering in microarray experiments, in European Signal Processing Conference (EUSIPCO), 2002.
- [18] G. FLEURY, A. HERO, S. ZAREPARSI, AND A. SWAROOP, Gene discovery using Pareto depth sampling distributions, Journal of the Franklin Institute, 341 (2004), pp. 55–75.
- [19] ——, Pareto depth sampling distributions for gene ranking, in IEEE International Symposium on Biomedical Imaging: Nano to Macro, vol. 1, Apr. 2004, pp. 65–68.
- [20] C.M. FONSECA AND P.J. FLEMING, Genetic algorithms for multiobjective optimization : formulation, discussion and generalization, Proceedings of the Fifth International Conference on Genetic Algorithms, 1

(1993), pp. 416-423.

- [21] ——, An overview of evolutionary algorithms in multiobjective optimization, Evolutionary Computation, 3 (1995), pp. 1–16.
- [22] M. GARAVELLO AND P. SORAVIA, Optimality principles and uniqueness for bellman equations of unbounded control problems with discontinuous running cost, Nonlinear Differential Equations and Applications (NoDEA), 11 (2004), pp. 271–298.
- [23] ——, Representation formulas for solutions of the HJI equations with discontinuous coefficients and existence of value in differential games, Journal of Optimization Theory and Applications, 130 (2006), pp. 209–229.
- [24] J.M. HAMMERSLEY, A few seedlings of research, in Proceedings of the Sixth Berkeley Symposium on Mathematical Statistics and Probability, vol. 1, 1972, pp. 345–394.
- [25] JULIA HANDL AND JOSHUA KNOWLES, Exploiting the trade-off The benefits of multiple objectives in data clustering, in Evolutionary Multi-Criterion Optimization, Springer, 2005, pp. 547–560.
- [26] A.O. HERO, Gene selection and ranking with microarray data, in IEEE International Symposium on Signal Processing and its Applications, vol. 1, July 2003, pp. 457–464.
- [27] A. HERO AND G. FLEURY, Posterior Pareto front analysis for gene filtering, in Proceedings of the Workshop on Genomic Signal Processing and Statistics (GENSIPS), 2002.
- [28] ——, Pareto-optimal methods for gene ranking., VLSI Signal Processing, 38 (2004), pp. 259–275.
- [29] DONALD KOSSMANN, FRANK RAMSAK, STEFFEN ROST, ET AL., Shooting stars in the sky: an online algorithm for skyline queries, in Proceedings of the 28th International Conference on Very Large Data Bases, 2002, pp. 275–286.
- [30] A. KUMAR AND A. VLADIMIRSKY, An efficient method for multiobjective optimal control and optimal control subject to integral constraints, Journal of Computational Mathematics, 28 (2010), pp. 517–551.
- [31] B. F. LOGAN AND L. A. SHEPP, A variational problem for random Young tableaux, Advances in Mathematics, 26 (1977), pp. 206–222.
- [32] R.D. LOU AND M. SARRAFZADEH, An optimal algorithm for the maximum three-chain problem, SIAM Journal on Computing, 22 (1993), pp. 976–993.
- [33] N.K. MADAVAN, Multiobjective optimization using a Pareto differential evolution approach, in Proceedings of the Congress on Evolutionary Computation (CEC), 2002.
- [34] I.M. MITCHELL AND S. SASTRY, Continuous path planning with multiple constraints, in IEEE Conference on Decision and Control, vol. 5, Dec. 2003, pp. 5502–5507.
- [35] D. MUMFORD AND J. SHAH, Optimal approximations by piecewise smooth functions and associated variational problems, Communications on Pure and Applied Mathematics, 42 (1989), pp. 577–685.
- [36] RT NEWCOMB AND J. SU, Eikonal equations with discontinuities, Differential and Integral Equations-Athens, 8 (1995), pp. 1947–1947.
- [37] DIMITRIS PAPADIAS, YUFEI TAO, GREG FU, AND BERNHARD SEEGER, Progressive skyline computation in database systems, ACM Transactions on Database Systems (TODS), 30 (2005), pp. 41–82.
- [38] P. PEVZNER, Computational Molecular Biology, The MIT Press, 2000.
- [39] TIMO SEPPÄLÄINEN, A growth model in multiple dimensions and the height of a random partial order, in Asymptotics: particles, processes and inverse problems, vol. 55 of IMS Lecture Notes Monograph Series, Institute for Mathematical Statistics, Beachwood, OH, 2007, pp. 204–233.
- [40] P. SORAVIA, Boundary value problems for Hamilton-Jacobi equations with discontinuous Lagrangian, Indiana University Mathematics Journal, 51 (2002), pp. 451–477.
- [41] T. SPEED, Statistical analysis of gene expression microarray data, vol. 11, Chapman & Hall/CRC, 2003.
- [42] N. SRINIVAS AND K. DEB, Muiltiobjective optimization using nondominated sorting in genetic algorithms, Evolutionary Computation, 2 (1994), pp. 221–248.
- [43] A. TOURIN, A comparison theorem for a piecewise Lipschitz continuous Hamiltonian and application to shape-from-shading problems, Numerische Mathematik, 62 (1992), pp. 75–85.
- [44] SM ULAM, *Monte carlo calculations in problems of mathematical physics*, Modern Mathematics for the Engineers, (1961), pp. 261–281.
- [45] A.M. VERSHIK AND S.V. KEROV, Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tables, Soviet Doklady Mathematics, 18 (1977), p. 38.
- [46] G. VIENNOT, Chain and antichain families, grids and Young tableaux, in Orders: description and roles, vol. 99 of North-Holland Mathematics Studies, 1984, pp. 409–463.