

Appendix B. Extension of Continuous Functions

Let $f(x)$ be a continuous function on a compact set $K \subset \mathbb{R}^d$. Then it is uniformly continuous on K , i.e. its **modulus of continuity**

$$\omega(\rho) := \sup \left\{ |f(x) - f(y)| : x, y \in K, |x - y| \leq \rho \right\} \searrow 0 \quad \text{as } \rho \searrow 0. \quad (1)$$

Lemma 1. *If K is convex, then $\omega(\rho)$ is subadditive, i.e.*

$$\omega(\rho_1 + \rho_2) \leq \omega(\rho_1) + \omega(\rho_2) \quad \text{for } \rho_1, \rho_2 \geq 0. \quad (2)$$

Proof. For arbitrary $x, y \in K$ with $|x - y| \leq \rho_1 + \rho_2$, the segment $[x, y]$ lies in K , and there is a point $z \in [x, y]$ such that $|x - z| \leq \rho_1$, $|y - z| \leq \rho_2$. Therefore,

$$|f(x) - f(y)| \leq |f(x) - f(z)| + |f(z) - f(y)| \leq \omega(\rho_1) + \omega(\rho_2),$$

and (2) follows. □

Remark 2. *The property (2) fails in general if K is not convex. In polar coordinates $x = r \cos \theta$, $y = r \sin \theta$, an easy example is*

$$f(x, y) := \theta \quad \text{on } K := \{1 \leq r \leq 2, \quad \varepsilon \leq \theta \leq 2\pi - \varepsilon\} \quad \text{with a small } \varepsilon > 0.$$

Lemma 3. *For any continuous function f on a compact $K \subset \mathbb{R}^d$ with the modulus of continuity $\omega(\rho)$ in (1), the function*

$$\bar{\omega}(\rho) := \sup_{s \geq 1} \frac{\omega(\rho s)}{s}, \quad \rho \geq 0, \quad (3)$$

satisfies the properties:

- (i) $\bar{\omega}(\rho) \geq \omega(\rho)$,
- (ii) $\bar{\omega}(\rho) \searrow 0$ as $\rho \searrow 0$,
- (iii) $\bar{\omega}(\rho)$ is subadditive, i.e. it satisfies (2).

Proof. (i) is obvious.

(ii) Since $\omega(\rho)$ is non-decreasing, the function $\bar{\omega}(\rho)$ is also non-decreasing for $\rho \geq 0$. We also have

$$\omega(\rho) \leq C_0 := 2 \sup_K |f|, \quad \text{and} \quad \bar{\omega}(\rho) \leq C_0.$$

For an arbitrary $A > 1$, we can write

$$\bar{\omega}(\rho) = \max \left\{ \sup_{A \geq s \geq 1} \frac{\omega(\rho s)}{s}, \quad \sup_{s \geq A} \frac{\omega(\rho s)}{s} \right\} \leq \max \left\{ \omega(A\rho), \frac{C_0}{A} \right\}.$$

This implies

$$\limsup_{\rho \rightarrow 0^+} \bar{\omega}(\rho) \leq \frac{C_0}{A},$$

and since $A > 1$ can be chosen arbitrarily large, we have $\bar{\omega}(\rho) \searrow 0$ as $\rho \searrow 0$.

(iii) We can write

$$\bar{\omega} := \sup_{s \geq 1} \frac{\omega(\rho s)}{s} = \rho \cdot \sup_{t \geq \rho} \frac{\omega(t)}{t} \quad \text{for } \rho > 0.$$

Therefore,

$$\begin{aligned} \bar{\omega}(\rho_1 + \rho_2) &= (\rho_1 + \rho_2) \cdot \sup_{t \geq \rho_1 + \rho_2} \frac{\omega(t)}{t} \\ &\leq \rho_1 \cdot \sup_{t \geq \rho_1} \frac{\omega(t)}{t} + \rho_2 \cdot \sup_{t \geq \rho_2} \frac{\omega(t)}{t} \\ &= \bar{\omega}(\rho_1) + \bar{\omega}(\rho_2). \end{aligned}$$

□

Theorem 4. *Let f be a continuous function on a compact $K \subset \mathbb{R}^d$. Then the function*

$$F(x) := \inf_{y \in K} \left[f(y) + \bar{\omega}(|x - y|) \right], \quad x \in \mathbb{R}^d, \quad (4)$$

where $\bar{\omega}(\rho)$ is defined in (3), satisfies the properties:

- (i) $F \equiv f$ on K ,
- (ii) F provides a continuous extension of f from K to \mathbb{R}^d , and

$$|F(x_1) - F(x_2)| \leq \bar{\omega}(|x_1 - x_2|), \quad \forall x_1, x_2 \in \mathbb{R}^d. \quad (5)$$

Proof. (i) Note that $\forall x, y \in K$, we have

$$f(x) \leq f(y) + \omega(|x - y|) \leq f(y) + \bar{\omega}(|x - y|),$$

with the equality at $y = x$. Therefore, $f(x) = F(x)$, $\forall x \in K$.

- (ii) By monotonicity and subadditivity of $\bar{\omega}(\rho)$, we have $\forall x_1, x_2, y \in K$:

$$\bar{\omega}(|x_1 - y|) \leq \bar{\omega}(|x_1 - x_2| + |x_2 - y|) \leq \bar{\omega}(|x_1 - x_2|) + \bar{\omega}(|x_2 - y|).$$

Hence the function $F(x)$ in (4) satisfies

$$F(x_1) \leq \bar{\omega}(|x_1 - x_2|) + F(x_2).$$

Interchanging x_1 and x_2 , we get the desired property (5). □