

Dedicated to Nina N. Ural'tseva

Non-divergence Elliptic Equations of Second Order with Unbounded Drift

M. V. Safonov

ABSTRACT. We consider uniformly elliptic equations and inequalities of second order in the *nondivergence* form $Lu = a_{ij}D_{ij}u + b_iD_iu = 0$ (≤ 0 , ≥ 0), in a domain $\Omega \subset \mathbb{R}^n$. We derive an interior Harnack inequality, a boundary Harnack inequalities and a comparison theorem for Lipschitz boundaries in the case $b_i \in L^n$, and a Hopf-Oleinik type estimate near flat boundary in the case when the normal component of the vector (b_1, \dots, b_n) belongs to L^q , $q > n$. Our main tools are special Landis-type growth lemmas.

1. Introduction

In this paper, we derive interior and boundary pointwise estimates for solutions to the equation

$$(1.1) \quad Lu := a_{ij}D_{ij}u + b_iD_iu = 0 \quad \text{in a domain } \Omega \subset \mathbb{R}^n,$$

and the inequalities $Lu \leq 0$ or $Lu \geq 0$, under minimal assumptions on the functions a_{ij} and b_i . We use the summation convention over repeated indices and the notations $D_i := \partial/\partial x_i$, $D_{ij} := D_iD_j$. We assume that a_{ij} and b_i are measurable functions on \mathbb{R}^n , a_{ij} satisfy the *uniform ellipticity condition*

$$(1.2) \quad a_{ij}(x) = a_{ji}(x), \quad \nu |\xi|^2 \leq a_{ij}(x)\xi_i\xi_j \leq \nu^{-1}|\xi|^2 \quad \forall \xi, x \in \mathbb{R}^n,$$

with a constant $\nu \in (0, 1]$, and $b_i, f \in L^n(\Omega)$. Throughout the paper, the operator L in (1.1) is applied to functions u in the class $W(\Omega) := W_{loc}^{2,n}(\Omega) \cap C(\overline{\Omega})$, which implies, in particular, that $u, D_iu, D_{ij}u$ belong to the Lebesgue space $L^n(\Omega')$ for any bounded open set $\Omega' \subset \overline{\Omega'} \subset \Omega$.

The following estimate is crucial for our considerations. It can be considered as a particular case (when $m = n$) of the estimate (5.6) in the paper by A.D. Aleksandrov [A63] (see also [GT83], (9.14)).

THEOREM 1.1. *Let Ω be a bounded open set in \mathbb{R}^n , and let u be a function in $W(\Omega)$ such that $Lu \geq f$ in Ω . Suppose that the coefficients a_{ij} satisfy (1.2), and $b_i, f \in L^n(\Omega)$. Then*

$$(1.3) \quad \sup_{\Omega} u \leq \sup_{\partial\Omega} u + N \operatorname{diam} \Omega \cdot e^{NS} \|f_-\|_{n,\Omega},$$

where $f_- := \max\{-f, 0\}$,

$$(1.4) \quad S := S(\Omega) := \int_{\Omega} |\mathbf{b}|^n dx, \quad \mathbf{b} := (b_1, \dots, b_n),$$

1991 *Mathematics Subject Classification.* Primary 35J15, 35J67; Secondary 35B05.

Key words and phrases. Second-order elliptic equations, Harnack inequality, boundary Hopf lemma, measurable coefficients.

$$\|g\|_{n,\Omega} := \left(\int_{\Omega} |g|^n dx \right)^{1/n} \quad \text{-- the norm of } g \text{ in } L^n(\Omega),$$

and N is a positive constant depending only on n and ν .

COROLLARY 1.2. *Let Ω be a bounded open set in \mathbb{R}^n , and let u be a function in $W(\Omega)$ such that $Lu \geq 0$ in Ω . Then the maximum of u on $\bar{\Omega}$ is attained on $\partial\Omega$.*

REMARK 1.3. Note that the equality $Lu = f$ in (1.1) and the inequality $Lu \geq f$ in Theorem 1.1 hold true almost everywhere in Ω with respect to the Lebesgue measure on \mathbb{R}^n . It is easy to see that the function $u(x) := 1 - |x|^2$ satisfies

$$\Delta u + b_i D_i u = 0 \quad \text{in } B_1 := \{x \in \mathbb{R}^n : |x| < 1\},$$

where $b_i(x) := -n x_i \cdot |x|^{-2} \in L^{n-\varepsilon}(B_1)$ for arbitrary small $\varepsilon > 0$, and the maximum of u on \bar{B}_1 is not attained on ∂B_1 . Therefore, the assumption $b_i \in L^n$ is essential for the whole theory, which is based on the maximum principle.

REMARK 1.4. Note that the quantity $S(\Omega)$ in (1.4) is scaling invariant in the following sense. If $u = u(x)$ satisfies the equation $Lu = 0$ in Ω , then for any constant $k > 0$, the function $\tilde{u}(y) := u(ky)$ satisfies $D_i \tilde{u} = k D_i u$, $D_{ij} \tilde{u} = k^2 D_{ij} u$, hence

$$\tilde{L}\tilde{u} := \tilde{a}_{ij} D_{ij} \tilde{u} + \tilde{b}_i D_i \tilde{u} = 0 \quad \text{in } \tilde{\Omega} := \{y \in \mathbb{R}^n : ky \in \Omega\},$$

where $\tilde{a}_{ij}(y) := a_{ij}(ky)$, $\tilde{b}_i(y) := k b_i(ky)$. Therefore, $|\tilde{\mathbf{b}}| := (\sum \tilde{b}_i^2)^{1/2}$ satisfies

$$\tilde{S}(\tilde{\Omega}) := \int_{\tilde{\Omega}} |\tilde{\mathbf{b}}(y)|^n dy = \int_{\tilde{\Omega}} |\mathbf{b}(ky)|^n k^n dy = \int_{\Omega} |\mathbf{b}(x)|^n dx =: S(\Omega).$$

Obviously, this argument also works for inequalities $Lu \leq 0$ or $Lu \geq 0$ in Ω .

In the next section, we will use the scaling invariance of $S(\Omega)$, together with the maximum principle, in order to prove some special *growth lemmas* (Lemmas 2.1 and 2.5) and a *doubling property* of solutions (Lemma 2.2) in the case $|\mathbf{b}| \in L^n$, i.e. $S(\Omega) < \infty$. The growth lemmas were first introduced by E.M. Landis [La67, La71]. Among other applications of these lemmas, Landis gave alternative proofs of results by De Giorgi and Moser on Hölder regularity and Harnack inequalities for solutions to second order elliptic equations in the *divergence* form. These results were extended to the equations in the *non-divergence* form with $|\mathbf{b}| \in L^\infty$ in our joint work with N.V. Krylov [KS80, S80] (see also the book [K85]). For this purpose, we also used some variants of growth lemmas. Now this technique became standard, see the paper by H. Aimar, L. Forzani, and R. Toledano [AFT01], in which they treat Hölder and Harnack properties from an abstract point of view.

In Section 3, we use the growth lemmas and the doubling property of solutions in order to derive the interior and boundary Harnack inequalities and the Hölder estimates for solutions of equations (1.1) with $|\mathbf{b}| \in L^n$. Note that the Hölder regularity in the case $|\mathbf{b}| \in L^n$ was proved earlier by O.A. Ladyzhenskaya and N.N. Ural'tseva [LU85]. However, the Hölder constants in that paper depend on the modulus of continuity of $|\mathbf{b}|$ in L^n , or more precisely, on the constant $\rho > 0$ for which the norms of $|\mathbf{b}|$ in $L^n(\Omega \cap B_\rho(x))$ are small enough for all $x \in \Omega$. In our estimates, all the constants depend on \mathbf{b} only through the quantity $S(\Omega)$ in (1.4).

Finally, in Section 4, we prove a Hopf-Oleinik type estimate near flat boundary $\{x_n = 0\}$ for positive solutions of the equation $a_{ij} D_{ij} u + b_i D_i u = 0$ with $|\mathbf{b}| \in L^n$, $b_n \in L^q$, $q > n$. A close result is contained in the paper by O.A. Ladyzhenskaya

and N.N. Ural'tseva [LU88], Lemma 4.4, under a bit stronger assumption $|\mathbf{b}| \in L^q$, $q > n$. See also a recent work by A.I. Nazarov and N.N. Ural'tseva [NU09] for further results concerning elliptic and parabolic, divergence and nondivergence, equations with unbounded lower-order coefficients.

Notations. $x = (x_1, \dots, x_{n-1}, x_n) = (x', x_n)$ are vectors or points in \mathbb{R}^n , $(x, y) := x_i y_i$ – the scalar product of $x, y \in \mathbb{R}^n$ (the summation convention is implied), and $|x| := (x, x)^{1/2}$ – the length of $x \in \mathbb{R}^n$. The standard orthonormal basis in \mathbb{R}^n consists of n vectors $\{\mathbf{e}_1, \dots, \mathbf{e}_n\}$, so that $x = x_i \mathbf{e}_i$ for all $x \in \mathbb{R}^n$. $\text{tr } a := \sum_i a_{ii}$ – the trace of a $n \times n$ matrix $a = [a_{ij}]$. For $c \in \mathbb{R}^1$, $[c]$ denotes its integer part (the maximal integer $\leq c$), $c_+ := \max\{c, 0\}$, $c_- := \max\{-c, 0\}$.

$B_r(x) := \{y \in \mathbb{R}^n : |y - x| < r\}$ – a ball of radius $r > 0$ centered at $x \in \mathbb{R}^n$.

$B'_r(x') := \{y' \in \mathbb{R}^{n-1} : |y' - x'| < r\}$ is a similar ball in \mathbb{R}^{n-1} centered at $x' \in \mathbb{R}^{n-1}$. $B_r := B_r(0)$, $B'_r := B'_r(0)$.

$\partial\Omega$ is the boundary of an open set $\Omega \subset \mathbb{R}^n$, $|\Omega|$ – its Lebesgue measure.

The notation $A := B$, or $B =: A$, means “ $A = B$ by definition”. Throughout the paper, N, c (with indices or without) denote different constants depending only on the prescribed quantities, such that n, ν , etc. This dependence is indicated in the parentheses: $N = N(n, \nu, \dots)$, $c = c(n, \nu, \dots)$.

2. Growth Lemmas

The first growth lemma can be treated as a growth lemma for “narrow” domains.

LEMMA 2.1 (First growth lemma). *Let Ω be an open set in \mathbb{R}^n , and let $u \in W(\Omega)$, $x_0 \in \mathbb{R}^n$, and $r > 0$ be such that*

$$u \geq 0, \quad Lu \geq 0 \quad \text{in } \Omega; \quad \text{and} \quad u = 0 \quad \text{on} \quad (\partial\Omega) \cap B_{2r}(x_0).$$

We claim that for arbitrary constant $\beta_1 \in (0, 1)$, there is a constant $\mu_1 \in (0, 1)$, depending only on n, ν , and S , such that from the estimate for the Lebesgue measure

$$(2.1) \quad |\Omega \cap B_{2r}(x_0)| \leq \mu_1 \cdot |B_{2r}|$$

it follows

$$(2.2) \quad M_r := \sup_{\Omega \cap B_r(x_0)} u \leq \beta_1 \cdot M_{2r}.$$

PROOF. Using translation in \mathbb{R}^n , and multiplying u by a constant, we can assume $x_0 = 0$, $M_r > 0$, and $M_{2r} = 1$. Moreover, since $S(\Omega)$ is invariant with respect to rescaling (Remark 1.4), we can choose any convenient value $r > 0$.

(a) We first take $r = 1$ and assume that $S := S(\Omega) \leq 1$. By Corollary 1.2, the maximum of u on $\overline{\Omega} \cap \overline{B_1}$ is attained at some point $y_0 \in \Omega \cap (\partial B_1)$. Consider the function $v(x) := u(x) - |x - y_0|^2$ on the set $\Omega' := \Omega \cap B_1(y_0)$. Since $v \leq u = 0$ on $(\partial\Omega) \cap B_2$, and $v = u - 1 \leq 0$ on $\overline{\Omega} \cap (\partial B_1(y_0))$, we have $v \leq 0$ on $\partial\Omega'$. Moreover,

$$Lv = Lu - 2 \text{tr } a - 2(\mathbf{b}, x - y_0) \geq f := -2n\nu^{-1} - 2|\mathbf{b}| \quad \text{in } \Omega'.$$

Applying Theorem 1.1 to the function v in Ω' , and having in mind that $S(\Omega) \leq 1$ and $|\Omega'| \leq \mu_1 \cdot |\Omega \cap B_2| \leq \mu_1 \cdot |B_2|$, we get

$$(2.3) \quad M_1 := \sup_{\Omega \cap B_1} u = u(y_0) = v(y_0) \leq N_0 \cdot \|f\|_{n, \Omega'} \leq N_1 \cdot (\mu_1^{1/n} + S^{1/n}),$$

with constants $N_0, N_1 \geq 1$, depending only on n and ν .

Now fix an arbitrary constant $\beta_1 \in (0, 1)$, and set $\mu_0 := S_0 := (\beta_1/2N_1)^n$. If $S \leq S_0$, we can take $\mu_1 = \mu_0$, and then the desired estimate $M_1 \leq \beta_1$ follows from (2.3).

(b) In the remaining case $S > S_0$, we set $m := [S/S_0] + 1 > S/S_0 > 1$. It is convenient to take $r := 4m$. Then we divide $\Omega \cap (B_{8m} \setminus \overline{B_{4m}})$ into m disjoint sets

$$\Omega_k := \Omega \cap (B_{4m+4k} \setminus \overline{B_{4m+4k-4}}) \quad \text{for } k = 1, 2, \dots, m.$$

We have

$$\sum_{k=1}^m \int_{\Omega_k} |\mathbf{b}|^n dx \leq \int_{\Omega} |\mathbf{b}|^n dx =: S.$$

Since $S < mS_0$, at least one of the integrals on the left side $\leq S_0$. Fix the corresponding index $k = k_0$ for which this is the case. Finally, we set $\mu_1 := (4m)^{-n} \mu_0$. By the maximum principle,

$$(2.4) \quad M_r := \sup_{\Omega \cap B_{4m}} u \leq \sup_{\Omega \cap B_{4m+4k_0-2}} u = u(y)$$

for some $y \in \Omega \cap (\partial B_{4m+4k_0-2})$. Note that the set $\Omega \cap B_2(y)$ is contained in Ω_{k_0} hence the integral of $|\mathbf{b}|^n$ over this set $\leq S_0$, and moreover, from (2.1) it follows

$$|\Omega \cap B_2(y)| \leq |\Omega \cap B_{8m}| \leq \mu_1 \cdot |B_{8m}| = \mu_0 \cdot |B_2|.$$

From the previous part (a) of the proof it follows that

$$u(y) \leq \sup_{\Omega \cap B_1(y)} u \leq \beta_1 \cdot \sup_{\Omega \cap B_2(y)} u \leq \beta_1 \cdot M_{2r}.$$

Together with (2.4), this completes the proof. \square

LEMMA 2.2 (Doubling property). *Let $x_0 \in \mathbb{R}^n$, $r > 0$, and a function v in $W(B_{4r}(x_0))$ be such that*

$$(2.5) \quad v \geq 0, \quad Lv \leq 0 \quad \text{in } B_{4r}(x_0); \quad \text{and } v \geq 1 \quad \text{in } B_r(x_0).$$

Then

$$(2.6) \quad v \geq \lambda = \lambda(n, \nu, S) > 0 \quad \text{in } B_{2r}(x_0).$$

PROOF. Without loss of generality, we assume $x_0 = 0$, $r = 1$. In the part (a) below, we consider the case $n = 1$, part(b) deals with $n \geq 2$ and small S , and part (c) – with general $S < \infty$. We need to treat the case $n = 1$ separately, because in our approach, we use the fact that the spheres $\partial B_r \subset \mathbb{R}^n$ are connected. This fails if $n = 1$, because in this case ∂B_r consists of two disjoint point $\pm r$.

(a) In the case $n = 1$, we have

$$v \geq 0, \quad av'' + bv' \leq 0 \quad \text{in } (-4, 4); \quad \text{and } v \geq 1 \quad \text{on } [-1, 1],$$

where $0 < \nu \leq a \leq \nu^{-1}$, and $b \in L^1(-4, 4)$. Using properties of weak derivatives (the *product formula* (7.18) and the *chain rule* in Lemma 7.5, [GT83]), one can rewrite the above differential inequality as follows:

$$(2.7) \quad (Av')' \leq 0, \quad \text{where } A(x) := \exp \left[\int_0^x \frac{b(y)}{a(y)} dy \right].$$

The function $A(x)$ satisfies

$$(2.8) \quad N_0^{-1} \leq A(x) \leq N_0 := \exp(\nu^{-1}S) \quad \text{in } (-4, 4), \quad S := \int_{-4}^4 |b(x)|dx.$$

It suffices to get the estimate $v \geq \lambda = \lambda(n, \nu, S) > 0$ on the interval $[-2, -1]$, because a similar estimate on the symmetric interval $[1, 2]$ can be obtained by replacing x with $-x$. Denote $\kappa := \min_{[-2, -1]} v$. We can assume $\kappa \leq 1/2$, because otherwise the estimate $v \geq \lambda$ holds true with $\lambda = 1/2$. By the maximum principle, i.e. Corollary 1.2 applied to the function $-v$, we have $\kappa = v(-2)$. Since v is continuous and $v(-2) = \kappa \leq 1/2 < 1 \leq v(-1)$, there is $h \in (0, 1]$ such that

$$v(-2+h) = 1, \quad \text{and} \quad v(x) < 1 \quad \text{for all } x \in [-2, -2+h].$$

Furthermore, for each $y \in (-4, -2+h)$ the minimum of $v(x)$ on $[y, -2+h]$ is attained at the point y . This means that the function $v(x)$ is non-decreasing on $(-4, -2+h]$, and therefore $v' \geq 0$ in $(-4, -2+h)$. From (2.7) it follows that the function Av' is non-increasing. Together with (2.8), this implies

$$\begin{aligned} \kappa = v(-2) &\geq \int_{-2-h}^{-2} v' dx \geq \frac{1}{N_0} \int_{-2-h}^{-2} Av' dx \geq \frac{1}{N_0} \int_{-2}^{-2+h} Av' dx \\ &\geq \frac{1}{N_0^2} \int_{-2}^{-2+h} v' dx = \frac{1}{N_0^2} [v(-2+h) - v(-2)] \geq \frac{1}{2N_0^2} =: \lambda. \end{aligned}$$

This proves the estimate (2.6) in the case $n = 1$.

(b) In the rest of the proof of this lemma, we assume $n \geq 2$. First consider the case when $S := S(\Omega) \leq S_0 = S_0(n, \nu)$ – a small positive constant to be chosen below in (2.10). Note that

$$\begin{aligned} L_0(|x|^{-\gamma_0}) &:= a_{ij}D_{ij}(|x|^{-\gamma_0}) = \gamma_0 \cdot \left[\frac{(\gamma_0 + 2)a_{ij}x_i x_j}{|x|^2} - \text{tr } a \right] \cdot |x|^{-\gamma_0-2} \\ &\geq \gamma_0 \cdot [(\gamma_0 + 2)\nu - n\nu^{-1}] \cdot |x|^{-\gamma_0-2} \geq 0 \quad \text{for } x \neq 0, \end{aligned}$$

provided $\gamma_0 + 2 \geq n\nu^{-2}$. Fix such $\gamma_0 = \gamma_0(n, \nu) > 0$, and consider the function

$$w(x) := \frac{|x|^{-\gamma_0} - 4^{-\gamma_0}}{1 - 4^{-\gamma_0}} \quad \text{in } \Omega_1 := \{x \in \mathbb{R}^n : 1 < |x| < 4\}.$$

We have $L_0 w \geq 0$ in Ω_1 , $w \leq v$ on $\partial\Omega_1$. Therefore,

$$L(w - v) \geq Lw = L_0 w + (b, Dw) \geq (b, Dw) \quad \text{in } \Omega_1,$$

and by Theorem 1.1,

$$(2.9) \quad \sup_{\Omega_1} (w - v) \leq N_0 \cdot \|(b, Dw)\|_{n, \Omega_1} \leq N_1 \cdot \|b\|_{n, \Omega} = N_1 \cdot S^{1/n},$$

with some constants $N_0, N_1 \geq 1$ depending only on n and ν . Next, fix the constants

$$(2.10) \quad \lambda_0 := \frac{2^{-\gamma_0} - 4^{-\gamma_0}}{2 \cdot (1 - 4^{-\gamma_0})} \in \left(0, \frac{1}{2}\right), \quad \text{and} \quad S_0 := \left(\frac{\lambda_0}{N_1}\right)^n,$$

which also depend only on n and ν . Then from $S \leq S_0$ and (2.9) it follows $w - v \leq \lambda_0$ in Ω_1 . On the other hand, obviously $w \geq 2\lambda_0$ on the set $\{1 \leq |x| \leq 2\} \subset \Omega_1$. Therefore, $v(x) \geq \lambda_0$ for $1 \leq |x| \leq 2$. Since also $v(x) \geq 1$ for $|x| < 1$, the desired estimate (2.6) holds true with $\lambda = \lambda_0(n, \nu) > 0$, provided $S \leq S_0 = S_0(n, \nu)$.

(c) Now it remains to consider the case $n \geq 2$, $S > S_0$. As in the proof of Lemma 2.1, we set $m := [S/S_0] + 1 > S/S_0 > 1$. Choose $\rho_0 \in (0, 1/2)$ of order $1/2$ and $y_1, \dots, y_m \in \partial B_{1/2}$ such that the balls $B_{\rho_0}(y_k)$ are disjoint for $k = 1, \dots, m$. Then the “tubes”

$$T_k := B_4 \cap \left(\bigcup_{t \geq 1} B_{\rho_0}(ty_k) \right), \quad k = 1, \dots, m,$$

are also disjoint. Since $S < mS_0$, we can fix $k = k_1$ such that the integral of $|\mathbf{b}|^n$ over T_{k_1} does not exceed S_0 . Further, divide $B_4 \setminus \overline{B_2}$ into m disjoint shells

$$\Omega_k := B_{r_k} \setminus \overline{B_{r_{k-1}}}, \quad \text{where } r_k := 2 + \frac{2k}{m}; \quad k = 1, \dots, m,$$

and choose $k = k_2$ such that the integral of $|\mathbf{b}|^n$ over Ω_{k_2} does not exceed S_0 . Finally, set

$$R := 2 + \frac{2k_2 - 1}{m} \in \left(2 + \frac{1}{m}, 4 - \frac{1}{m} \right), \quad \rho := \frac{1}{4} \min \left\{ \rho_0, \frac{1}{m} \right\}.$$

Fix an arbitrary $y \in \partial B_R$. One can move the ball $B_{4\rho}(z)$ continuously inside T_{k_1} from $z = y_{k_1} \in \partial B_{1/2}$ to $z = 2R \cdot y_{k_1} \in \partial B_R$, and then inside Ω_{k_2} to $z = y$. Therefore, there is a sequence

$$z_0 := y_{k_1}, z_1, \dots, z_{j_0}, z_{j_0+1}, \dots, z_{m_0} := y,$$

such that

$$(2.11) \quad B_{4\rho}(z_j) \subset T_{k_1} \quad \text{for } j \leq j_0; \quad B_{4\rho}(z_j) \subset \Omega_{k_2} \quad \text{for } j \geq j_0 + 1,$$

$|z_{j+1} - z_j| \leq \rho$ for all $j = 0, 1, \dots, m_0 - 1$, and m_0 is of order m .

We claim that

$$(2.12) \quad v \geq \lambda_0^j \quad \text{in } B_\rho(z_j) \quad \text{for } j = 0, 1, \dots, m_0,$$

where $\lambda_0 = \lambda_0(n, \nu) > 0$ is a constant in part (b), i.e. in (2.10), which corresponds to $S = S_0$. For $j = 0$, this inequality is contained in (2.5), because $B_\rho(z_0) = B_\rho(y_{k_1}) \subset B_{1/2}(y_{k_1}) \subset B_1$. Moreover, if (2.12) is true for some j , then the function $v_j := \lambda_0^{-j} v$ satisfies (2.5) with $x_0 = z_j$ and $r = \rho$. By our construction of the sets T_{k_1} and Ω_{k_2} , from (2.11) it follows $S(B_{4\rho}(z_j)) \leq S_0$ for all j . Therefore, we can apply the preceding part (b) of this proof, which yields $v_j \geq \lambda_0$ in $B_{2\rho}(z_j)$. Since $|z_{j+1} - z_j| \leq \rho$, we have $B_\rho(z_{j+1}) \subset B_{2\rho}(z_j)$, hence $v \geq \lambda_0^{j+1}$ on $B_\rho(z_{j+1})$. By induction, (2.12) holds true for all $j = 0, 1, \dots, m_0$. Here m_0 does not exceed a constant m_1 depending only on n, ν , and S . Therefore,

$$v(y) = v(z_{m_0}) \geq \lambda_0^{m_0} \geq \lambda_0^{m_1} =: \lambda = \lambda(n, \nu, S) > 0.$$

Here y is an arbitrary point in ∂B_R . From the inequalities $Lv \leq 0$ in B_R and $v \geq \lambda$ on ∂B_R it follows $v \geq \lambda$ on B_R . Since $R > 2 = 2r$, the desired estimate (2.6) follows. \square

COROLLARY 2.3. *Let $x_0 \in \mathbb{R}^n$, $r > 0$, and a function v in $W(B_r(x_0))$ be such that*

$$(2.13) \quad v \geq 0, \quad Lv \leq 0 \quad \text{in } B_r(x_0); \quad \text{and } v \geq 1 \quad \text{in } B_{\varepsilon r}(x_0),$$

where $0 < \varepsilon \leq 1/2$. Then

$$(2.14) \quad v \geq \varepsilon^\gamma \quad \text{in } B_{r/2}(x_0),$$

where $\gamma = \gamma(n, \nu, S) = -\log_2 \lambda > 0$, and λ is the constant in the previous lemma.

PROOF. We assume $x_0 = 0$. Set $r_j := 2^{-j}r$ for $j = 1, 2, \dots$, and choose natural k such that $2^{-k-1} < \varepsilon \leq 2^{-k}$, so that $r_{k+1} < \varepsilon r \leq r_k$. By our assumptions, $v \geq 1$ on $B_{\varepsilon r} \supset B_{r_{k+1}}$. The previous lemma with $r = r_{k+1}$ yields $v \geq \lambda$ in B_{r_k} . Repeatedly using this lemma again, we get

$$v \geq \lambda^2 \quad \text{in } B_{r_{k-1}}, \quad \dots, \quad v \geq \lambda^k \quad \text{in } B_{r_1} = B_{r/2}.$$

Since $\lambda^k = 2^{-k\gamma} \geq \varepsilon^\gamma$, the corollary is proved. \square

The following technical lemma will help us to deal with functions defined on a ball B_r rather than on a general open set $\Omega \subset \mathbb{R}^n$.

LEMMA 2.4 (Extension lemma). *Let Ω be an open set in \mathbb{R}^n , and let u be a function in $W(\Omega)$, such that*

$$u \geq 0, \quad Lu \geq 0 \quad \text{in } \Omega; \quad \text{and} \quad u = 0 \quad \text{in } (\partial\Omega) \cap B_r,$$

where $B_r := B_r(x_0)$ for some $r > 0$ and $x_0 \in \mathbb{R}^n$. We claim that there are functions $u_\varepsilon \in W(B_r)$ defined for each $\varepsilon > 0$, such that

$$u_\varepsilon \geq 0, \quad Lu_\varepsilon \geq 0 \quad \text{in } B_r; \quad u_\varepsilon \equiv 0 \quad \text{in } \overline{B_r} \setminus \Omega,$$

and $u_\varepsilon \rightarrow u$ as $\varepsilon \rightarrow 0^+$ uniformly on $\Omega \cap B_r$.

PROOF. We partially follow [CS07], pp. 6–7. Fix a standard function

$$0 \leq \eta \in C^\infty(\mathbb{R}^1), \quad \text{such that} \quad \eta(t) \equiv 0 \quad \text{for } |t| \geq 1, \quad \text{and} \quad \int_{\mathbb{R}^1} \eta(t) dt = 1;$$

and set $\eta_\varepsilon(t) := \varepsilon^{-1}\eta(\varepsilon^{-1}t - 2)$ for $\varepsilon > 0$, $t \in \mathbb{R}^1$. These are smooth functions vanishing on $\mathbb{R}^1 \setminus [\varepsilon, 3\varepsilon]$. Further, by repeated integration of η_ε , define the functions $G_\varepsilon \in C^\infty(\mathbb{R}^1)$ satisfying the properties

$$G_\varepsilon \equiv 0 \quad \text{on } (-\infty, \varepsilon], \quad G_\varepsilon'' \equiv \eta_\varepsilon \quad \text{on } \mathbb{R}^1.$$

Now we set

$$u_\varepsilon := G_\varepsilon(u) \quad \text{in } \Omega, \quad u_\varepsilon \equiv 0 \quad \text{on } \overline{B_r} \setminus \Omega.$$

From the properties of η_ε it follows $G_\varepsilon' \geq 0$, $G_\varepsilon'' \geq 0$, and $(u - 3\varepsilon)_+ \leq u_\varepsilon \leq (u - \varepsilon)_+$. Since $u = 0$ on the set $(\partial\Omega) \cap B_r$, the functions u_ε vanish near this set. Hence we have $u_\varepsilon \in W(B_r)$, $u_\varepsilon \geq 0$, and $|u_\varepsilon - u| \leq 3\varepsilon$ on Ω . Finally,

$$Lu_\varepsilon = LG_\varepsilon(u) = G_\varepsilon'(u) \cdot Lu + G_\varepsilon''(u) \cdot a_{ij}D_i u D_j u \geq 0 \quad \text{in } \Omega.$$

Lemma is proved. \square

In comparison with the first growth lemma (Lemma 2.1), the next lemma states that, roughly speaking, from (2.1) with $\mu_1 < 1$ it follows (2.2) with $\beta_1 < 1$.

LEMMA 2.5 (Second growth lemma). *Let Ω be an open set in \mathbb{R}^n , and let $u \in W(\Omega)$, $x_0 \in \mathbb{R}^n$, and $r > 0$ be such that*

$$u \geq 0, \quad Lu \geq 0 \quad \text{in } \Omega; \quad \text{and} \quad u = 0 \quad \text{on } (\partial\Omega) \cap B_{2r}(x_0).$$

We claim that for arbitrary $\mu_2 \in (0, 1)$, there is a constant $\beta_2 = \beta_2(n, \nu, S, \mu_2) \in (0, 1)$, such that from

$$(2.15) \quad |\Omega \cap B_r(x_0)| \leq \mu_2 \cdot |B_r|$$

it follows

$$(2.16) \quad M_r := \sup_{\Omega \cap B_r(x_0)} u \leq \beta_2 \cdot M_{2r}.$$

Replacing Ω by $\Omega \cap B_{2r}(x_0)$, and u by $\text{const} \cdot u$ we can assume that $0 \leq u \leq M_{2r} = 1$ in Ω . Taking

$$v := 1 - u, \quad \mu := 1 - \mu_2 \in (0, 1), \quad \text{and} \quad \beta := 1 - \beta_2 \in (0, 1),$$

we see that this lemma follows from the following one.

LEMMA 2.6. *Let Ω be an open set in \mathbb{R}^n , and let $v \in W(\Omega)$, $x_0 \in \mathbb{R}^n$, and $r > 0$ be such that*

$$v \geq 0, \quad Lv \leq 0 \quad \text{in} \quad \Omega; \quad \text{and} \quad v \geq 1 \quad \text{on} \quad (\partial\Omega) \cap B_{2r}(x_0).$$

We claim that for arbitrary $\mu \in (0, 1)$, there is a constant $\beta = \beta(n, \nu, S, \mu) \in (0, 1)$ such that from $|B_r(x_0) \setminus \Omega| \geq \mu \cdot |B_r|$ it follows $v \geq \beta$ on $\Omega \cap B_r(x_0)$.

PROOF. We follow the lines of the proof of Lemma 2.3 in [S80], though some details are different. As before, we assume $x_0 = 0$. Moreover, applying the extension lemma (Lemma 2.4) to the function $u := 1 - v$, we can also assume that the function v belongs to $W(B_{2r})$ and satisfies $v \geq 0$, $Lv \leq 0$ in B_{2r} . One needs to show that from $|B_r \cap \{v \geq 1\}| \geq \mu \cdot |B_r|$ with $0 < \mu < 1$ it follows $v \geq \beta = \beta(n, \nu, S, \mu) \in (0, 1)$ on B_r .

First consider the case

$$(2.17) \quad |B_r \cap \{v \geq 1\}| \geq \mu_0 \cdot |B_r|, \quad \text{where} \quad \mu_0 := 1 - \mu_1(n, \nu, S, 1/2),$$

i.e. μ_1 is the constant in Lemma 2.1 corresponding to $\beta_1 = 1/2$. Then

$$\begin{aligned} |B_r \cap \{u > 0\}| &= |B_r \cap \{v < 1\}| = |B_r| - |B_r \cap \{v \geq 1\}| \\ &\leq (1 - \mu_0) \cdot |B_r| = \mu_1 \cdot |B_r|. \end{aligned}$$

By Lemma 2.1 applied to $\Omega' := \Omega \cap B_r \cap \{u > 0\}$ and $r/2$ in place of r , we get $u \leq 1/2$, and $v = 1 - u \geq 1/2$ on $B_{r/2}$. Now by the doubling property (Lemma 2.2), $v \geq \beta_0 = \beta_0(n, \nu, S) := \lambda/2 > 0$ on B_r . We have proved that from (2.17) it follows $v \geq \beta_0 = \beta_0(n, \nu, S) > 0$ on B_r , so that the estimate $v \geq \beta > 0$ holds true for $\mu \geq \mu_0$ with $\beta = \beta_0$.

Now consider the remaining case when the set $\Gamma := B_r \cap \{v \geq 1\}$ satisfies $\mu \cdot |B_r| \leq |\Gamma| < \mu_0 \cdot |B_r|$. Almost every point x in the set Γ is its density point, which implies that $B_\rho(x) \subset B_r$ and $|\Gamma \cap B_\rho(x)| > \mu_0 \cdot |B_\rho|$ for small $\rho > 0$. One can include $B_\rho(x)$ into a monotone continuous family of balls B^θ , $0 \leq \theta \leq 1$, such that $B^0 = B_\rho(x)$ and $B^1 = B_r$. Then $\varphi(\theta) := |\Gamma \cap B^\theta|/|B^\theta|$ is a continuous function on $[0, 1]$ with the boundary values $\varphi(0) > \mu_0 > \varphi(1)$. Therefore, for some intermediate value $\theta_0 \in (0, 1)$ we have $\varphi(\theta_0) = \mu_0$, and the corresponding ball $B := B^{\theta_0}$ satisfies

$$(2.18) \quad B \subset B_r, \quad |B \cap \Gamma| = |B \cap \{v \geq 1\}| = \mu_0 \cdot |B|.$$

Denote by Γ_1 the union of all the balls B satisfying (2.18). By the previous argument $v \geq \beta_0 > 0$ on Γ_1 . Obviously $|\Gamma \setminus \Gamma_1| = 0$, because Γ_1 contains all the density points of Γ . Further, by the *simple Vitali lemma*, there is a finite collection of *disjoint* balls $B^{(1)}, B^{(2)}, \dots, B^{(m)}$, satisfying (2.18), such that

$$\sum_{j=1}^m |B^{(j)}| \geq c_0 \cdot |\Gamma_1|, \quad \text{where} \quad c_0 = c_0(n) > 0.$$

This implies

$$\begin{aligned} |\Gamma_1 \setminus \Gamma| &\geq \sum_{j=1}^m |B^{(j)} \cap (\Gamma_1 \setminus \Gamma)| = \sum_{j=1}^m |B^{(j)} \setminus \Gamma| \\ &= (1 - \mu_0) \sum_{j=1}^m |B^{(j)}| \geq (1 - \mu_0)c_0 \cdot |\Gamma_1| \geq (1 - \mu_0)c_0 \cdot |\Gamma|, \end{aligned}$$

and

$$(2.19) \quad |B_r \cap \{v \geq \beta_0\}| \geq |\Gamma_1| \geq c_1 \cdot |\Gamma| = c_1 \cdot |B_r \cap \{v \geq 1\}| \geq c_1 \mu \cdot |B_r|,$$

where $c_1 = c_1(n, \nu, S) := 1 + (1 - \mu_0)c_0 > 1$. If the left side is $< \mu_0 \cdot |B_r|$, then we can apply this estimate again with the function $\beta_0^{-1}v$ in place of v , then to $\beta_0^{-2}v$, etc, which gives us

$$|B_r \cap \{v \geq \beta_0^k\}| \geq c_1^k \mu \cdot |B_r| \quad \text{for } k = 1, 2, \dots$$

This iteration stops when the left side becomes $\geq \mu_0 \cdot |B_r|$. Since $c_1^k \mu \leq 1$, we must have $k \leq k_0 = k_0(n, \nu, S, \mu) := \lceil -\ln \mu / \ln c_1 \rceil$. Finally, since the function $\beta_0^{-k_0}v$ satisfies (2.17), it follows $\beta_0^{-k_0}v \geq \beta_0 > 0$ on B_r , and the estimate $v \geq \beta > 0$ on B_r holds true with $\beta = \beta(n, \nu, S, \mu) := \beta_0^{k_0+1} > 1$. Lemma is proved. \square

Hölder regularity of solutions to non-homogeneous equations $Lu = f$ with $f \in L^n$ follows from Theorem 1.1 and Lemma 2.5 in the same way as Theorem IV.2.5 in [K85], or Theorem 4.1 in [S80], are derived from the corresponding statements. The next theorem is quite similar to these results, therefore we formulate it without proof. We only note that the proof uses approximation of u by solutions of equations with regular coefficients, so that some auxiliary boundary value problems have solutions. This part is provided by the approximation Lemma 4.2 below.

THEOREM 2.7. *Let u be a function in $W(B_{2r})$, $r > 0$, such that $Lu = f$ in B_{2r} , where $f \in L^n(B_{2r})$. Then there are constants $\alpha \in (0, 1)$ and $N > 0$, depending only on n, ν , and S , such that*

$$(2.20) \quad \sup_{x, y \in B_r} \frac{|u(x) - u(y)|}{|x - y|^\alpha} \leq \frac{N}{r^\alpha} \cdot \left(\sup_{B_{2r}} |u| + r \cdot \|f\|_{n, B_{2r}} \right).$$

3. Interior and boundary Harnack inequalities

THEOREM 3.1 (Interior Harnack inequality). *Let u be a function in $W(B_{8r})$ satisfying $u > 0$, $Lu := a_{ij}D_{ij}u + b_i D_i u = 0$ in B_{8r} for some $r > 0$. Then*

$$(3.1) \quad \sup_{B_r} u \leq N_1 \cdot \inf_{B_r} u, \quad \text{where } N_1 = N_1(n, \nu, S) \geq 1, \quad S := \int_{B_{8r}} |\mathbf{b}|^n dx.$$

PROOF. We partially follow the proof of Theorem 3.1 in [S80].

Without loss of generality, we assume $r = 1$. Let $\gamma = \gamma(n, \nu, S) > 0$ be the constant in Corollary 2.3. Since $2 - |x| \geq 1$ in the ball $B_1 := B_1(0)$, we have

$$(3.2) \quad \sup_{B_1} u \leq M := \sup_{B_2} (2 - |x|)^\gamma u = (2 - |x_0|)^\gamma u(x_0)$$

for some $x_0 \in B_2$. Further, consider the function

$$u_0 := u - \frac{u(x_0)}{2} \quad \text{in the ball } B_\rho(x_0), \quad \text{where } \rho := \frac{2 - |x_0|}{2}.$$

Since $2 - |x| \geq \rho$ in $B_\rho(x_0)$, we also have

$$\sup_{B_\rho(x_0)} u_0 < \sup_{B_\rho(x_0)} u \leq \sup_{B_\rho(x_0)} \left(\frac{2 - |x|}{\rho} \right)^\gamma u \leq \rho^{-\gamma} M = 2^\gamma u(x_0) = 2^{\gamma+1} u_0(x_0),$$

and

$$\sup_{B_{\rho/2}(x_0)} u_0 \geq u_0(x_0) > \beta_1 \cdot \sup_{B_\rho(x_0)} u_0, \quad \text{where } \beta_1 = \beta_1(n, \nu, S) := 2^{-\gamma-1} > 0.$$

Now we can use Lemma 2.1 in an equivalent form “if (2.2) fails, then (2.1) fails”, with $\Omega := B_8 \cap \{u_0 > 0\}$, $r := \rho/2$, and u_0 in place of u . By this lemma, there is a constant $\mu_1 > 0$ depending only on n, ν , and S , such that

$$(3.3) \quad |B_\rho(x_0) \cap \{u_0 > 0\}| > \mu_1 \cdot |B_\rho|.$$

Next, the function $v := u/u_0(x_0)$ satisfies

$$v > 0, \quad Lv = 0 \quad \text{in } \Omega' := B_8 \cap \{v < 1\} = B_8 \setminus \bar{\Omega},$$

and $v = 1$ on $(\partial\Omega') \cap B_8$. Moreover, by virtue of (3.3),

$$|B_\rho(x_0) \setminus \Omega'| = |B_\rho(x_0) \cap \{v \geq 1\}| = |B_\rho(x_0) \cap \{u_0 \geq 0\}| > \mu_1 \cdot |B_\rho|.$$

Applying Lemma 2.6 to the function v in Ω' , with $r := \rho$, we obtain the estimate

$$v \geq \beta = \beta(n, \nu, S, \mu_1) > 0 \quad \text{in } B_\rho(x_0).$$

By the choice of x_0 and ρ ,

$$u = u(x_0) \cdot v \geq c_1 := \beta \cdot (2\rho)^{-\gamma} M \quad \text{in } B_\rho(x_0).$$

Finally, we apply Corollary 2.3 to the function $c_1^{-1}u$ with $r = 6$ and $\varepsilon := \rho/6 \in (0, 1/6)$. In our case, $B_3(x_0) = B_{r/2}(x_0) \subset B_6(x_0) \subset B_8$, so that the estimate (2.14) implies $c_1^{-1}u \geq \varepsilon^\gamma$ in the ball $B_3(x_0)$, which contains $B_1 := B_1(0)$. Therefore,

$$\inf_{B_1} u \geq \inf_{B_3(x_0)} u \geq \varepsilon^\gamma c_1 = \left(\frac{\rho}{6}\right)^\gamma \cdot \beta \cdot (2\rho)^{-\gamma} M = \frac{\beta M}{12^\gamma}.$$

This estimate together with (3.2) imply the Harnack inequality (3.1) with $N_1 = N_1(n, \nu, S) := 12^\gamma \cdot \beta^{-1} \geq 1$. \square

As a standard consequence of the interior Harnack inequality, we have

THEOREM 3.2 (Liouville). *Let u be a bounded from above or from below function in $W(\mathbb{R}^n) := W^{2,n}(\mathbb{R}^n) \cap C(\mathbb{R}^n)$ such that $Lu := a_{ij}D_{ij}u + b_iD_iu = 0$ in the entire space \mathbb{R}^n , with a_{ij} satisfying (1.2) and $|\mathbf{b}| \in L^n(\mathbb{R}^n)$. Then $u = \text{const}$ in \mathbb{R}^n .*

Indeed, replacing u by $\text{const} \pm u$ is necessary, we can reduce the proof to the case $\inf_{\mathbb{R}^n} u = 0$. In this case, taking the limit in (3.1) as $r \rightarrow \infty$, we get $u \equiv 0$ in \mathbb{R}^n .

The following theorem is a more general form of the interior Harnack inequality, which is convenient for applications.

THEOREM 3.3. *Let Ω be a bounded open set in \mathbb{R}^n , and let u be a function in $W(\Omega)$ satisfying $u > 0$, $Lu = 0$ in Ω . Then for arbitrary constant $\delta > 0$, such that the set $\Omega^\delta := \{x \in \Omega : \text{dist}(x, \partial\Omega) > \delta\}$ is nonempty and connected, we have*

$$(3.4) \quad \sup_{\Omega^\delta} u \leq N_2 \cdot \inf_{\Omega^\delta} u, \quad \text{where } N_2 = N_2(n, \nu, S, \delta/\text{diam } \Omega), \quad S := \int_{\Omega} |\mathbf{b}|^n dx.$$

PROOF. Fix points $x, y \in \Omega^\delta$. One can choose a sequence $x^{(0)}, x^{(1)}, \dots, x^{(m)}$ in Ω^δ , such that

$$x^{(0)} = x, \quad x^{(m)} = y, \quad \text{and} \quad |x^{(k)} - x^{(k-1)}| < \delta/8 \quad \text{for all} \quad k = 1, \dots, m;$$

and m does not exceed a number m_0 which depends only on n and $\delta/\text{diam} \Omega$. Applying Theorem 3.1 to the balls $B_r(x^{(k)})$, $k = 1, \dots, m \leq m_0$, we obtain

$$u(x) = u(x^{(0)}) \leq N_1 u(x^{(1)}) \leq \dots \leq N_1^m u(x^{(m)}) = N_1^m u(y) \leq N_1^{m_0} u(y),$$

where $N_1 = N_1(n, \nu, S)$ is the constant in (3.1). Since the points $x, y \in \Omega^\delta$ can be selected in an arbitrary way, the inequality (3.4) follows with $N_2 := N_1^{m_0}$. \square

We now proceed to the boundary estimates for positive solutions vanishing at the “bottom” of a Lipschitz cylinder. Here ψ is a Lipschitz function on \mathbb{R}^{n-1} with a Lipschitz constant $K \geq 0$, i.e.

$$(3.5) \quad |\psi(x') - \psi(y')| \leq K \cdot |x' - y'| \quad \text{for all} \quad x', y' \in \mathbb{R}^{n-1}.$$

For $r > 0$, denote

$$(3.6) \quad \begin{aligned} Q_r &:= \{x = (x', x_n) \in \mathbb{R}^n : |x'| < r, 0 < x_n - \psi(x') < r\}, \\ \Gamma_r &:= \{x = (x', x_n) \in \mathbb{R}^n : |x'| \leq r, x_n = \psi(x')\} \subset \partial Q_r. \end{aligned}$$

The following theorem contains a Carleson type estimate for equations (1.1) with $|\mathbf{b}| \in L^n$.

THEOREM 3.4 (Boundary Harnack inequality). *Let ψ be a function on \mathbb{R}^{n-1} satisfying the Lipschitz condition (3.5), $\psi(0) = 0$, and let u be a function in $W(Q_{2r})$ for some $r > 0$, such that*

$$(3.7) \quad u > 0, \quad Lu = 0 \quad \text{in} \quad Q_{2r}; \quad \text{and} \quad u = 0 \quad \text{on} \quad \Gamma_{2r},$$

where Q_{2r} and Γ_{2r} are defined according to (3.6). Then

$$(3.8) \quad \sup_{Q_r} u \leq N_3 u(0, r), \quad \text{where} \quad N_3 = N_3(n, \nu, S, K) \geq 1.$$

PROOF. Since all the conditions here are invariant with respect to rescaling (see Remark 1.4), we can assume that $r = 1$.

(a) First we show that there is a constant $\gamma = \gamma(n, \nu, S, K) > 0$, such that

$$(3.9) \quad M := \sup_{Q_2} d^\gamma u \leq N u(P_1), \quad \text{where} \quad d = d(x) := \text{dist}(x, \partial Q_2),$$

the constant $N = N(n, \nu, S, K) \geq 1$, and $P_1 := (0, 1)$.

Fix $x \in Q_2$. The ball of radius $d(x)$ centered at x touches ∂Q_2 at some point y . A simple geometrical consideration shows that there is a smooth curve parameterized by the arc length $C := \{z = z(s), 0 \leq s \leq s_0\} \subset \overline{Q_2}$, connecting the points $y = z(0)$ and $P_1 = z(s_0)$, passing through the point x , i.e. $z(s_1) = x$ for some $s \in (0, s_0]$, and such that

$$(3.10) \quad cs \leq d(z(s)) \leq s \quad \text{for all} \quad s \in [0, s_0],$$

with a constant $c \in (0, 1)$ depending only on K . (This property means that Q_2 is a *John domain*)

Set $\theta := 1 - c/8$. Then for arbitrary $s \in (0, s_0]$ and $t \in [\theta s, s]$, we have

$$|z(t) - z(s)| \leq |t - s| \leq (1 - \theta)s = \frac{cs}{8} \leq \frac{d(z(s))}{8}.$$

From Theorem 3.1, with $r = d(z(s))/8$, it follows

$$(3.11) \quad N_1 u(z(s)) \geq u(z(t)), \quad 0 < \theta s \leq t \leq s \leq s_0.$$

Fix a constant $\gamma = \gamma(n, \nu, S, K) > 0$ such that $1 \geq \theta^\gamma N_1$. Then

$$\varphi(s) := s^\gamma u(z(s)) \geq N_1 (\theta s)^\gamma u(z(s)) \geq (\theta s)^\gamma u(z(\theta s)) = \varphi(\theta s), \quad 0 < s \leq s_0.$$

For each $s \in (0, s_0]$, there is an integer $k \geq 0$, such that $\theta s_0 < \theta^{-k} s \leq s_0$. Using the previous inequalities, including (3.11) with $s = s_0$, we get

$$\varphi(s) \leq \varphi(\theta^{-1} s) \leq \dots \leq \varphi(\theta^{-k} s) \leq s_0^\gamma N_1 u(z(s_0)) = s_0^\gamma N_1 u(P_1).$$

Note that $s_0 \leq c^{-1} d(z(s_0)) \leq c^{-1}$. Therefore, at the point $x = z(s_1)$,

$$d^\gamma u(x) = d^\gamma u(z(s_1)) \leq s_1^\gamma u(z(s_1)) = \varphi(s_1) \leq N u(P_1),$$

where $N := c^{-\gamma} N_1$. Since $x \in Q_2$ can be chosen in an arbitrary way, the estimate (3.9) follows.

(b) Our next step is to prove the estimate

$$(3.12) \quad M_0 := \sup_{Q_2} d_0^\gamma u \leq N_0 M, \quad \text{where} \quad d_0 = d_0(x) := \text{dist}(x, (\partial Q_2) \setminus \Gamma_2),$$

with a constant $N_0 = N_0(n, \nu, S, K) \geq 1$. Note that $d_0^\gamma u = 0$ on ∂Q_2 , hence the *supremum* in (3.12) is attained at some point $x_0 \in Q_2$, i.e. $d_0^\gamma u(x_0) = M_0$. We claim that

$$(3.13) \quad d(x_0) \geq \varepsilon_0 d_0(x_0) \quad \text{with} \quad \varepsilon_0 = \varepsilon_0(n, \nu, S, K) \in (0, 1/4].$$

The constant ε_0 will be specified below. Suppose (3.13) fails, i.e.

$$\rho := d(x_0) < \varepsilon_0 \rho_0, \quad \text{where} \quad \rho_0 := d_0(x_0), \quad 0 < \varepsilon_0 \leq 1/4.$$

Since $4\rho < 4\varepsilon_0 d_0(x_0) < d_0(x_0)$, the intersection $(\partial Q_2) \cap B_{4\rho}(x_0)$ lies in Γ_2 , so that $u = 0$ on this set. Further, the ball $B_\rho(x_0)$ touches ∂Q_2 at some point $y_0 \in \Gamma_2$. By (3.6), Γ_2 is the graph of a Lipschitz function $x_n = \psi(x')$ restricted to $|x'| \leq 2$. It is easy to see that the measure

$$|B_{2\rho}(x_0) \setminus Q_2| \geq |B_\rho(y_0) \setminus Q_2| \geq \mu \rho^n \quad \text{with} \quad \mu = \mu(n, K) \in (0, 1).$$

Now we can apply Lemma 2.5 with $r := 2\rho$ and $\mu_2 := 1 - 2^{-n}\mu$. By this lemma,

$$u(x_0) \leq \sup_{Q_2 \cap B_{2\rho}(x_0)} u \leq \beta \cdot \sup_{Q_2 \cap B_{4\rho}(x_0)} u, \quad \text{where} \quad \beta = \beta(n, \nu, S, K) \in (0, 1).$$

By the triangle inequality,

$$d_0(x) \geq \rho_0 - 4\rho > (1 - 4\varepsilon_0)\rho_0 \quad \text{on} \quad Q_2 \cap B_{4\rho}(x_0).$$

Combining these inequalities, we obtain

$$M = \rho_0^\gamma u(x_0) \leq (1 - 4\varepsilon_0)^{-\gamma} \beta \cdot \sup_{Q_2 \cap B_{4\rho}(x_0)} d_0^\gamma u \leq (1 - 4\varepsilon_0)^\gamma \beta \cdot M.$$

For small enough $\varepsilon_0 = \varepsilon_0(n, \nu, S, K) \in (0, 1)$, the right side is strictly less than M , and we get the desired contradiction.

The above argument proves the estimate (3.13), which in turn implies (3.12) with $N_0 := \varepsilon_0^{-\gamma} \geq 1$ as follows:

$$M_0 = d_0^\gamma u(x_0) \leq \varepsilon_0^{-\gamma} d^\gamma u(x_0) \leq \varepsilon_0^{-\gamma} M.$$

(c) Both “top” and “bottom” portions of ∂Q_2 are graphs of Lipschitz functions $x_n = \psi(x') + c$, with $c = 0$ or 2 . An elementary geometric reasoning shows that $d_0(x) \geq (1 + K^2)^{-1/2}$ on Q_1 . Hence

$$\sup_{Q_1} u \leq (1 + K^2)^{\gamma/2} \sup_{Q_1} d_0^\gamma u \leq (1 + K^2)^{\gamma/2} M_0.$$

This estimate together with (3.12) and (3.9) yields the desired estimate (3.4). Lemma is proved. \square

In the following Theorem 3.6, which is preceded with a technical Lemma 3.5, we deal with ratios u_1/u_2 of positive solutions. Note that only the numerator u_1 vanishes on Γ_{2r} , while u_2 is just a positive solution. In particular, in the case $u_2 \equiv 1$, Theorem 3.6 is reduced to Theorem 3.4.

Let $\psi = \psi(x')$ be a function on \mathbb{R}^{n-1} satisfying the Lipschitz condition (3.5), and $\psi(0) = 0$. For $r > 0$ and $h \geq 0$, denote

$$\begin{aligned} Q_{r,h} &:= \{x = (x', x_n) \in \mathbb{R}^n : |x'| < r, \quad 0 < x_n - \psi(x') < h\}, \\ Q_{r,h}^+ &:= \{x = (x', x_n) \in \mathbb{R}^n : |x'| < r, \quad h/2 < x_n - \psi(x') < h\}, \\ \Gamma_{r,h} &:= \{x = (x', x_n) \in \mathbb{R}^n : |x'| \leq r, \quad x_n = \psi(x') + h\}, \\ S_{r,h} &:= \{x = (x', x_n) \in \mathbb{R}^n : |x'| = r, \quad 0 < x_n - \psi(x') < h\}. \end{aligned} \tag{3.14}$$

Comparing these notations with (3.6), we see that $Q_r = Q_{r,r}$, $\Gamma_r = \Gamma_{r,0}$. For $r, h > 0$, the boundary $\partial Q_{r,h}$ of the “cylinder” $Q_{r,h}$ is the union of three disjoint sets: the “top” $\Gamma_{r,h}$, the “bottom” Γ_r , and the “lateral side” $S_{r,h}$. If $\psi \equiv 0$, this terminology is understood in the usual sense.

LEMMA 3.5. *Let w be a function in $W(Q_{r,h})$ for some $0 < h \leq r$, such that*

$$Lw = 0 \quad \text{in } Q_{r,h}, \quad w \geq 0 \quad \text{on } \Gamma_r, \tag{3.15}$$

and

$$\inf_{\Gamma_{r,h}} w \geq \sup_{S_{r,h}} (-w)_+. \tag{3.16}$$

We claim that there is a constant $\varepsilon_1 = \varepsilon_1(n, \nu, S, K) \in (0, 1/4]$, such that from $h \leq \varepsilon_1 r$ it follows

$$w(0, x_n) \geq 0 \quad \text{for } 0 \leq x_n \leq h. \tag{3.17}$$

PROOF. Without loss of generality, we assume $r = 1$, and the left side of (3.16) is equal to 1. Then

$$w \geq 1 \quad \text{on } \Gamma_{1,h}, \quad w \geq 0 \quad \text{on } \Gamma_1, \quad \text{and } w \geq -1 \quad \text{on } S_{1,h}. \tag{3.18}$$

(a) Consider the function

$$u := -w \quad \text{on the set } \Omega := Q_{1,h} \cap \{w < 0\}.$$

Obviously, it satisfies

$$0 < u \leq 1, \quad Lu = 0 \quad \text{in } \Omega; \quad u = 0 \quad \text{on } (\partial\Omega) \cap Q_{1,h}.$$

For each $x_0 \in \Omega \cap S_{3/4,h}$, the measure

$$|\Omega \cap B_{1/4}(x_0)| \leq |Q_{1,h}| \leq N_0 \cdot h \quad \text{with } N_0 = N_0(n) > 0,$$

so that we can apply Lemma 2.1 with $x_0 \in \Omega \cap S_{3/4,h}$ and $r = 1/8$ to the function $u := -w$. Since also $u \leq 0$ on the remaining part of $\partial Q_{3/4,h}$, we obtain the estimate

$$(3.19) \quad \sup_{Q_{3/4,h}} (-w)_+ = \sup_{S_{3/4,h}} (-w)_+ \leq \beta_1 = \beta_1(n, \nu, S, h) \rightarrow 0^+ \quad \text{as } h \rightarrow 0^+.$$

(b) Next, set $w_1 := w + \beta_1$. By the maximum principle, (3.18), and (3.19), it follows

$$w_1 \geq 0, \quad Lw_1 = 0 \quad \text{in } Q_{3/4,h}; \quad w_1 \geq 0 \quad \text{on } \Gamma_{3/4}, \quad w_1 \geq 1 \quad \text{on } \Gamma_{3/4,h}.$$

For an arbitrary $x_0 = (x'_0, x_{0n}) \in \Gamma_{1/2,h}$ and $\rho := (1 + K^2)^{-1/2}h/2$, the intersection $(\partial Q_{3/4,h}) \cap B_{2\rho}(x_0)$ lies in the set $\Gamma_{3/4,h}$, which is the graph of a Lipschitz function. Therefore, the measure $|B_\rho(x_0) \setminus Q_{3/4,h}| \geq \mu \cdot |B_\rho|$ with a constant $\mu = \mu(n, K) \in (0, 1)$. Using Lemma 2.6 with $Q_{3/4,h}$, w_1 , ρ in place of Ω , v , r respectively, we get the estimate

$$w_1 \geq \beta = \beta(n, \nu, S, K) > 0 \quad \text{on } Q_{3/4,h} \cap B_\rho(x_0).$$

Further, by Theorem 3.3, applied to the function w_1 on the intersection $Q_{3/4,h} \cap \{|x' - x'_0| < \rho\}$, it follows

$$w_1(x_0 - t\mathbf{e}_n) \geq \beta_0 = \beta_0(n, \nu, S, K) > 0 \quad \text{for } 0 \leq t \leq \frac{h}{2}.$$

Since x_0 is an arbitrary point in $\Gamma_{1/2,h}$, we get the estimate

$$(3.20) \quad w_1 \geq \beta_0 > 0 \quad \text{in } Q_{1/2,h}^+.$$

(c) It is important that the constant β_0 in (3.20) does not depend on h . By virtue of (3.19), one can choose the constant $\varepsilon_1 = \varepsilon_1(n, \nu, s, K) \in (0, 1/4]$ in such a way that from $h \in (0, \varepsilon_1]$ it follows $2\beta_1 \leq \beta_0$. Then the estimates (3.20) and (3.19) imply

$$(3.21) \quad \inf_{Q_{1/2,h}^+} w = \inf_{Q_{1/2,h}^+} w_1 - \beta_1 \geq \beta_1 \geq \sup_{Q_{1/2,h}^+} (-w)_+.$$

Note that $\partial Q_{1/2,h}^+$ contains the set $\Gamma_{1/2,h/2} = \Gamma_{r/2,h/2}$, and $Q_{1/2,h}$ contains $S_{1/2,h/2} = S_{r/2,h/2}$. Therefore

$$\inf_{\Gamma_{r/2,h/2}} w \geq \sup_{S_{r/2,h/2}} (-w)_+.$$

This simply means that the inequality (3.16) remains true with r, h being replaced by $r/2, h/2$. Iterating this procedure, we can replace r, h by $2^{-k}r, 2^{-k}h$ for $k = 1, 2, \dots$. Correspondingly, the first inequality in (3.21) implies

$$w \geq 0 \quad \text{on } Q_{2^{-k-1}r, 2^{-k}h}^+ \quad \text{for } k = 0, 1, 2, \dots,$$

and (3.17) follows by continuity of w . Lemma is proved. \square

THEOREM 3.6 (Comparison theorem). *Let ψ be a function on \mathbb{R}^{n-1} satisfying the Lipschitz condition (3.5), $\psi(0) = 0$, and let u_1 and u_2 be functions in $W(Q_{3r})$, $r > 0$, such that*

$$(3.22) \quad u_{1,2} > 0, \quad Lu_{1,2} = 0 \quad \text{in } Q_{3r}; \quad \text{and } u_1 = 0 \quad \text{on } \Gamma_{3r}.$$

Then

$$(3.23) \quad \sup_{Q_r} \frac{u_1}{u_2} \leq N_4 \cdot \frac{u_1(0, r)}{u_2(0, r)}, \quad \text{where } N_4 = N_4(n, \nu, S, K) \geq 1.$$

PROOF. Multiplying $u_{1,2}$ by appropriate constants if necessary, we can assume $u_1(0, r) = u_2(0, r) = 1$. By Theorem 3.4

$$(3.24) \quad u_1 \leq N_3 \quad \text{in} \quad Q_r,$$

where $N_3 = N_3(n, \nu, S, K) \geq 1$ is the constant in this theorem. Then set $h := \varepsilon_1 r$, where $\varepsilon_1 = \varepsilon_1(n, \nu, S, K) \in (0, 1)$ is the constant in the previous lemma. Applying the Harnack inequality, Theorem 3.3, to the function u_2 , we obtain

$$(3.25) \quad u_2 \geq c_0 = c_0(n, \nu, S, K) > 0 \quad \text{on} \quad Q_r \setminus Q_{r,h}.$$

Finally, we set

$$N_4 := 2N_3/c_0, \quad \text{and} \quad w := N_4 u_2 - u_1.$$

Then

$$w \geq N_4 c_0 - N_3 \geq N_3 \quad \text{on} \quad Q_r \setminus Q_{r,h}, \quad \text{and} \quad -w \leq u_1 \leq N_3 \quad \text{in} \quad Q_r.$$

Therefore, the function w satisfies all the assumption of the previous lemma, which implies $w(0, x_n) \geq 0$ for $0 \leq x_n \leq r$.

This construction remains valid if we move the origin $0 \in \mathbb{R}^n$ to any point $y = (y', y_n) \in \Gamma_r$. Under this translation, the set Q_{2r} will be replaced by

$$Q_{2r}(y) := \{x = (x', x_n) \in \mathbb{R}^n : |x' - y'| < 2r, \quad 0 < x_n - \psi(x') < 2r\},$$

which is a subset of Q_{3r} . As a result, we have $w \geq 0$, or equivalently, $u_1/u_2 \leq N_4$, on the whole set Q_1 . Theorem is proved. \square

COROLLARY 3.7. *Let u_1 and u_2 be functions in $W(Q_{3r})$, $r > 0$, satisfying (3.22), and in addition, $u_2 = 0$ on Γ_{3r} . Then*

$$(3.26) \quad \sup_{Q_r} \frac{u_1}{u_2} \leq N_4^2 \cdot \inf_{Q_r} \frac{u_1}{u_2},$$

where $N_4 = N_4(n, \nu, S, K) \geq 1$ is the constant in (3.23).

PROOF. Since both functions u_1 and u_2 vanish on Γ_{3r} , we can interchange u_1 and u_2 in (3.23), so that

$$\left(\inf_{Q_r} \frac{u_1}{u_2} \right)^{-1} = \sup_{Q_r} \frac{u_2}{u_1} \leq N_4 \cdot \frac{u_2(0, r)}{u_1(0, r)}.$$

Multiplying both sides of this inequality by the corresponding sides of (3.23), we get the desired estimate (3.26). \square

4. Boundary Hopf-Oleinik estimates

In a particular case $\psi \equiv 0$ and $b \equiv 0$, the function $u_2(x) = u_2(x', x_n) := x_n$ satisfies $Lu = 0$ and vanishes on the $\{x_n = 0\}$. In this case $Q_r := \{|x'| < r, 0 < x_n < r\}$, and the estimate (3.26) provides both upper and lower bounds for the ratio $u_1(x)/x_n$ near the origin in \mathbb{R}^n . In 1952, the lower bounds of such kind for solutions of uniformly elliptic equations $Lu = 0$ with $|\mathbf{b}| \in L^\infty$ were independently obtained by E. Hopf [H52] and O.A. Oleinik [O52]. They considered domains Ω satisfying the *interior sphere condition* at a point $y_0 \in \partial\Omega$, i.e. there exists a ball $B_r(x_0) \subset \Omega$ such that $(\partial\Omega) \cap \partial B_r(x_0) = \{y_0\}$. The Hopf-Oleinik estimates state that for any function $u \in C^2(\Omega) \cap C(\bar{\Omega})$ satisfying $u > 0$, $Lu \leq 0$ in Ω , and $u(y_0) = 0$, and any vector $\mathbf{v} \in \mathbb{R}^n$ such that $(\mathbf{v}, x_0 - y_0) > 0$, we have $t^{-1}u(y_0 + t\mathbf{v}) \geq \text{const} > 0$ for small $t > 0$. See the books by M.H. Protter and H.F. Weinberger [PW67], and by D. Gilbarg and N.S. Trudinger [GT83], for further references on this subject.

Here we only mention that in a majority of sources, such kind of estimates are obtained by means of more or less standard *barrier technique*, which requires the boundedness (at least locally) of coefficients.

Our approach uses special iterations based on Theorem 1.1 and Lemma 2.1. It allows us to derive a Hopf-Oleinik type estimate in the case $|\mathbf{b}| \in L^q, q > n$. On the other hand, this estimate fails if $|\mathbf{b}| \in L^n$, as the following example shows.

EXAMPLE 4.1. Consider the functions

$$u_1(x) := \frac{x_n}{|\ln|x||} \quad \text{and} \quad u_2(x) := x_n \cdot |\ln|x||$$

in the cylinder $Q := \{x = (x', x_n) \in \mathbb{R}^n : |x'| < 1/2, 0 < x_n < 1/2\}$, extended as $u_1 = u_2 = 0$ on $(\partial Q) \cap \{x_n = 0\}$. Each of these two functions can be considered as a positive solution to the equation

$$\Delta u + (b, Du) := \Delta u + b_i D_i u = 0 \quad \text{in } Q,$$

where the vector function $b := -\Delta u \cdot |Du|^{-2} Du$ satisfies

$$|\mathbf{b}| = \frac{|\Delta u|}{|Du|} \leq \frac{\text{const}}{|x| \cdot |\ln|x||} \in L^n(Q) \quad \text{for } n \geq 2.$$

However, in any neighborhood of $0 \in \mathbb{R}^n$, $\inf(u_1/x_n) = 0$ and $\sup(u_2/x_n) = \infty$.

In fact, if $|\mathbf{b}| \in L^q$ with $q > n$, then any positive solution of the equation $Lu = 0$ in Q_{2r} vanishing on Γ_{2r} (we follow notations (3.6) with $\psi \equiv 0$) satisfies a two-sided estimate

$$(4.1) \quad 0 < \inf_{Q_r} \frac{u}{x_n} \leq \sup_{Q_r} \frac{u}{x_n} < \infty.$$

Here both the upper estimate $\sup(u/x_n) < \infty$, and the lower estimate $\inf(u/x_n) > 0$ are contained the in paper by O.A. Ladyzhenskaya and N.N. Ural'tseva [LU88], Lemmas 2.3 and 4.4 correspondingly). In Theorem 4.3 below, the lower bound is extended to a bit more general case $|\mathbf{b}| \in L^n, b_n \in L^q$ with $q > n$. The upper estimate can be obtained similarly, with some simplifications. The following lemma helps to reduce the proofs of such kind of results to the case when the coefficients of L are smooth.

LEMMA 4.2 (Approximation lemma). *Let Ω be a bounded open set in \mathbb{R}^n satisfying an interior cone condition, i.e. there a constants $K > 0$ and $r_0 > 0$ such that for each $y \in \partial\Omega$, there is a Cartesian coordinate system centered at the point $y = 0$, such that the cone*

$$(4.2) \quad C := \{x = (x', x_n) \in \mathbb{R}^n : K \cdot |x'| < x_n < Kr_0\} \subset \Omega^c := \mathbb{R}^n \setminus \Omega.$$

Let u be a function in $W^{2,n}(\Omega) \cap C(\bar{\Omega})$ satisfying the inequality

$$(4.3) \quad Lu := a_{ij} D_{ij} u + b_i D_i u \leq f \quad \text{in } \Omega,$$

where a_{ij} satisfy (1.2), $|\mathbf{b}| \in L^n(\Omega), f \in L^n(\Omega)$.

We claim that there are approximations of functions a_{ij}, b_i, f by functions $a_{ij}^\varepsilon, b_i^\varepsilon, f^\varepsilon \in C^\infty(\bar{\Omega}), \varepsilon > 0$, such that a_{ij}^ε satisfy (1.2),

$$(4.4) \quad a_{ij}^\varepsilon \rightarrow a_{ij} \quad \text{a.e. in } \Omega; \quad b_i^\varepsilon \rightarrow b_i, \quad f^\varepsilon \rightarrow f \quad \text{in } L^n(\Omega)$$

as $\varepsilon \rightarrow 0^+$, and solutions $u^\varepsilon \in C^\infty(\Omega) \cap C(\bar{\Omega})$ of the Dirichlet problems

$$(4.5) \quad L^\varepsilon u^\varepsilon := a_{ij}^\varepsilon D_{ij} u^\varepsilon + b_i^\varepsilon D_i u^\varepsilon = f^\varepsilon \quad \text{in } \Omega, \quad u^\varepsilon = u \quad \text{on } \partial\Omega,$$

satisfy

$$(4.6) \quad \sup_{\Omega} (u^{\varepsilon} - u) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0^+.$$

The solvability of the problems (4.5) in Lipschitz domains for equations with smooth, or even Hölder or just continuous, coefficients is well known; see e.g. [M67], Theorem 3.

PROOF. (a) We first consider the case $|\mathbf{b}| \in L^{\infty}(\Omega)$. The coefficients a_{ij} are defined on the whole space \mathbb{R}^n , and also b_i, f can be extended from Ω to \mathbb{R}^n by setting $b_i, f \equiv 0$ on $\mathbb{R}^n \setminus \Omega$. One can approximate a_{ij}, b_i, f by convolutions $a_{ij}^{\varepsilon} := a_{ij} * \eta^{\varepsilon}$, $b_i^{\varepsilon} := b_i * \eta^{\varepsilon}$, $f^{\varepsilon} := f * \eta^{\varepsilon}$ with standard kernels η^{ε} , $\varepsilon > 0$, satisfying

$$0 \leq \eta^{\varepsilon} \in C^{\infty}(\mathbb{R}^n), \quad \eta^{\varepsilon} \equiv 0 \quad \text{on } \mathbb{R}^n \setminus B_{\varepsilon}, \quad \text{and} \quad \int_{\mathbb{R}^n} \eta^{\varepsilon}(x) dx = 1.$$

Then $a_{ij}^{\varepsilon}, b_i^{\varepsilon}, f^{\varepsilon} \in C^{\infty}(\mathbb{R}^n)$, a_{ij}^{ε} satisfy (1.2), and

$$(4.7) \quad a_{ij}^{\varepsilon} \rightarrow a_{ij}, \quad b_i^{\varepsilon} \rightarrow b_i \quad \text{as } \varepsilon \rightarrow 0^+ \quad \text{a.e. in } \Omega.$$

This convergence follows from the properties of the Lebesgue sets (see [St70], Sec. I.1.8). Moreover, $f^{\varepsilon} \rightarrow f$ in $L^n(\mathbb{R}^n)$ as $\varepsilon \rightarrow 0^+$ (see [GT83], Sec. 7.2). Having in mind the application of Theorem 1.1 to $u^{\varepsilon} - u$, we need to estimate from below the functions

$$L^{\varepsilon}(u^{\varepsilon} - u) = L^{\varepsilon}u^{\varepsilon} - Lu + (L - L^{\varepsilon})u \geq F^{\varepsilon} := f^{\varepsilon} - f + (L - L^{\varepsilon})u.$$

We have

$$(L - L^{\varepsilon})u := (a_{ij} - a_{ij}^{\varepsilon})D_{ij}u + (b_i - b_i^{\varepsilon})D_i u \rightarrow 0 \quad \text{in } L^n(\Omega) \quad \text{as } \varepsilon \rightarrow 0^+.$$

(Here we use our additional assumption $b_i \in L^{\infty}$). By Theorem 1.1, from the property (4.7) it follows:

$$\sup_{\Omega} (u^{\varepsilon} - u) \leq N \cdot \|F^{\varepsilon}\|_{n,\Omega} \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0^+.$$

(b) In the general case $|\mathbf{b}| \in L^n(\Omega)$, fix a small constant $\delta > 0$, and choose an open subset $\Omega_{\delta} \subset \Omega$, such that $|\mathbf{b}| \in L^{\infty}(\Omega \setminus \overline{\Omega_{\delta}})$, and the norms in $L^n(\Omega_{\delta})$,

$$\|a_{ij}D_{ij}u\|_{n,\Omega_{\delta}}, \quad \|b_i\|_{n,\Omega_{\delta}}, \quad \|f\|_{n,\Omega_{\delta}} \leq \delta.$$

Define the functions \bar{b}_i, \bar{f} as follows

$$\bar{b}_i \equiv 0, \quad \bar{f} := a_{ij}D_{ij}u \quad \text{on } \Omega_{\delta}, \quad \bar{b}_i \equiv b_i, \quad \bar{f} \equiv f \quad \text{on } \Omega \setminus \Omega_{\delta}.$$

The inequality (4.3) remains valid with the modified b_i and f :

$$\bar{L}u := a_{ij}D_{ij}u + \bar{b}_iD_i u \leq \bar{f}.$$

By the choice of Ω_{δ} , we also have

$$(4.8) \quad \|\bar{b}_i - b_i\|_{n,\Omega} = \|b_i\|_{n,\Omega_{\delta}} \leq \delta, \quad \|\bar{f} - f\|_{n,\Omega} = \|a_{ij}D_{ij}u - f\|_{n,\Omega_{\delta}} \leq 2\delta.$$

Since $\bar{b}_i \in L^{\infty}$, we can apply the argument in (a) correspondingly with

$$b_i^{\delta,\varepsilon} := \bar{b}_i * \eta^{\varepsilon}, \quad f^{\delta,\varepsilon} := \bar{f} * \eta^{\varepsilon}, \quad u^{\delta,\varepsilon}, \quad \text{in place of } b_i^{\varepsilon}, f^{\varepsilon}, u^{\varepsilon}.$$

One can go through this construction for each $\delta_k := 2^{-k}$, $k = 1, 2, \dots$. Therefore, there is a sequence $\varepsilon_k \searrow 0$ such that

$$\|b_i^{\delta_k,\varepsilon} - \bar{b}_i\|_{n,\Omega} \leq \delta_k, \quad \|f^{\delta_k,\varepsilon} - \bar{f}\|_{n,\Omega} \leq \delta_k; \quad \text{and} \quad \sup_{\Omega} (u^{\delta_k,\varepsilon} - u) \leq \delta_k$$

for all $\varepsilon \in (0, \varepsilon_k]$. Here \bar{b}_i and \bar{f} are defined above for $\delta = \delta_k$. Together with (4.8), the first two of these inequalities imply

$$\|b_i^{\delta_k, \varepsilon} - b_i\|_{n, \Omega} \leq 2\delta_k, \quad \|f^{\delta_k, \varepsilon} - f\|_{n, \Omega} \leq 3\delta_k.$$

Finally, for each $\varepsilon > 0$, we take $a_{ij}^\varepsilon := a_{ij} * \eta^\varepsilon$, as in the part (a), and define $b_i^\varepsilon, f^\varepsilon$ in two steps: (i) find k from the relations $\varepsilon_{k+1} < \varepsilon \leq \varepsilon_k$; then (ii) take $b_i^\varepsilon := b_i^{\delta_k, \varepsilon}, f^\varepsilon := f^{\delta_k, \varepsilon}$. This choice of $a_{ij}^\varepsilon, b_i^\varepsilon, f^\varepsilon$ satisfies all the requirements of our lemma. Lemma is proved. \square

THEOREM 4.3. *Let Ω be an open set in \mathbb{R}^n , and let u be a function in $W(\Omega)$ satisfying $u > 0$ and $Lu = 0$ in Ω . Let $y_0 \in \partial\Omega$ and $r > 0$ be such that in a Cartesian coordinate system centered at $y_0 = 0$, the cylinder*

$$(4.9) \quad Q_{2r} := B'_{2r} \times (0, 2r) = \{x = (x', x_n) \in \mathbb{R}^n : |x'| < 2r, 0 < x_n < 2r\}$$

is contained in Ω , and $u(0) = 0$. Suppose that

$$(4.10) \quad S := \int_{Q_{2r}} |\mathbf{b}|^n dx < \infty, \quad \text{and} \quad S_1 := r^{q-n} \int_{Q_{2r}} |b_n|^q dx < \infty,$$

where $q = \text{const} > n$. Then

$$(4.11) \quad x_n^{-1}u(0, x_n) \geq c_1 \cdot r^{-1}u(0, r) \quad \text{for} \quad 0 < x_n \leq r,$$

where $c_1 = c_1(n, \nu, S, S_1, q) \in (0, 1]$.

REMARK 4.4. This theorem provides a Hopf-Oleinik type estimate for equations with $|\mathbf{b}| \in L^q, q > n$. One can modify this formulation in different directions.

(i) One can replace the condition $Q_{2r} \subset \Omega$ with $y_0 \in (\partial Q_{2r}) \cap (\partial\Omega)$ by the interior sphere condition $B_r \subset \Omega$ with $y_0 \in (\partial B_r) \cap (\partial\Omega)$, in both cases $u(y_0) = 0$. One case is reduced to another by an appropriate C^2 -transformation of coordinates.

(ii) Using Lemma 4.2, and the comparison principle, one can replace the equality $Lu = 0$ in Ω by the inequality $Lu \leq 0$ in Ω .

(iii) By the interior Harnack inequality, from (4.11) it follows that for any vector $\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{R}^n$ such that $v_n > 0$, we have $t^{-1}u(t\mathbf{v}) \geq \text{const} > 0$ for small $t > 0$.

PROOF OF THEOREM 4.3. All the quantities in the above formulation are invariant with respect to transformations $x \rightarrow \text{const} \cdot x$ and $u \rightarrow \text{const} \cdot u$. Without loss of generality, we assume $r = 1$ and $u(0, 1) = 1$. By the previous lemma, $u \geq \limsup_{\varepsilon \rightarrow 0^+} u^\varepsilon$, where u^ε are solutions to the Dirichlet problem (4.5) with coefficients $a_{ij}^\varepsilon, b_i^\varepsilon \in C^\infty(\bar{\Omega})$, and $f^\varepsilon \equiv 0$. Therefore, we can also assume $a_{ij}, b_i \in C^\infty(\bar{\Omega})$.

Set $\delta := 1 - n/q \in (0, 1)$, $\alpha := \delta/2$, and for $k = 1, 2, \dots, r_k := 2^{-k}, h_k := r_k^{1+\alpha}$,

$$\Omega_k := Q_{r_k} := B'_{r_k} \times (0, r_k), \quad \Omega_k^+ := \Omega_k \cap \{x_n > h_k\}, \quad \Omega_k^- := \Omega_k \cap \{x_n < h_k\}.$$

By the Harnack inequality, the quantities

$$(4.12) \quad m_k := \inf_{\Omega_k^+} x_n^{-1}u > 0, \quad k = 1, 2, \dots,$$

are estimated from below by a positive constants depending on the prescribed quantities and k . Our goal is to eliminate the dependence on k . From the definition of m_k it follows

$$v_k := m_k x_n - u \leq 0 \quad \text{on} \quad \bar{\Omega}_k \cap (\{x_n = 0\} \cup \{x_n = h_k\}).$$

We can "split" v_k into two functions: $v_k = w_k + z_k$ on Ω_k^- , where w_k and z_k are solutions of the problems

$$(4.13) \quad \begin{aligned} Lw_k = Lv_k = m_k b_n & \quad \text{in } \Omega_k^-, & w_k = 0 & \quad \text{on } \partial\Omega_k^-; \\ Lz_k = 0 & \quad \text{in } \Omega_k^-, & z_k = v_k & \quad \text{on } \partial\Omega_k^-. \end{aligned}$$

These problems have classical solutions because of our smoothness assumptions on the coefficients a_{ij}, b_i . By Theorem 1.1,

$$\sup_{\Omega_k^-} w_k \leq Nr_k m_k \cdot \|b_n\|_{n, \Omega_k^-} \quad \text{with } N = N(n, \nu, S).$$

The last factor is estimated by Hölder's inequality ($1/n = 1/q + 1/p$):

$$\|b_n\|_{n, \Omega_k^-} \leq \|b_n\|_{q, \Omega_k^-} \cdot \|1\|_{p, \Omega_k^-} \leq NS_1^{1/q} r_k^{n/p}, \quad N = N(n).$$

Since $n/p = 1 - n/q = \delta > 0$, we get

$$(4.14) \quad w_k \leq Nr_k^{1+\delta} m_k \quad \text{in } \Omega_k^-.$$

Here and in the rest of the proof, N denotes different constants depending only on n, ν, S, S_1, q . Our next step is to evaluate, for $0 < \rho \leq r_k$ and fixed k ,

$$M(\rho) := \sup_{D(\rho)} (z_k)_+, \quad \text{where } D(\rho) := B'_\rho \times (0, h_k).$$

Consider the case $M(\rho) > 0$, and $0 < \rho \leq r_k - h_k$. Since $z_k = v_k \leq 0$ on $\{x_n = 0\} \cup \{x_n = h_k\}$, by the maximum principle, the value $M(\rho)$ is attained by z_k on the lateral side of $D(\rho)$:

$$M(\rho) = z_k(x(\rho)) = z_k(x'(\rho), x_n(\rho)), \quad \text{where } |x'(\rho)| = \rho, \quad 0 \leq x_n(\rho) \leq h_k.$$

In a subcase $0 \leq x_n(\rho) \leq h_k/2$, one can apply Lemma 2.5 to the function z_k with $\Omega := \Omega_k^- \cap \{z_k > 0\}$, $x_0 := (x'(\rho), 0)$, and $r := h_k/2$. Obviously, the measure condition (2.15) holds true with $\mu_2 = 1/2$. From this lemma and the maximum principle, it follows

$$M(\rho) = z_k(x(\rho)) \leq \sup_{\Omega \cap B_{h_k/2}(x_0)} z_k \leq \beta \cdot \sup_{\Omega \cap B_{h_k}(x_0)} z_k \leq \beta \cdot M(\rho + h_k),$$

where $\beta = \beta(n, \nu, S) \in (0, 1)$. If $h_k/2 \leq x_n \leq h_k$, then these inequalities remain valid with $x_0 := (x'(\rho), h_k)$.

Let k be large enough, so that the integer part $j := [r_{k+1}/h_k] = [2^{k\alpha-1}] \geq 1$. Then $r_{k+1} + jh_k \leq 2r_{k+1} = r_k$, and iterating the previous estimate, we get

$$M(r_{k+1}) \leq \beta \cdot M(r_{k+1} + h_k) \leq \dots \leq \beta^j \cdot M(r_{k+1} + jh_k) \leq \beta^j M(r_k) \leq \beta^j m_k h_k.$$

Here

$$\beta^j \leq \beta^{-1} \cdot \beta^{2^{k\alpha-1}} = \beta^{-1} \cdot \exp(-cr_k^{-\alpha}), \quad \text{where } c := -(\ln \beta)/2 > 0.$$

This estimate together with (4.14) yield the estimate for $m_k x_n - u =: v_k = w_k + z_k$:

$$(4.15) \quad m_k x_n - u \leq N \cdot \left(r_k^{1+\delta} + \exp(-cr_k^{-\alpha}) \right) \cdot m_k \quad \text{in } D(r_{k+1}).$$

From the definition (4.12) of m_k it follows that if $m_k > m_{k+1}$, then

$$(4.16) \quad m_{k+1} = \inf_{\Delta_k} x_n^{-1} u, \quad \text{where } \Delta_k := (\Omega_{k+1}^+ \setminus \overline{\Omega_k^+}) \subset D(r_{k+1}),$$

Since $x_n \geq h_{k+1} = r_{k+1}^{1+\alpha} = 2^{-1-\alpha} r_k^{1+\alpha}$, and $\delta = 2\alpha$, the properties (4.15) and (4.16) imply

$$m_k - m_{k+1} \leq N \cdot \left(r_k^\alpha + r_k^{-1-\alpha} \exp(-cr_k^{-\alpha}) \right) \cdot m_k.$$

We proved the last inequality in the case $m_k > m_{k+1}$. In the opposite case $m_k \leq m_{k+1}$ it is obvious, so this inequality holds true without any restrictions.

Since the exponential term converges to 0 as $k \rightarrow \infty$ faster than any positive power of r_k , there are large natural numbers k_0 and N_0 , depending only on n, ν, S, S_1, q , such that

$$m_k - m_{k+1} \leq N_0 r_k^\alpha m_k, \quad \text{and} \quad N_0 r_k^\alpha \leq 1/2 \quad \text{for all} \quad k \geq k_0.$$

From $m_{k+1} \geq (1 - N_0 r_k^\alpha) m_k$, together with an elementary inequality $\ln(1-t) \geq -2t$ for $0 \leq t \leq 1/2$, we obtain

$$\ln m_{k+1} - \ln m_k \geq \ln(1 - N_0 r_k^\alpha) \geq -2N_0 r_k^\alpha, \quad k \geq k_0.$$

For any integer $k_1 > k_0$,

$$\ln m_{k_1} - \ln m_{k_0} = \sum_{k=k_0}^{k_1-1} (\ln m_{k+1} - \ln m_k) \geq -2N_0 \sum_{k=k_0}^{\infty} r_k^\alpha =: N_1.$$

This implies the estimate

$$m_k \geq c_0 = c_0(n, \nu, S, S_1, q) > 0 \quad \text{for all} \quad k > k_0.$$

In particular, $x_n^{-1} u(0, x_n) \geq c_0$ for all $x_n \in (0, r_{k_0+1})$. By the Harnack inequality, this estimate holds true for $x_n \geq r_{k_0+1}$ as well, possibly with a different constant, because k_0 depends only on the prescribed quantities. This proves the desired estimate (4.11). \square

REMARK 4.5. In our recent paper [S08], the Oleinik-Hopf type estimate (4.11) was extended to the case when Q_r is represented in the form (3.6) with

$$(4.17) \quad \psi(x') := \psi_0(|x'|), \quad \text{where} \quad \psi_0 \geq 0, \quad \psi'_0 \geq 0, \quad \text{and} \quad \int_0^1 t^{-2} \psi(t) dt < \infty.$$

This condition is sharp in a sense that if the integral in (4.17) diverges, then (4.11) fails. However, in [S08] we assumed $b_i \equiv 0$. In the present paper, we impose the ‘‘complementary’’ conditions $\psi \equiv 0$, and $|\mathbf{b}| \in L^q, q > n$. In our forthcoming work, we plan to combine the results and techniques of these two papers in order to cover the general case: ψ satisfying (4.17), and $|\mathbf{b}| \in L^q, q > n$.

Acknowledgements. The author is thankful to Nicolai V. Krylov for stimulating conversations. A part of this paper was completed during June 2009, when the author took part in the INdAM intensive period in Milan, Italy, organized by Ugo Gianazza, Sandro Salsa, and Vincenzo Vespri. The author thanks the organizers for invitation and useful discussions. Special thanks are due to Nina N. Ural'tseva, who raised some open problems which motivated the whole research related to this paper. Finally, the author wishes to thank the referee for corrections and valuable advices.

References

- [AFT01] H. Aimar, L. Forzani, and R. Toledano, *Hölder regularity of solutions of PDE's: a geometrical view*. Comm. Partial Differential Equations **26** (2001), no. 7–8, 1145–1173.
- [A63] A.D. Aleksandrov, *Uniqueness conditions and estimates for the solution of the Dirichlet problem*, Vestnik Leningrad Univ. **18** (1963), no. 3, 5–29 (in Russian). English transl. in Amer. Math. Soc. Transl. (2) **68** (1968), 89–119.
- [CS07] S. Cho and M.V. Safonov, *Hölder regularity of solutions to second order elliptic equations in non-smooth domains*, Journal of Boundary Value Problems, Special Volume 2007, 24 pp.
- [GT83] D. Gilbarg and N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, 2nd ed., Springer-Verlag, Berlin–Heidelberg–New York–Tokyo, 1983.
- [H52] E. Hopf, *A remark on linear elliptic differential equations of second order*, Proc. Amer. Math. Soc., **3** (1952), 791–793.
- [K85] N.V. Krylov, *Nonlinear Elliptic and Parabolic Equations of Second Order*, Nauka, Moscow, 1985 (in Russian). English transl.: Reidel, Dordrecht, 1987.
- [KS80] N.V. Krylov and M.V. Safonov, *A certain property of solutions of parabolic equations with measurable coefficients*, Izvestia Akad. Nauk SSSR, ser. Matem. **44**(1980), 161–175 (in Russian). English transl. in Math. USSR Izvestija, **16** (1981), 151–164.
- [La67] E.M. Landis, *A new proof of E. De Giorgi's theorem*, Trudy Moskov. Mat. Obshch. **16** (1967), 319–328 (in Russian). English transl. in Trudy Moscow Math. Soc. **1967** (1968).
- [La71] E.M. Landis, *Second Order Equations of Elliptic and Parabolic Type*, “Nauka”, Moscow, 1971 (in Russian). English transl.: Amer. Math. Soc., Providence, RI, 1997.
- [LU85] O.A. Ladyzhenskaya and N.N. Ural'tseva, *Estimates of the Hölder constant for functions satisfying a uniformly elliptic or a uniformly parabolic quasilinear inequality with unbounded coefficients*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI) **147** (1985), 72–94 (in Russian). English transl. in J. Soviet Math. **37** (1987), 837–851.
- [LU88] O.A. Ladyzhenskaya and N.N. Ural'tseva, *Estimates on the boundary of a domain for the first derivatives of functions satisfying an elliptic or parabolic inequality*, Trudy Mat. Inst. Steklov **179** (1988), 102–125 (in Russian). English transl. in Proc. Steklov Inst. Math. **179** (1989), 109–135.
- [M67] K. Miller, *Barriers on cones for uniformly elliptic operators*, Ann. Mat. Pura Appl. (4) **76** (1967), 93–105.
- [NU09] A.I. Nazarov and N.N. Ural'tseva, *Qualitative properties of solutions to elliptic and parabolic equations with unbounded lower-order coefficients*, Preprints of the St. Petersburg Math. Soc. 2009-05, 9 pp.
- [O52] O.A. Oleinik, *On properties of solutions of certain boundary problems for equations of elliptic type*, Mat. Sb. (N. S.) **30** (1952), 695–702 (in Russian).
- [PW67] M.H. Protter and H.F. Weinberger, *Maximim Principles in Differential Equations*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1967.
- [S80] M.V. Safonov, *Harnack inequality for elliptic equations and the Hölder property of their solutions*, Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov (LOMI) **96** (1980), 272–287 (in Russian). English transl. in J. Soviet Math. **21**, no. 5 (1983), 851–863.
- [S08] M.V. Safonov, *Boundary estimates for positive solutions to second order elliptic equations*, arXiv:0810.0522v2 [math.AP], 20 p.
- [St70] E. Stein, *Singular Integrals and Differentiability Properties of Functions*. Princeton. Princeton University Press, 1970.

SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA

E-mail address: safonov@math.umn.edu