

ZEROS OF GENERALIZED KRAWTCHOUK POLYNOMIALS

LAURA CHIHARA* AND DENNIS STANTON**

ABSTRACT. The zeros of generalized Krawtchouk polynomials are studied. Some interlacing theorems for the zeros are given. A new infinite family of integral zeros is given, and it is conjectured that these comprise most of the non-trivial zeros. The integral zeros for two families of q -Krawtchouk polynomials are classified.

1. Introduction.

Given a sequence of orthogonal polynomials $p_n(x)$, it is well-known [5, p. 27] that the zeros of $p_n(x)$ are real, simple and lie inside the interval of orthogonality. The Krawtchouk polynomials are a finite sequence of orthogonal polynomials $p_n(x)$, $0 \leq n \leq N$, whose interval of orthogonality is $[0, N]$. In this paper we shall consider the zeros of Krawtchouk polynomials, and in particular investigate the integral zeros.

The integrality of zeros for orthogonal polynomials has combinatorial importance. If the polynomials are naturally related to an association scheme, then the location of the zeros is critical for combinatorial properties of the scheme. For example, if the scheme has a configuration called a perfect e -code, then the polynomial of degree e has e integral zeros [7, Chapter 5]. As another example, generalized Radon transforms can be defined on association schemes [9]. Such a transform is invertible if and only if the relevant polynomial does not have integral zeros.

The most important association scheme is the Hamming scheme of classical coding theory, and its polynomials are the Krawtchouk polynomials. Thus, the zeros of these polynomials and their q -analogs are important. We give some elementary interlacing properties of the zeros in §3. A new infinite family of integral zeros is given in Theorem 4.6. Numerical evidence supports Conjecture 4.7 that asymptotically these are most of the non-trivial integral zeros. In Theorem 6.1 we show that a family of q -Krawtchouk polynomials never has integral zeros. Finally in §6 we consider a second family of q -Krawtchouk polynomials, and classify its integral zeros.

For the (generalized) Krawtchouk polynomials, we need to introduce the standard notation for (basic) hypergeometric series,

$$(1.1) \quad {}_{r+1}F_r \left(\begin{matrix} a_1 & a_2 & \cdots & a_{r+1}; & x \\ b_1 & b_2 & \cdots & b_r \end{matrix} \right) = \sum_{j=0}^{\infty} \frac{(a_1)_j (a_2)_j \cdots (a_{r+1})_j x^j}{(b_1)_j (b_2)_j \cdots (b_r)_j j!}$$

*Department of Mathematics, St. Olaf College, Northfield, MN 55057.

**School of Mathematics, University of Minnesota, Minneapolis, MN 55455. This work was partially supported by a fellowship from the Sloan Foundation, and by NSF grant DMS:8700995.

where

$$(1.2) \quad (a)_j = a(a+1) \cdots (a+j-1)$$

and

$$(1.3) \quad {}_{r+1}\phi_r \left(\begin{matrix} a_1 & a_2 & \cdots & a_{r+1}; & q, & x \\ b_1 & b_2 & \cdots & b_r \end{matrix} \right) = \sum_{j=0}^{\infty} \frac{(a_1)_j (a_2)_j \cdots (a_{r+1})_j x^j}{(b_1)_j (b_2)_j \cdots (b_r)_j (q)_j}$$

where

$$(1.4) \quad (a)_j = (a; q)_j = (1-a)(1-aq) \cdots (1-aq^{j-1}).$$

2. Preliminaries.

In this section we review the necessary facts about the Krawtchouk polynomials. We refer the reader to [2] or [11] for more details.

We shall use notation for Krawtchouk polynomials which agrees with the Hamming scheme $H(N, q)$. The Krawtchouk polynomial $k_n(x, q, N)$ of degree n in x is orthogonal on $x = 0, 1, \dots, N$ with respect to the measure $\binom{N}{x} (q-1)^x$. We may take $q > 1$, although q is integral for $H(N, q)$. This defines the polynomials up to a normalization constant. They may also be defined by the three-term recurrence relation

$$(2.1) \quad (n+1)k_{n+1}(x) = [(N-n)(q-1) + n - qx]k_n(x) - (q-1)(N-n+1)k_{n-1}(x)$$

for $n = 1, \dots, N-1$, with the initial conditions $k_0(x) = 1$ and $k_1(x) = -qx + N(q-1)$. Their generating function is

$$(2.2) \quad \sum_{n=0}^N k_n(x, q, N) z^n = (1 + (q-1)z)^{N-x} (1-z)^x.$$

From (2.2), it is easy to see

$$(2.3) \quad k_n(x, q, N) = \sum_{j=0}^n (-1)^j (q-1)^{n-j} \binom{N-x}{n-j} \binom{x}{j}$$

and thus deduce

$$(2.4) \quad k_n(0, q, N) = \binom{N}{n} (q-1)^n.$$

Also

$$(2.5) \quad k_n(x, q, N) = k_n(N-x, q/(q-1), N) (1-q)^n.$$

The Krawtchouk polynomials can also be expressed as a ${}_2F_1$

$$(2.6) \quad k_n(x, q, N) = \binom{N}{n} (q-1)^n {}_2F_1 \left(\begin{matrix} -n & -x; & q \\ & -N & q-1 \end{matrix} \right).$$

Because a ${}_2F_1$ is a symmetric function of the two numerator parameters, if $x = i$ is integral we have the self-dual relation

$$(2.7) \quad \binom{N}{i} (q-1)^i k_n(i, q, N) = \binom{N}{n} (q-1)^n k_i(n, q, N) \quad n, i = 0, 1, \dots, N.$$

Clearly the three-term relation (2.1) and the self-dual relation (2.7) imply the difference equation

$$(2.8) \quad (q-1)(N-x)k_n(x+1, q, N) - [(q-1)(N-x) + x - qn]k_n(x, q, N) + xk_n(x-1, q, N) = 0.$$

In later sections we will need the following propositions. Proposition 2.1 follows from the generating function (2.2), and Proposition 2.2 follows from parts (1) and (2) of Proposition 2.1.

Proposition 2.1. For $0 \leq n \leq N$,

- (1) $k_n(x, q, N+1) = k_n(x, q, N) + (q-1)k_{n-1}(x, q, N)$
- (2) $k_n(x, q, N+1) = k_n(x-1, q, N) - k_{n-1}(x-1, q, N)$
- (3) $k_n(x, q, N) - k_n(x-1, q, N) + (q-1)k_{n-1}(x, q, N) + k_{n-1}(x-1, q, N) = 0.$
- (4) $k_n(x, q, N) - k_n(x+1, q, N) = qk_{n-1}(x, q, N-1)$

Proposition 2.2. If $k_r(s, q, N) = 0$, where s is real, then

- (1) $k_r(s, q, N+1) = (q-1)k_{r-1}(s, q, N)$
- (2) $k_{r+1}(s, q, N+1) = k_{r+1}(s, q, N)$
- (3) $k_r(s+1, q, N+1) = -k_{r-1}(s, q, N)$

Finally we need a special proposition for the $q = 2$ case.

Proposition 2.3. We have

$$k_r(x, 2, N) - k_r(x+2, 2, N) = 4k_{r-1}(x, 2, N-2).$$

Proof. From Proposition 2.1(4) we obtain the two equations

$$\begin{aligned} k_r(x, 2, N) - k_r(x+1, 2, N) &= 2k_{r-1}(x, 2, N-1) \\ k_r(x+1, 2, N) - k_r(x+2, 2, N) &= 2k_{r-1}(x+1, 2, N-1) \end{aligned}$$

Adding these two equations together yields

$$\begin{aligned} k_r(x, 2, N) - k_r(x+2, 2, N) &= 2[k_{r-1}(x, 2, N-1) + k_{r-1}(x+1, 2, N-1)] \\ &= 2[-k_r(x+1, 2, N-1) + k_r(x, 2, N-1)] \\ &= 2[2k_{r-1}(x, 2, N-2)] \end{aligned}$$

where the second equality follows from Proposition 2.1(3) and the third equality follows from Proposition 2.1(4). \square

3. Interlacing of the zeros.

In this section we concentrate on properties of the zeros which do not involve integrality. Let $x_{n,1}^N < x_{n,2}^N < \dots < x_{n,n}^N$ denote the zeros of $k_n(x, q, N)$. Recall that $0 < x_{n,1}^N < x_{n,n}^N < N$, there is some integer in any open interval $(x_{n,k}^N, x_{n,k+1}^N)$ [16], and that the zeros of $k_n(x, q, N)$ and $k_{n-1}(x, q, N)$ interlace [16]. First we give an interlacing theorem which is analogous to [10, Th.4] for Hahn polynomials.

Theorem 3.1. *The zeros of $k_n(x, q, N)$ and $k_n(x, q, N + 1)$ interlace,*

$$x_{n,i}^N < x_{n,i}^{N+1} < x_{n,i+1}^N < x_{n,i+1}^{N+1}, \quad \text{for } i = 1, \dots, n-1.$$

Proof. This follows from the interlacing property for $k_n(x, q, N)$ and $k_{n-1}(x, q, N)$ and Proposition 2.1(1). The details are similar to the proof of Theorem 3.2. \square

We now have two zeros, $x_{n,i}^{N+1}$ and $x_{n-1,i}^N$, which lie in the interval $[x_{n,i}^N, x_{n,i+1}^N]$. These zeros also interlace.

Theorem 3.2. *The zeros of $k_{n-1}(x, q, N)$ and $k_n(x, q, N + 1)$ interlace,*

$$x_{n,i}^{N+1} < x_{n-1,i}^N < x_{n,i+1}^{N+1}, \quad \text{for } i = 1, \dots, n-1.$$

Proof. We must show that $x_{n,i}^{N+1} < x_{n-1,i}^N$, so we first consider the $i = 1$ case. Clearly the interlacing property for $k_n(x, q, N)$ and $k_{n-1}(x, q, N)$, and Proposition 2.1(1) imply that $x_{n,1}^{N+1} \neq x_{n-1,1}^N$. So assume that $x_{n-1,1}^N < x_{n,1}^{N+1}$. Because the leading terms of $k_n(x, q, N)$ and $k_n(x, q, N + 1)$ have the same sign, $k_n(x, q, N)$ and $k_n(x, q, N + 1)$ have opposite signs at $x_{n-1,1}^N$. This contradicts Proposition 2.2(2).

The proof continues by induction on i noting that $k_n(x, q, N)$ and $k_n(x, q, N + 1)$ have opposite signs on the interval $(x_{n,i}^N, x_{n,i}^{N+1})$. So the assumption $x_{n-1,i}^N < x_{n,i}^{N+1}$ would contradict Proposition 2.2(2). \square

The next theorem shows that the interlacing in Theorem 3.1 is “close”.

Theorem 3.3. *For $1 \leq i \leq n \leq N$, $x_{n,i}^N - x_{n,i}^{N+1} < 1$.*

Proof. Since $k_n(x_{n,i}^N, q, N) = 0$, Proposition 2.2(1) implies

$$k_n(x_{n,i}^N, q, N + 1) = (q - 1)k_{n-1}(x_{n,i}^N, q, N)$$

and Proposition 2.2(2) implies

$$k_n(x_{n,i}^N + 1, q, N + 1) = -k_{n-1}(x_{n,i}^N, q, N).$$

Thus, we have

$$k_n(x_{n,i}^N, q, N + 1) = -(q - 1)k_n(x_{n,i}^N + 1, q, N + 1),$$

and if $q > 1$, then $k_n(x, q, N + 1)$ must have a zero between $x_{n,i}^N$ and $x_{n,i}^N + 1$. \square

If $q = 2$, we can say more about the distance between consecutive zeros. Levit [10, Th.3] has a similar result for Hahn polynomials.

Theorem 3.4. *Let $x_1 < x_2$ be consecutive zeros of $k_r(x, 2, N)$ and suppose $r < N/2$. Then*

$$x_1 + 2 < x_2.$$

Proof. First we note that the difference equation (2.8) when $q = 2$ is

$$(3.1) \quad (N - 2r)k_r(x) = (N - x)k_r(x + 1) + xk_r(x - 1).$$

Assume that x_1 is not an integer (the argument for x_1 an integer is similar). Suppose there is only one integer s separating x_1 and x_2 , that is $s - 1 \leq x_1 < s < x_2 \leq s + 1$. (Recall that the open interval between two consecutive zeros of an orthogonal polynomial must contain at least one spectral point.) Then we must have

$$\operatorname{sgn}(k_r(s-1)) = \operatorname{sgn}(k_r(s+1)) = -\operatorname{sgn}(k_r(s))$$

which contradicts (3.1) if $x = s$. This contradiction forces there to be at least two integers in the interval (x_1, x_2) , or $x_2 > x_1 + 1$.

Now suppose $x_2 \leq x_1 + 2$, and let $x = x_1 + 1$ in (3.1). Then

$$(N - 2r)k_r(x_1 + 1) = (N - x_1 - 1)k_r(x_1 + 2)$$

which implies

$$(3.2) \quad \operatorname{sgn}(k_r(x_1 + 1)) = \operatorname{sgn}(k_r(x_1 + 2)).$$

Then (3.2) and $x_1 + 1 < x_2 \leq x_1 + 2$ implies that there must be another zero of $k_r(x)$ between x_2 and $x_1 + 2$. But from the previous paragraph, we know then that $(x_2, x_1 + 2)$ must contain at least two integers, thus giving us four integers in the interval $(x_1, x_1 + 2)$. This is clearly impossible; thus $x_2 > x_1 + 2$. \square

The condition $r < N/2$ in Theorem 3.4 cannot be relaxed because the zeros of $k_N(x, 2, 2N)$ are $1, 3, \dots, 2N - 1$.

There is a discrete form of Markoff's theorem [16, p. 115] which states the following. If the weight function $w(x, q)$ is purely discrete, and the logarithmic derivative $w_q(x, q)/w(x, q)$ is increasing, then the i th of the corresponding orthogonal polynomials is an increasing function of q . For the Krawtchouk polynomials we then have the following theorem.

Theorem 3.5. *Let $x_{i,n}^N(q)$ be the i th zero of $k_n(x, q, N)$. Then $x_{i,n}^N(q)$ is an increasing function of q , for $q > 1$.*

4. Integral zeros.

We next consider the number theoretic conditions which are necessary for a Krawtchouk polynomial to have an integral zero. We also give a non-trivial family of integral zeros in Theorem 4.6.

Theorem 4.1. *If $k_r(s, q, N) = 0$, s an integer, then q divides $\binom{N}{r}$.*

Proof. From Proposition 2.1(4), we have

$$k_r(0, q, N) - k_r(1, q, N) = qk_{r-1}(0, q, N - 1)$$

$$k_r(1, q, N) - k_r(2, q, N) = qk_{r-1}(1, q, N - 1)$$

$$\vdots$$

$$k_r(s-1, q, N) - k_r(s, q, N) = qk_{r-1}(s-1, q, N - 1)$$

The telescoping sum yields

$$k_r(0, q, N) = q \sum_{j=0}^{s-1} k_r(j, q, N - 1).$$

Recalling (2.4) yields the desired conclusion, since $k_r(j, q, N - 1)$ is integral. \square

If q is a prime number, the value of $\binom{N}{r} \pmod{q}$ is well-known from Lucas' Theorem [6, p. 65], and we have the following corollary.

Corollary 4.2. *Let $r_1 \cdots r_a$ and $N_1 \cdots N_b$ be the base q representations of r and N respectively. If q is a prime number and $k_r(s, q, N) = 0$ for some integer s , then for some i , $r_i > N_i$.*

The divisibility condition in Theorem 4.1 can be improved if $q = 2$.

Theorem 4.3. *If s is a positive integer and $k_r(s, 2, N) = 0$, then*

$$k_r(s \pmod{2^t}, 2, N) \equiv 0 \pmod{2^{t+1}},$$

for any integer $t \geq 1$.

Proof. Proposition 2.3 immediately gives the result for $t = 1$. It also shows

$$(4.1) \quad k_r(x, 2, N) - k_r(x + 4, 2, N) = 4[k_{r-1}(x, 2, N - 2) + k_{r-1}(x + 2, 2, N - 2)].$$

If x is integral, then Proposition 2.3 implies that $k_{r-1}(x, 2, N - 2)$ and $k_{r-1}(x + 2, 2, N - 2)$ are the same modulo four, so the left side of (4.1) $\equiv 0 \pmod{8}$. Again a telescoping sum shows that this is the $t = 2$ case. The general case follows from $k_r(x, 2, N) - k_r(x + 2^t, 2, N)$ being the sum of 2^{t-1} terms, all of which are the same modulo four. \square

We next consider the integral zeros of $k_n(x, 2, N)$. By symmetry in n and x we can assume that $n \leq x$. From (2.5), we can clearly take $x \leq N/2$. There is another set of zeros to which we will refer as trivial. For n odd, (2.5) also implies that $x = N/2$ is an integral zero for N even.

For polynomials of small degree n , the non-trivial integral zeros (n, x, N) , $1 \leq n \leq x \leq N/2$, can be found explicitly (see [9]).

Proposition 4.4. *The integral zeros for degrees 1, 2, and 3 are*

- (1) $(1, k, 2k)$, $k \geq 1$,
- (2) $(2, k(k-1)/2, k^2)$, $k \geq 3$,
- (3) $(3, k(3k \pm 1)/2, 3k^2 + 3k + 3/2 \pm (k + 1/2))$, $k \geq 2$.

Strictly speaking, (1) consists of trivial zeros. Note that the pentagonal numbers appear as the integral zeros in (3).

Even though the zeros can be given for degree four, it is a difficult number theoretic problem to classify when these zeros are integral. Graham and Diaconis [9] give non-trivial values of $(4, 7, 17)$, $(4, 10, 17)$, $(4, 30, 66)$, $(4, 36, 66)$, $(4, 715, 1521)$, $(4, 806, 1521)$, $(4, 7476, 15043)$, $(4, 7567, 15043)$, along with the trivial values $(4, 1, 8)$, $(4, 3, 8)$, $(4, 5, 8)$, $(4, 7, 8)$. They conjecture that this is the complete list for degree four. Laurent Habsieger has shown that any other possible value of N must have at least 1000 digits. We can use Theorem 4.3 to restrict the possible values of N . The following corollary takes $t = 2$.

Corollary 4.5. *Suppose $k_4(x, 2, N) = 0$, for some integer x .*

- (1) *If $x \equiv 0 \pmod{4}$, then $N \equiv 0, 1, 2, \text{ or } 3 \pmod{32}$.*
- (2) *If $x \equiv 1 \pmod{4}$, then $N \equiv 1, 2, 3, \text{ or } 8 \pmod{32}$.*
- (3) *If $x \equiv 2 \pmod{4}$, then $N \equiv 0, 2, 3, \text{ or } 17 \pmod{32}$.*
- (4) *If $x \equiv 3 \pmod{4}$, then $N \equiv 2, 3, 8, \text{ or } 17 \pmod{32}$.*

Proof. Use the explicit form of $k_4(i, 2, N) = 0, 0 \leq i \leq 3$, as a polynomial of degree i in N and Theorem 4.3. \square

Using MACSYMA we found all of the non-trivial integral zeros for $q = 2$ and $N \leq 700$. The data suggested the following theorem, for an infinite family of non-trivial zeros.

Theorem 4.6. *For any integer $h \geq 1$,*

$$k_{2h}(4h - 1, 2, 8h + 1) = 0.$$

Proof. This theorem can be proven using the theory of hypergeometric series in [1]. Instead we give a simple proof from the $q = 1$ version of (6.6) below. Let $N = 2n + t$ in (6.6), and assume that x is even. If $t = 3$ the two allowed values of r in (6.6) are $r = x/2$ and $r = x/2 - 1$. If these two terms sum to zero, we find that $x = (n + 1)/2 = 2h$, so that $n = 4h - 1$ and $N = 8h + 1$. \square

Using MACSYMA we found all of the non-trivial integral zeros for $q = 2$ and $N \leq 700$. The following table lists all non-trivial integral zeros for $q = 2, N \leq 700$, and $1 \leq n \leq x \leq N/2$, which do not follow from Proposition 4.4 or Theorem 4.6.

(5, 14, 36)	(23, 31, 67)	(31, 103, 214)	(34, 254, 514)
(5, 22, 67)	(14, 47, 98)	(5, 133, 289)	(84, 286, 576)
(5, 28, 67)	(19, 62, 132)	(6, 155, 345)	
(6, 31, 67)	(61, 86, 177)	(44, 230, 465)	

Note that many of these zeros have $N = 2n + t$, for small values of t . In fact, with Laurent Habsieger we have found five more infinite families of zeros, for $t = 4, 5, 6$, and 8 . These families contain all of the zeros in the table, except for $(5, 22, 67), (5, 28, 67), (5, 133, 289)$, and $(6, 155, 345)$. The details will appear elsewhere. Nevertheless, these zeros are more sparse than those in Theorem 4.6, and we make the following conjecture.

Conjecture 4.7. *The number of non-trivial integral zeros of Krawtchouk polynomials $k_n(x, 2, M)$ for $M \leq N$ is asymptotic to $N/8$.*

One may also look for families of integral zeros of $k_n(x, q, N), q \geq 3$. In this case we may assume $1 \leq n \leq \min(x, N - x)$. Again using MACSYMA this was done for $q \leq 20$ and $N \leq 100$.

Proposition 4.8. *The integral zeros (n, x, N) of $k_n(x, q, N)$ for $n \geq 3, N \leq 100$, and $3 \leq q \leq 20$ are*

- (1) $(3, 14, 28), (5, 16, 30), (4, 35, 57), (4, 65, 93)$, for $q = 3$,
- (2) $(3, 55, 66), (5, 55, 68)$, for $q = 4$,
- (3) $(4, 66, 80)$ for $q = 5$, and
- (4) $(4, 52, 66)$ for $q = 6$.

5. Summation theorems.

In this section we give a few summation theorems for Krawtchouk polynomials which follow from the recurrence relations in §2.

From Proposition 2.1(4) we can deduce the following identity for the Krawtchouk polynomials

$$(5.1) \quad \sum_{j=0}^N k_r(j, q, N) = \frac{1}{q} [(q-1)^{r+1} + (-1)^r] \binom{N+1}{r+1}.$$

This can be generalized for $q = 2$ by using Lemma 4.3 to find if N is even and r odd,

$$(5.2) \quad \sum_{j=0}^{N/2} k_r(2j, 2, N) = 0 = \sum_{j=1}^{N/2} k_r(2j-1, 2, N),$$

if N and r are both even,

$$(5.3) \quad \begin{aligned} \sum_{j=0}^{N/2} k_r(2j, 2, N) &= \frac{N+2}{2(N-r+1)} \binom{N+1}{r+1} \\ \sum_{j=1}^{N/2} k_r(2j-1, 2, N) &= \frac{N-2r}{2(N-r+1)} \binom{N+1}{r+1}, \end{aligned}$$

if N and r are both odd,

$$(5.4) \quad \sum_{j=0}^{\frac{N-1}{2}} k_r(2j, 2, N) = \frac{1}{2} \binom{N+1}{r} = - \sum_{j=1}^{\frac{N+1}{2}} k_r(2j-1, 2, N),$$

and if N is odd and r is even

$$(5.5) \quad \sum_{j=0}^{\frac{N-1}{2}} k_r(2j, 2, N) = \frac{1}{2} \binom{N+1}{r+1} = \sum_{j=1}^{\frac{N+1}{2}} k_r(2j-1, 2, N).$$

From (5.1) and the product formula for Krawtchouk polynomials it is easy to see that

$$(5.6) \quad \sum_{k=0}^N k_r(k, 2, N) k_s(k, 2, N) = \sum_{m=0}^{N/2} \binom{N-2m}{\frac{r+s}{2}-m} \binom{2m}{\frac{r-s}{2}-m} \binom{N+1}{2m+1}.$$

The binomial coefficients in (5.6) which have non-integral arguments are defined to be zero.

6. q -analogs.

There are two families of orthogonal polynomials which have been called q -Krawtchouk polynomials. In this section we shall consider the zeros of these polynomials, and compare the results to §4.

The first family has been called the *affine q -Krawtchouk* polynomials, because it arises from association schemes with natural translations [14, 15],

$$(6.1) \quad K_n^{Aff}(x, a, N, q) = \nu_n {}_3\phi_2 \left(\begin{matrix} q^{-x} & q^{-n} & 0 \\ q^{-N} & a \end{matrix} ; q, q \right), \text{ for } 0 \leq n \leq N,$$

where ν_n is non-zero. Clearly (6.1) implies that $K_n^{Aff}(x, a, N, q)$ is a polynomial in q^{-x} of degree n . The three cases for which these polynomials are realized from association schemes all have q equal to a prime power. They are

- (A1) $K_n^{Aff}(x, q^{-M}, N, q)$, M integral and $N \leq M$,
- (A2) $K_n^{Aff}(x, q^{2\lfloor(N+1)/2\rfloor-1}, \lfloor N/2 \rfloor, q^2)$, and
- (A3) $K_n^{Aff}(x, -(-q)^{-N}, N, -q)$.

Note that the effect of the greatest integer functions in (2) is just to put q^{-N} and q^{1-N} as denominator parameters in the ${}_3\phi_2$. Note also that if ν_n is defined appropriately,

$$\lim_{q \rightarrow 1} K_n^{Aff}(x, a, N, q) = k_n(x, 1/a, N).$$

However this limit does not apply to the three cases above, because the value of a depends upon q .

The other family of q -Krawtchouk polynomials is a q -analog of the *binary* Krawtchouk polynomials (the $q = 2$ case in §2). It is

$$(6.2) \quad K_n(x, c, N, q) = \nu_n {}_3\phi_2 \left(\begin{matrix} q^{-x} & q^{-n} & -q^{n-N-c} \\ q^{-N} & 0 & \end{matrix} ; q, q \right), \text{ for } 0 \leq n \leq N,$$

where ν_n is non-zero. These polynomials arise from six association schemes [13], giving five sets of polynomials

- (B1) $K_n(x, 0, N, q)$
- (B2) $K_n(x, 1, N, q)$
- (B3) $K_n(x, 2, N, q)$
- (B4) $K_n(x, 1/2, N, q^2)$
- (B5) $K_n(x, 3/2, N, q^2)$.

This time

$$\lim_{q \rightarrow 1} K_n(x, c, N, q) = k_n(x, 2, N).$$

if ν_n is defined appropriately. There are also group theoretic reasons [14] for considering these polynomials as the correct q -analogs of $k_n(x, 2, N)$.

First we consider the affine q -Krawtchouk polynomials (A1)-(A3). Surprisingly, the next theorem states that these polynomials are never zero at integral values of x . It had previously been shown [4] that at least one zero must be non-integral.

Theorem 6.1. *Let $p_n(x)$ be one of the affine q -Krawtchouk polynomials (1)-(3), where q is a prime power. Then*

- (1) $p_n(i) \neq 0$ for $i, n = 0, 1, \dots, N$, for (1) and (3),
- (2) $p_n(i) \neq 0$ for $i, n = 0, 1, \dots, \lfloor N/2 \rfloor$, for (2).

Proof. We shall prove case (2), and leave the other two cases to the reader. By symmetry we can assume that $i \geq n \geq 1$ and $p_n(i) = 0$. Multiply the definition (6.1) of $p_n(i)$ by $(q^{-N}; q^2)_n (q^{-N+1}; q^2)_n (q^2; q^2)_n q^{2nN}$ to obtain

$$(6.3) \quad \sum_{j=0}^n (q^{2n}; q^{-2})_j (q^{2i}; q^{-2})_j (q^{N-2j}; q^{-2})_{n-j} (q^{N-1-2j}; q^{-2})_{n-j} \\ \times (q^{2j+2}; q^2)_{n-j} q^{(N-2n)j + (N-2i)j + 2n^2 - n + j} \\ = \sum_{j=0}^n c_j(q) = 0.$$

For each j , $c_j(q)$ is a polynomial in q with integer coefficients. The lowest power of q that appears in $c_j(q)$ is $(N-2n)j + (N-2i)j + 2n^2 - n + j$ and the highest power

of q that appears is $2Nn + n(n+1) - j^2 + j$. Since $i, j \leq \lfloor N/2 \rfloor$, (6.3) implies that the lowest power of q appearing is q^{2n^2-n} . Thus, if we divide both sides by q^{2n^2-n} , we obtain $0 \equiv 1 \pmod{q}$. This contradiction implies that $p_n(i) \neq 0$ for $i \geq n$. \square

Next we turn to the q -Krawtchouk polynomials $K_n(x, c, N, q)$. These polynomials are not symmetric in n and x . We shall need the generating function [13]

$$(6.4) \quad \sum_{x=0}^N K_n(x, c, N, q) z^x = (z)_n (-q^c z)_{N-n},$$

which implies the following q -analog of (2.3) via the q -binomial theorem [1]

$$(6.5) \quad K_n(x, c, N, q) = \sum_{j=0}^x (-1)^j \begin{bmatrix} n \\ j \end{bmatrix}_q \begin{bmatrix} N-n \\ x-j \end{bmatrix}_q q^{\binom{x}{2} + (c-j)(x-j)}.$$

We first give the q -analog of the trivial zeros of x odd for $k_N(x, 2, 2N)$.

Proposition 6.2. *If x is an odd integer satisfying $1 \leq x \leq 2N$, then $K_N(x, 0, 2N, q) = 0$.*

Proof. This is clear from (6.4) and $(z)_N (-z)_N = (z^2; q^2)_N$. \square

Finally, the argument of Theorem 6.1 implies that these polynomials are otherwise non-zero.

Theorem 6.3. *If q is a prime power, the following values of q -Krawtchouk polynomials are non-zero*

- (1) $K_n(i, 0, N, q)$ for $0 \leq i, n \leq N, 2n \neq N$,
- (2) $K_n(i, 1, N, q)$ for $0 \leq i, n \leq N$,
- (3) $K_n(i, 2, N, q)$ for $0 \leq i, n \leq N$,
- (4) $K_n(i, 1/2, N, q^2)$ for $0 \leq i, n \leq N$,
- (5) $K_n(i, 3/2, N, q^2)$ for $0 \leq i, n \leq N$,

Proof. First we take $c = 0$ and assume that $2n < N$, so that the generating function is $(z^2; q^2)_n (-zq^n; q)_{N-2n}$. Then the q -binomial theorem [1] implies

$$(6.6) \quad \begin{aligned} K_n(x, 0, N, q) &= \sum_r \begin{bmatrix} n \\ r \end{bmatrix}_{q^2} q^{2\binom{r}{2}} \begin{bmatrix} N-2n \\ x-2r \end{bmatrix}_q q^{\binom{x-2r}{2} + n(x-2r)} (-1)^r \\ &= \sum_r c_r(q), \end{aligned}$$

where $c_r(q)$ is a polynomial in q with integral coefficients. The degree of the lowest degree term in $c_r(q)$ is

$$L(r) = nx + x^2/2 - x/2 + 3r^2 - 2rx - 2rn.$$

As a function of r , $L(r)$ has its minimum value at $r = (x+n)/3$. Thus there is a unique term in (6.6) containing the smallest power of q , and as in Theorem 6.1, this implies $K_n(x, 0, N, q) \neq 0$. The $2n > N$ case can be done similarly.

The proof for $c = 1$ ($c = 2$) is similar, this time two (three) terms naturally occur in the sum that corresponds to (6.6). Nevertheless, again there is a unique term with the term of minimum degree.

For $c = 1/2$ or $c = 3/2$, we can use (6.5), for which

$$L(r) = x^2/2 - x/2 + r^2 - rx - rc + cx.$$

The minimum value for $L(r)$ occurs at $r = (x + c)/2$. If $c = 1/2$ or $c = 3/2$, this insures a unique term in (6.5) of minimum degree. Note that this argument also does the $c = 0$ case with x even, but not for x odd. \square

REFERENCES

1. W. Bailey, *Generalized Hypergeometric Series*, Cambridge University Press, Cambridge, 1935.
2. E. Bannai and T. Ito, *Algebraic Combinatorics I*, Benjamin/Cummins, Menlo Park California, 1984.
3. M.R. Best, *A contribution to the non-existence of perfect codes*, PhD thesis, Amsterdam, 1982.
4. L. Chihara, *On the zeros of the Askey-Wilson polynomials, with applications to coding theory*, SIAM J. Math. Anal. **18** (1987), 191–207.
5. T. Chihara, *An Introduction to Orthogonal Polynomials*, Gordon and Breach, New York, 1978.
6. L. Comtet, *Analyse Combinatoire, Tome Second*, Presses Universitaires de France, Paris, 1970.
7. P. Delsarte, *An algebraic approach to the association schemes of coding theory*, Philips Res. Repts. Supp. **10**.
8. ———, *Properties and applications of the recurrence $F(i+1, k+1, n+1) = q^{k+1}F(i, k+1, n) - q^kF(i, k, n)$* , SIAM J. Appl. Math. **31** (1976), 262–270.
9. P. Diaconis and R.L. Graham, *The Radon transform on Z_2^k* , Pac. J. Math. **118** (1985), 323–345.
10. R.J. Levit, *The zeros of the Hahn polynomials*, SIAM Review **9** (1967), 191–203.
11. F. MacWilliams and N.J.A. Sloane, *The Theory of Error Correcting Codes*, North Holland Mathematical Library, New York, 1977.
12. R.J. McEliece, E.R. Rodemich, H.C. Rumsey and L.R. Welch, *New upper bounds on the rate of a code via the Delsarte-MacWilliams inequalities*, IEEE Trans. Inf. Theory **It-23** (1977), 157–166.
13. D. Stanton, *Some q -Krawtchouk polynomials on Chevalley groups*, Amer. J. Math. **102** (1980), 625–662.
14. ———, *Orthogonal polynomials and Chevalley groups*, Special Functions: Group Theoretic Aspects and Applications, R. Askey et al. (eds.), Reidel, Dordrecht, 1984, pp. 87–128.
15. ———, *A partially ordered set and q -Krawtchouk polynomials*, J. Comb. Th. A. **30** (1981), 276–284.
16. G. Szegő, *Orthogonal Polynomials*, AMS Soc. Colloquium Pub. Vol. 23, Fourth Edition, Providence, 1975.